

Bibliography of Indonesian Geology

BIBLIOGRAPHY OF THE GEOLOGY OF INDONESIA AND SURROUNDING AREAS

Edition 8.0, February 2026

J.T. VAN GORSEL

X. PALEONTOLOGY, BIOSTRATIGRAPHY



www.vangorselslist.com

X. PALEONTOLOGY, BIOSTRATIGRAPHY, INDONESIA + SE ASIA

X. REFERENCES INDONESIA- SE ASIA: PALEONTOLOGY, BIOSTRATIGRAPHY 60
X.1. Quaternary/Recent faunas-microfloras and distribution, Biogeography 60
X.2. Cenozoic 136
X.3. Jurassic- Cretaceous 185
X.4. Triassic 197
X.5. Paleozoic 206
X.6. Quaternary Hominids and Mammals, Archeology 219

This chapter X of the Bibliography 8.0 contains 367 pages with 2618 papers. These are mainly papers of a more general or regional nature. There are numerous additional papers in this Bibliography with paleontological content, but if they are focused on with faunas/ floras from specific localities, they are listed under those areas .

It is organized in six sub-chapters:
- chapter X.1 on modern and sub-recent fauna and flora distribtions, information that serves as analog data for interpretations of depositional facies and paleoclimate from fossil assemblages;
- chapters X.2- X.5 on fossil faunas and floras, organized by age;
- chapter X.6 is a large final chapter on mammals and hominids, mainly of Quaternary age (1267 papers).

Four relatively recent introductory review papers on Indonesian fossils were published in 2014 in Berita Sedimentologi (FOSI/ IAGI): Cenozoic biostratigraphy (Van Gorsel, Lunt and Morley BS 29), Cenozoic macrofossils (Van Gorsel, BS 30), Paleozoic and Mesozoic faunas and floras (Van Gorsel, BS31 a, b). Most of this introduction is taken from these papers.

There is a vast amount of literature on the biostratigraphy and paleontology of Indonesia and surrounding Southeast Asia. A 2014 comprehensive reference list of 337 pages, selected from an earlier edition of this Bibliography and issued as a supplement of Berita Sedimentologi 29, contained over 3200 papers on Paleozoic- Recent biostratigraphy and paleontology of the region. (online at:
http://www.iagi.or.id/fosi/files/2014/04/BS29A_Bibliography_biostrat_hvg_final2.pdf)

Early papers on fossil groups (late 1800's- early 1900's but also later) tended to focus on fossil identifications and taxonomic descriptions. Many of these are still valuable today, especially since many of these studies have not been updated and many of the remote regions have not been revisited. Modern papers tend to deal more with faunal and floral successions (biostratigraphy), interpretation of paleoenvironments (paleoecology), paleoclimate, and geographic faunal/floral provinces (paleobiogeography).

A remarkable comprehensive listing of all fossil species known from Indonesia around 1930 was published by Escher et al. (1931, Professor Martin Memorial volume). This volume also includes reviews of stratigraphy of the Paleozoic by Brouwer, the Mesozoic by Wanner and Tertiary by Leupold and Van der Vlerk. A map of the more important fossil localities in the Indonesian region was compliled by Krijnen (1931), in the same volume.

Updated listings of fossil species reported from Indonesia include:
- foraminifera, by Harsono Pringgoprawiro, Kadar and Skwarko (1998; 3 volumes);
- mollusca, by Skwarko, Sufiati and. Limbong (1994; 3 volumes).

Biostratigraphy is still the main tool for age determinations of sedimentary rocks, and good age control is obviously critical for all geology interpretation. Paleobiogeographic patterns provide constraints on the reconstruction of the SE Asia mosaic of tectonic blocks and most papers on this are captured in Chapters I.1 - I.2).

Papers on the various fossil groups are listed in Tables 1-10. These are not complete listings of all relevant papers, and additional titles can be found scattered across the Bibliography 7.0..

X.1. Quaternary-Recent faunas-microfloras and modern distribution

This section X.1 of Bibliography 8.0 contains 76 pages with 646 papers, mainly on the taxonomy of modern or sub-recent microfaunas and microfloras as well on their distributions in Recent sediments of SE Asia. For a more extensive introduction see Van Gorsel, Lunt and Morley (2014, Bertia Sedimentologi 29).

An understanding of modern biofacies distributions is obviously significant for the interpretation of depositional environments and paleoclimates in the geological rock record. Different microfossil groups and species lived in different environments, and once the distributions of modern counterparts are understood, these can assist in the interpretation of ancient depositional environments, sequence stratigraphy, recognition of displaced faunal elements, etc.

Neogene benthic foram assemblages are sufficiently similar to modern faunas that direct comparisons may be made. Paleogene and older faunas consist mainly of extinct species, so interpretations rely more on comparisons with taxonomically related species or with similar morphotypes. Modern case studies using microfossil biofacies for Miocene paleoenvironmental interpretation in the region include Finger and Drugg (1992) in Central Sumatra, Simmons et al. (1999), Noad and Preece (2014) in North Borneo, etc..

Foraminifera

Foraminifera are the primary tool for facies analysis, particularly the benthic foraminifera. Today these are found from brackish coastal waters down to the deepest marine environments, and their environmental distributions are well documented in numerous studies. Foraminiferal assemblages contain information not only on paleo-water depth, but certain species may also be indicative of substrate, salinity, oxygen levels, paleotemperatures, etc. of sedimentary deposits.



Figure X.1. Recent reef flat miliolid foraminifera (Sorites, Peneroplis, planoconvex Cibicides) (Scholle 2003).

Early studies of modern foraminifera distribution off Java were by LeRoy (1939) and Myers (1945). Classic papers on the distribution of foraminifera in deep waters of East Indonesia include Hofker (1978; samples from the 1929 Snellius I expedition), Frerichs (1970; Andaman Sea), Van Marle (1988, 1991; Snellius II expedition) and Villain (1995; Mahakam Delta and Makassar Straits. Among more recent case studies on the distribution of foraminifera in the waters of the Indonesian archipelago are papers by Adisaputra, Barmawidjaya, Rositasari, Suhartati Natsir, etc.. For listings of papers on distributions of modern foraminifera in the waters of Indonesia and surrounding regions see Tables 1 and 2 and the Bibliography.

Major trends in distribution of modern foraminifera in Indonesia and nearby waters include:

1. increase in diversity from marginal marine to a maximum in deep shelf- upper slope environments,
2. increase in percentage of planktonic foraminifera with water depth from 0 in marginal marine to >80% of fauna in bathyal settings
3. decrease in calcareous species below ~2000-3000m where dissolution of calcareous tests starts to become significant (calcareous tests are sometimes subject to dissolution at much shallower depths adjacent to major deltas).

Table 1- FORAMINIFERA RECENT DISTRIBUTION

FAUNA	AREA	REFERENCES
Benthic foraminifera Monographs'	Challenger Exp.1873-1876	Brady 1884, Barker 1960 , Jones 1994
	Durrand Collection	Millett 1898-1904
	Albatross Exped. 1891	Cushman 1921
	Siboga Exped. 1899-1900	Hofker(1927, 1930, 1951
	Snellius Exped. 1929-30	Hofker 1978
	Snellius II Exp. 1984-85	Van Marle 1988-1991
	Recent Indonesia catalog	Adisaputra, Hendrizan & Kholoq 2010
Brackish, estuarine	SE Sulawesi	Horton et al. 2005
	N Borneo	Dhillon 1968, Bronnimann & Keij 1986
	Malay Peninsula	Mahani et al. 2010, 2011, Culver et al. 2012, 2013
Delta front, prodelta, bay	Mahakam Delta,	Villain 1995, Lambert 2003
	Java- north coast.	Rositasari 2010
	N Borneo	Ho Kiam Fui 1971
Carbonate environments	NW Java (incl. P. Seribu)	LeRoy 1938, Wijono 1991, Renema 2008, Suhartati 2009
	E Kalimantan	Renema 2006
	SW Sulawesi	Troelstra et al. 1996, Renema & Troelstra 2001, Renema et al. 2001
	Bali, Lombok	Barbin et al. 1987, Adisaputra 1998, 2000, Renema 2003
	PNG	Haig 1979-1993, Langer & Lipps 2003
	Great Barrier Reef	Jell et al. 1965, Renema et al. 2013
	NW Australia	Parker 2009, Parker & Gischler 2011
Shelf seas	Java Sea, Paternoster Platform	Myers 1945, Biswas 1976, Hofker 1978, Boichard et al. 1985, Isnaniawardhani 2009
	Sunda Shelf- S China Sea	Waller 1960, Szarek 2001, Szarek et al. 2006, Suhartati & Muchslin 2012
	Makassar Straits	Adisaputra & Rostyati 2003
	Arafura Sea	Suhartati & Rubiman 2010
Deep marine	East Indonesia	Hofker 1927, 1930, 1951, Van Marle 1988, 1991
	Makassar Straits	Rositasari 2010, Suleiman et al. 2011
	South China Sea-deep	Miao & Thunell 1993, 1996, Jian & Wang 1997, Marquez 1999, Szarek et al. 2009
	Sulu Sea	Rathburn et al. 1994, 1995, Szarek et al. 2007
	Andaman Sea	Frerichs 1970, 1971
	Timor Sea, Indian Ocean	Loeblich & Tappan 1994, Basov & Krashenninnikov 1995, Murgese & De Deckker 2005, De & Gupta 2010

Palynofacies

Palynomorphs can also be grouped into environmental categories, from terrestrial to marine environments, but these do not always not reflect local environment since all pollen and spores are designed to be transported water and wind, away from their source vegetation.

Pollen and spores can be divided into locally sourced and regionally sourced types, provided that the source plants are growing in the depositional locality. Interpretation of depositional environments is thus essentially an exercise in understanding pollen and spore transportation processes. They are, however, very helpful in augmenting environmental interpretations based on foraminifera. If foraminifera are absent, implying non-marine deposition, the locally derived pollen component may indicate the nature of the local depositional environment, but in open marine facies all the pollen is transported.

Pollen and spores are abundant in nearshore environments, but occur in low concentrations in holomarine sediments. However, in deep water turbidites, pollen and spores are as common as in coastal facies, reflecting downslope transport of sediments by gravity flow processes.

Also, it must be emphasized that vegetation is complex, and the same pollen genera may occur in unrelated environments. For instance, the genus *Casuarina* has species which are common along sandy coasts (*C. equisitifolia*) in kerapah swamps and kerangas (*C. nobilis*) and on mountains in Nusa Tenggara (*C junghuhniana*). The likely source taxon may therefore become clear by association: if it occurs with common *Dacrydium* pollen it is probably derived from kerapah/kerangas, if in sandy sediments with mangroves, probably from sandy coasts, and if from East Java, a montane source should be suspected.

Some common pollen and spore types are from some important vegetation types are as follows (dispersed pollen names in parentheses):

Mangrove belt:	*Rhizophora, Bruguiera (=Zonocostites ramonae), Avicennia, Sonneratia alba (=Florschuetzia meridionalis)*;
Back-mangrove:	*Acrostrichum, Sonneratia caseolaris (=Florschuetzia levipoli), Oncosperma, Nypa (= Spinizonocolpites echinatus), Acanthus;*
Freshwater peat swamp :	*Blumeodendron, Gonystulus, Cephalomappa, Lophopetalum, Alangium (= Lanagiopollis spp.);*
Seasonally inundated swamps:	*grasses, Barringtonia (= Marginopollis concinnus), Ceratopteris (= Magnastriatites grandiosus);*
Rain forests:	*Dipterocarpaceae, many Leguminosae, Rubiaceae and many others;*
Open savanna:	*Pinus, grasses.*
Mountains:	*Dacrycarpus imbricatus, Phyllocladus, Picea, Abies;*
Freshwater algae from lakes:	*Pediastrum, Botryococcus;*
Marine dinocysts:	*Spiniferites, Systematophora, Operculodinium.*

Table 2- OTHER MICROFAUNA/FLORA RECENT DISTRIBUTION		
FAUNA/ FLORA	**AREA**	**REFERENCES**
Ostracodes	Mahakam Delta	Carbonel & Moyes 1987, Carbonel and Hoibian 1988
	Sunda Shelf	Mostafawi 1992
	Java Sea, Bali	Dewi 1997, Mostafawi et al. 2005
	P. Seribu, Jakarta Bay	Watson 1988, Whatley & Watson 1988, Fauzielly et al 2013
	Malacca Straits	Whatley & Zhao 1987, 1988
	Other Indonesia	Muller 1906, Keij 1953, Dewi et al. (many papers)
Pollen, spores and kerogens	SE Asia case studies	Flenley 1973, Haseldonckx 1977, Gastaldo (op. div), Van der Kaars 1991, Chandra 2010, Sun et al. 1999
	N Borneo	Simmons et al. 1999
	E Kalimantan-Mahakam	Hardy & Wrenn 2009
	Sulawesi	Engelhardt et al. 2007
Diatoms (fresh-brackish water)	Lake Toba, Sumatra	Van der Marel 1947
	SE Sulawesi	Horton et al. 2007
	Malay Peninsula	Zong & Kamaludin 2004
Diatoms (deep marine)	Indonesia	Van Iperen et al. 1993, De Deckker & Gingele 2002
	South China sea	Wu et al. 2013
Calcareous nannofossils	SE Asia	Okada 1983, Varol 1985, Fernando et al. 2007
Radiolaria	Banda Sea	Harting 1863, Van de Paverd 1989
Dinoflagellate cysts	Java Sea, etc.	Lirdwitayaprasit 1997a,b, Matsuoka et al. 1999, Kawamura 2004, Furio et al. 2006, 2012, Hessler et al. 2013

X.2. Tertiary

Cenozoic sediments and volcanics cover about ¾ of the islands of Indonesia (Van Bemmelen, 1949), so it should not be surprising that the vast majority of paleontological studies has been on Cenozoic micro- and macrofossils. This Bibliography 8.0-section X.2 contains 48 pages with 445 papers on Cenozoic faunas and floras of Indonesia and surrounding regions. These tend to be papers of a more general nature; many additional titles on this topic may be found in the various area chapters if the fossils described were from a specific area only. Tables 3,4,5 and 6 list many of the key papers on the various fossil groups.

For more detailed overviews see the two relatively recent papers on Cenozoic fossils of Indonesia: Van Gorsel, Lunt and Morley (2014a; microfaunas/ biostratigraphy) and Van Gorsel (2014b, macrofossils).

Recent faunas from the Indonesian region boast some of the highest diversities in the world (a 'hotspot' region). This is probably also true for some Cenozoic faunas of Indonesia. In a series of monographs and papers between 1879 and 1937, professor Karl Martin of Leiden University and Museum ('Linnaeus of Java') identified 1412 species of Tertiary molluscs from Java alone! The collection of Tertiary corals from Indonesia at Naturalis Museum, Leiden, contains 271 species from 210 localities (Leloux and Renema, 2007).

Tertiary macrofossils in Indonesia are mainly molluscs and corals in marine deposits and plants, petrified wood and fresh-water molluscs in terrestrial deposits.

Today the principal tools for biostratigraphic subdivision of Tertiary sediments in Indonesia differ depending on paleo-environments:
- open marine sediments: planktonic foraminifera,calcareous nannofossils, radiolaria;
- very deep marine sediments, below carbonate compensation depth: radiolaria;
- shallow marine carbonate environments: larger foraminifera;
- shallow marine clastic marine environments: smaller benthic foraminifera, dinoflagellates;
- non-marine and marginal marine environments: palynology (spores and pollen).

Planktonic foraminifera

Planktonic foraminifera are a key group for biostratigraphy of Cretaceous- Recent open marine sediments. The series of evolutionary lineages of planktonic foraminifera (and the zonation schemes based on these), have been summarized in the classic books by Blow (1969, 1979), followed with minor modifications by Bolli and Saunders (1985). The zonation scheme of Blow, with the convenient 'N' (Neogene; 20) zones and 'P' (Paleogene 23 zones) zones has been used by most workers in the oil industry and academia in SE Asia.

The largely similar but 'competing' zonation of Berggren (1972, Wade et al., 2011), with its P (Paleocene), E (Eocene), O (Oligocene), M (Miocene) and Pl (Pliocene) zones, has not found many followers in SE Asia. This is partly because most of the modifications of the earlier Blow zonation proposed herein were more applicable to the Atlantic Ocean area than to the Indo-Pacific province.

The 'standard' zonations can be further refined when adding observations that reflect global or regional changes in climate or paleooceanographic patterns,such as changes in coiling directions of species of *Globorotalia, Neogloboquadrina, Pulleniatina*, etc. (Bolli & Saunders,1985), and climate-controlled events like incursions of cold water species.




Figure X.2. Example of Late Oligocene- Early Miocene planktonic foraminifera from Bulongan, NE Kalimantan (Koch 1926; Koch 1935).

While palynology is the tool of choice for non-marine deposits, it should be realized that spores-pollen are dispersed mainly in downstream direction from where they are produced. Therefore, mangrove pollen, on which much of the Neogene pollen zonation is based, are typically absent in fluvial-alluvial deposits away from the paleo-shoreline.

Smaller benthic foraminifera

Small benthic foraminifera are generally common in samples from all marine and marginal marine facies, and assemblages can be of very high diversity. About 2000 species and subspecies have been described, although many of these are probably synonyms (Lunt 2013). They are primarily used for paleoenvironment and paleobathymetric interpretations.

Classic early taxonomic descriptions of modern benthic foraminifera are the monographs on Recent material from the major oceanographic expeditions to the East Indonesia region, including the Challenger, Albatross, Siboga, Snellius I and Snellius II Expeditions (see Table 2).

Other significant papers include those by Millett (1898-1904) and Loeblich and Tappan (1994) on foraminifera from the Timor Sea. The principal taxonomic works on Neogene benthic foraminifera in the region are papers by LeRoy (1939-1944; Figure X.3), Belford (1966) and Whittaker and Hodgkinson (1979). For additional references see Table 3 and the Bibliography.




Figure X.3. Example of benthic foraminifera assemblage (Early- Middle Miocene Telisa Fm, Central Sumatra; LeRoy 1939).

Smaller benthic foraminifera species in SE Asia tend have relatively long age ranges and most of their age ranges have not been properly documented. The composition of benthic foraminifera assemblages has not changed much through the Neogene. In most instances highest and lowest occurrences of benthic species in a sedimentary section therefore reflect changes in facies, rather than evolutionary appearance or extinction levels. As a result, benthic foraminifera are of limited use for biostratigraphy age dating.

However, between about 1930 and 1960, the petroleum industry worldwide routinely used benthic foraminifera for correlations between exploration wells. In Indonesia BPM, Stanvac and Caltex all had laboratories with micropaleontologists that primarily used benthic foraminifera, and who tended to work in isolation. In the absence of detailed taxonomic studies, paleontologists generally identified species by genus name and numbers, such as *Rotalia* 5 or *Elphidium* 8. Due to the lack of communication between companies *Rotalia* 2 of BPM was not the same as *Rotalia* 2 of Stanvac.

Most of this early micropaleontological work was deemed proprietary and very little has ever been published. The taxonomic studies by LeRoy (1939 -1944) were an early effort to introduce formal species names for the Neogene foraminifera from the Caltex's oil wells of Sumatra, as well as outcrop sections of Java and East Kalimantan.

Over short distances, correlations based on 'tops' and 'bottoms' of benthic foraminifera were not unreasonable, but they were primarily correlations of facies changes or events like flooding surfaces, with no real age significance. Correlations across lateral facies changes were obviously very difficult. Such benthic foraminifera zonations are no longer used today.




Figure X.4. Recent 'prodelta' benthic foraminifera Pseudorotalia schroeteriana (top) and Asterorotalia trispinosa (bottom) (from Brady 1884, Challenger Foraminifera).

One of the exceptions to the above 'rule' is the use of the evolutionary developments in Late Miocene- Recent larger rotaliid foraminifera (Figure X.4). Rotaliids are common and dominant in delta front- prodelta facies, where planktonic foraminifera are absent or very rare. Following initial work by Huang (1964) in Taiwan, a zonation based on *Ammonia, Pseudorotalia* and *Asterorotalia* species was proposed by Billman and Witoelar (1974) in the East Kalimantan region, and subsequently expanded by Billman et al. (1980). It was further documented from the more marine deposits of the NE Java basin and calibrated to planktonic foram zones by Soeka et al. (1980) and Kadar (1992).

Table 3- CENOZOIC PLANKTONIC, SMALLER BENTHIC FORAMINIFERA		
FAUNA/FLORA	**AREA**	**REFERENCES**
General	Indonesia	Billman & Scrutton 1976, Van Gorsel 1988, Van Gorsel, Lunt and Morley 2014, Lunt 2013a,b
Neogene planktonic foraminifera	General	Blow 1969, 1979, Kennett & Srinivasan 1983, Bolli & Saunders 1984
	Java	Bolli 1966, Blow & Banner 1966, Van der Vlerk & Postuma 1967, Hartono 1960, 1969, Harsono P. 1968, 1983, Blow 1969, 1979, Wibisono 1972, Kadar 1972- 2008, Saint-Marc & Suminta 1979, Van Gorsel & Troelstra 1981, Harsono et al. 1998
	West Papua/ PNG	Belford 1962, 1966, 1967, 1984, 1988
	Sumatra	LeRoy 1948, 1952, Blow & Banner 1966, Kadar et al. 2008
	Andaman Sea	Frerichs 1971
Paleogene planktonic foraminifera	Kalimantan	Thalmann 1942, Baumann 1972
	Java	Hartono 1960, 1969, Purnamaningsih Siregar & Harsono P. 1981
	West Papua	Belford 1966, 1974, 1989
Smaller benthic foraminifera	Sumatra	LeRoy 1939, 1941, 1944
	Java	Koch 1923, Thalmann 1934, 1935, Yabe & Asano 1937, LeRoy 1941, 1944, Van der Sluis & Vletter 1942, Boomgaart 1949, Soeka et al. 1980, Sukandarrumidi 1990, Kadar 1992
	Kalimantan, Sabah	Koch 1926, LeRoy 1941, Mohler 1946, Billman and Witoelar 1974, Billman et al. 1980, Whittaker & Hodgkinson 1979
	Seram, Timor	Fischer 1921, 1927, Koch 1925, Rocha & Ubaldo 1964
	New Guinea	Belford 1962, Haig 1982

Larger foraminifera

By far the most publications on Tertiary microfaunas in Indonesia are dedicated to larger foraminifera. They are the dominant fossils in shallow marine carbonate deposits of Southeast Asia, especially of Eocene and Late Oligocene- Middle Miocene ages.They have been used in biostratigraphy in Indonesia for over a century, and are still the most useful group for age and environmental interpretations in carbonates, which generally contain very few planktonic foraminifera, nannofossils or palynomorphs.

Early classic works on larger foraminifera from Indonesia were by Verbeek, Rutten, Van der Vlerk, Tan Sin Hok, Caudri, Umbgrove, and others. The overall stratigraphic successions of genera and species and their evolutionary trends in most lineages had been identified by these workers by the late 1930's. More recent work on Indonesian larger foraminifera is by Boudagher-Fadel, Lunt, Renema and others. For selected references see Table 4 and the Bibliography.

Larger foraminifera are benthic foraminifera with relatively large and complex calcareous tests. They are generally 2-5mm in diameter; but some specimens of *Cycloclypeus, Eulepidina* and *Nummulites* may grow to 5 cm or more.

Most Recent larger foraminifera have a symbiotic relationship with algae or diatoms, which limits their presence to the photic zone (<100m water depth), in clear waters away from clastic influx. Examples of Recent larger foraminifera are shown in Figure X.5.




Figure X.5.. Modern tropical larger foraminifera: 1. Alveolina from Madang lagoon, PNG, 2. Baculogypsinoides? from Great Barrier Reef, Australia. 3. Marginopora from French Polynesia. 4. Foraminiferal beach sand from Kiribati (Western Pacific) (Langer 2008).




Figure X.6. Vertical sections of Cenozoic Indonesian larger foraminifera (Van der Vlerk & Dickerson, 1927).

Larger foraminifera comprise several taxonomically unrelated families. The complex inner structure generally requires study of oriented thin sections through the embryonic stage for accurate identication at the species level, or of the degree of development within an evolutionary series. However, identifications at genus level can generally be made from randomly oriented thin sections or from external features, using a hand lens. For examples of thin sections of Tertiary larger foraminifera from Indonesia see Figure X.6.

One of the earliest descriptions of larger foraminifera from SE Asia is on Late Eocene *Nummulites* limestones from SE Kalimantan, West Sumatra and Central Java (Verbeek (1871, 1892; Figure X.7).




Figure X.7. Middle Eocene Nummulites javanus (microspheric) from Nanggulan, Central Java (Verbeek and Fennema 1896).

Larger Foraminifera biozonations

The famous Eocene-Recent 'East Indies Letter Classification' is a zonation scheme (zones Ta-Th) based on larger foraminifera and in the youngest part originally also molluscs. It was first proposed by Van der Vlerk and Umbgrove (1927) (Figure X.8) It was further improved by Leupold and Van der Vlerk (1931) and Adams (1970).

Calibration of this larger foraminiferal zonation to the geological time scale has improved significantly over time, first in the 1970's-1980's through correlations of larger foram zones to planktonic foraminifera zones

(Clarke & Blow 1969, Haak and Postuma 1975, Adams 1984, Chaproniere 1984) and later through Strontium isotopes. For the latest ranges and calibration of the Letter Classification to the modern GPTS time scales see Lunt and Allen (2004), Lunt (2013) or Van Gorsel, Lunt and Morley (2014).




Figure X.8. Distribution chart of Tertiary larger foraminifera that define the 'East Indies Letter zonation' (Van der Vlerk, 1948).

Evolutionary series in larger foraminifera

Indonesia was the birthplace of classic studies on larger foraminifera evolution patterns of *Cycloclypeus*, *Miogypsina* and *Lepidocyclina* (Tan Sin Hok, 1932-1939), which were later refined by Van der Vlerk and others; Table 4).

Many larger foraminifera lineages started from a relativey simple, spirally arranged ancestor, then gradually evolved over millions of years into increasingly complex forms with concentric/ radial chamber growth patterns. These trends are well documented, and identification of evolutionary stages can be very helpful for age determinations.

Examples include (references see Table 4):
- *Heterostegina- Cycloclypeus* (Late Eocene- Recent)
- *Heterostegina- Tansinhokella- Spiroclypeus* (Late Oligocene- Early Miocene)
- *Neorotalia- Miogypsinoides- Miogypsina* (latest Oligocene- Middle Miocene) (Figure X.2.8)
- *Lepidocyclina* embryon evolution (Oligocene- Miocene) (Figure X.2.9).
- *Borelis- Flosculinella- Alveolinella* (Oligocene- Recent)




Figure X.9. Evolution from Late Oligocene Miogypsinoides complanata with long initial spiral on left (Van der Vlerk 1959) through Early Miocene Miogypsinoides dehaartii (Larat; Van der Vlerk 1924) to advanced Middle Miocene Miogypsina indonesiensis/antillea on right (Madura, Schipper and Drooger 1974).




Figure X.10 Evolution of Lepidocyclina embryonic chambers in Oligocene- Miocene Van der Vlerk (1959).

Table 4 - CENOZOIC LARGER FORAMINIFERA		
FAUNA/FLORA	**AREA**	**REFERENCES**
Larger foraminifera-	General	Douville 1905-1924, Mohler 1946, Adams 1965-1992, Lunt 2003, Lunt & Allen 2004, Renema 2002
	Letter zonation	Van der Vlerk & Umbgrove 1927, Leupold & Van der Vlerk 1931, Gerth 1935, Tan Sin Hok 1939, Van der Vlerk 1959, Clarke and Blow 1969, Adams 1970, Haak and Postuma 1975, Boudagher-Fadel & Banner 1999, Govindan 2003, Lunt and Allan 2004, Renema 2007, Matsumaru 2011
	Cycloclypeus	Tan Sin Hok 1932, MacGillavry 1962, Drooger 1955, Adams & Frame 1979
	Lepidocyclina	Van der Vlerk 1928, 1959-1973, Scheffen 1932, Tan Sin Hok 1936, Caudri 1939, Van der Vlerk & Postuma 1967, Ho Kiam Fui 1976, Van Vessem 1978, Adams 1987, Ghosh & Saraswati 2002
	Miogypsina	Tan Sin Hok 1936, 1937, Drooger 1953, 1963, 1993, Raju 1974, Schipper & Drooger 1974, De Bock 1976, Adisaputra-Sudinta et al. 1978
	Heterostegina.- Spiroclypeus	Van der Vlerk 1925, Tan Sin Hok 1930, 1937, Krijnen 1931, Lunt & Renema 2014
	Austrotrillina	Adams 1968
	Pellatispira	Umbgrove 1928
	Alveolinids	Bakx 1932, Mohler 1949, Ritsema 1951, Wonders & Adams 1991
	Nummulitids	Verbeek 1871, 1874, 1891, Doornink 1932, Renema et al. 2002
Larger foraminifera-faunal studies	Multiple areas	Rutten 1911-1934, Van der Vlerk 1922-1973, Adams 1965, 1992, Boudagher Fadel 1997-2008
	Sumatra	Verbeek 1871-1891, Brady 1878, Douville 1912, Tappenbeck 1936
	Java	Verbeek 1892, Martin 1881, Douville 1916, Hanzawa 1930, 1965, Caudri 1932, 1939, Doornink 1932, Mohler 1949, Cole 1975, Adisaputra et al. 1978, Adisaputra & Coleman 1983, Boudagher-Fadel 1997-2008, Renema 2002, 2007, Sharaf et al. 2006, Umiyatun et al. 2006, Irwansyah et al. 2011, Rahmawati et al. 2012, Lunt 2013
	Kalimantan-North Borneo	Verbeek 1871, Von Fritsch 1877, Provale 1908, Rutten 1911-1926, Van der Vlerk 1925, 1929, Tan Sin Hok 1930-1940, Adams 1965, Hashimoto et al. 1973, Hashimoto & Matsumaru 1981, Roohi (1998, Boudagher-Fadel et al. 2000, McMonagle et al. 2011, Cotton et al. 2014, Lunt 2014,
	Sulawesi	Crotty & Engelhardt 1993, Boudagher-Fadel 2002, Sudijono 2005
	West Papua-Papua New Guinea	Schlumberger 1894, Rutten 1914, 1921, 1925, 1936, Crespin 196), Binnekamp 1973, Bain & Binnekamp 1973, Beldord 1974, Brash et al. 1991, Sudijono 2000
	Sumba, Kei islands	Van der Vlerk 1922, 1966, Douville 1923, Caudri 1934, Bursch 1947
	Timor	Bakx 1932, Henrici 1934, Hayasaka & Ishizaki 1939, Ritsema 1951, Marks 1954

Calcareous nannoplankton

Calcareous nannofossils are minute calcite platelets that cover unicellular pelagic marine algae (coccolithophorids), and are common in low latitude open marine deposits. Because most of the taxonomic work on was done in the last thirty years, by relatively few workers, this group does not have the taxonomic confusion that existed among some of the foraminifera taxa.

Nannofossils have an advantage over planktonic foraminifera in that they may be found in larger numbers closer to shore, more specimens can be extracted from smaller samples (e.g. side-wall cores), and sample processing is relatively easy. They are relatively simple structures, so identification is relatively easy, although the often subtle differences between species do require some operator experience. On the negative side, due their small size they require high-powered optical microscopes and samples are prone to contamination, both from contaminated lab equipment and from geological reworking. The routine use of this group for age dating of oil well samples from Tertiary basins in Indonesia started in the mid-1970's.

Although some of the earliest pioneering work on nannofossils was done in Indonesia (Tan Sin Hok, 1927; Figure X.11), relatively little research work has been done on this group in Indonesia since then. For selected references see Table 5 and the Bibliography.




Figure X.11. Calcareous nannofossils of Discoaster group from the Plio-Pleistocene of Roti and Timor, from the pioneering study of Tan Sin Hok (1927).

Calcareous nannoplankton biozonations

The most widely used nannofossil zonation in the low-latitude Cenozoic is based on the work of Martini (1971). It has a resolution similar to that of planktonic foraminifera, and integration of both tools typically provide a higher stratigraphic resolution either of the methods alone.

The Martini zones are numbered in a similar fashion as used in planktonic foraminifera: Neogene zones NN 1-21 and Paleogene zones NP1-25. Another frequently used zonal scheme is that of Okada & Bukry (1980), using similar zones, but numbered with 'CN' and 'CP' prefixes. Varol (l983) proposed another slightly modified scheme for the Miocene- Recent in Southeast Asia..

In the petroleum basins of Western Indonesia open marine facies with diverse nannofossils can be rare. Prodelta shales may yield only impoverished, *Discoaster*-dominated nannofossil assemblages. For this reason Lambert and Laporte-Galaa (2005) proposed a modified Miocene nannofossil zonation for the Mahakam Delta area, based on species of the *Discoaster* group only.

Ostracoda

Ostracode fossils are the calcareous valves of a class of small crustaceans. Although found in a wide variety of marine and non-marine environments, ostracodes have been much less studied than foraminifera, and are not routinely used in SE Asia biostratigraphy. This is partly because they are usually associated with more numerous and apparently more useful foraminifera, partly because there are very few ostracode specialists. Stratigraphic ranges of ostracode species appears to be poorly documented and many of the Cenozoic species appear to be long-ranging (e.g. Guernet, 1993).

The first paper on fossil ostracodes in Indonesia is by Fyan (1916) from the Pliocene of Timor (Figure X.12). The most comprehensive study on Indonesian material is by Kingma (1948), who described 94 species of Late Cenozoic ostracodes from North Sumatra, the Bojonegoro 1 well and the Kendeng zone in East Java, as well as 20 Recent species from the Java Sea. Series of more recent, smaller papers by LeRoy (1939-1945) and Keij (1953-1979) are also of interest. A review of studies and extensive listings of recent and fossil ostracode species described from SE Asia is by Hanai et al. (1980).




Figure X.12. Pliocene ostracodes from Atambua, Timor (Fyan, 1916).

The outer shells of ostracod species may vary between relatively thin, smooth types or thicker and highly ornamented varieties. Both types occur in brackish and marine environments, but smooth types appear to be relatively more common in lower salinity settings.

For selected references on ostracods see Table 5 and the Bibliography. For papers on Recent species and distributions see Carbonel et al. (1987, 1988), Dewi (1993-2008), Mostafawi, Titterton et al. and others.

Diatoms

Diatoms are photosynthetic single-celled algae that build intricate siliceous tests. Some species are benthic forms, others led a planktonic existence, but, being dependent on light, all lived within the photic zone. They are present in a broad range of environments, from fresh water deposits to marginal marine and pelagic oceanic environments. Many taxa have near-global distributions across multiple climate zones.

Fossil diatom floras have been commonly used for biostratigraphic and paleoenvironmental interpretations in oceanic sections, like the Neogene of the Pacific Ocean (Burckle 1972, 1978), but there are very few studies from Indonesia. The first and still most elaborate paper on Cenozoic diatoms from Indonesia is Reinhold (1937), describing 208 species and varieties of marine and lacustrine diatoms from the Neogene of Java (Figure X.13). Other examples of applications of fossil diatom biostratigraphic zonation include papers by Ninkovich and Burckle (1979), Siregar (1981) and Burckle (1982), also on Late Neogene Central and East Java material.

Although diatoms are an important part of Recent marine plankton worldwide, they are generally rare in most samples of marine sediment ary rocks in Indonesia, because:

1. diatoms are more abundant in high latitudes; low latitude pelagic oozes are dominated by planktonic foraminifera and calcareous nannofossils, except in areas of diatom algal blooms, as in cold-water upwelling zones.
2. diatoms are relative susceptible to dissolution, both in the water column and under burial diagenesis.

Dissolution of the opaline diatom tests is widespread below burial depths of ~600-1000m (>40-50°C ; Opal-A to Opal CT transition), below which much of the diatom shells are converted to siliceous cement and are the main cause of a Bottom-Simulating Reflector (BSR) on seismic lines over deep water deposits. At greater depths all that remains of diatom tests is microcrystalline quartz in bedded cherts. Diatoms may still be seen occasionally in micropaleontology samples from greater depths as pyritized discs ('*Coscinodiscus*').

Accumulations of the large marine diatom *Ethmodiscus rex* have been documented in Pleistocene glacial intervals in deep marine cores in the Indian Ocean off Sumatra (DeDeckker & Gingele 2002) and further West (Broecker et al. 2000), and also in the Equatorial Pacific and Atlantic Oceans. These represent times of increased salinity and supply of silica to the oceanic surface water, under a more arid climate.

Quaternary lacustrine diatomites are known from the Lake Toba area in Sumatra (Samosir Island; Van der Marel 1947) and from Central Java (Darma area, Grandjean & Reinhold, 1933; Kalioso, Sangiran, Reinhold 1937) and East Java (Upper Kalibeng Fm near Kabuh, Krian in Kendeng zone). Such lacustrine diatom-rich deposits are commonly associated with volcanic tuffs that provided additional silica source.

The distribution of modern pelagic marine diatoms in East Indonesia deep water sediments was studied by Van Iperen et al. (1993). Horton et al. (2007) described modern diatom distribution in mangrove swamps of the Tukang Besi islands, SE Sulawesi.

Recent field experiments in Indonesia have shown the potential of producing biofuels from cultures of marine diatoms (Nurachman et al. 2012), which underlines the potential importance of diatoms for marine algal hydrocarbon source rocks.




Figure X.13. Diatoms from Late Miocene- Pleistocene of Central Java (Reinhold, 1937).

Radiolaria

Like diatoms, radiolara are not common in Cenozoic 'normal' marine' sedimentary rocks from Indonesia, and Cenozoic and Recent radiolaria have received very little attention here. Radiolaria appear to be relatively more common in marine sediments of Mesozoic age in the Indonesian- SE Asia region, and a larger body of work exists for these (Hinde 1902, 1908 and later work by Wakita et al., Jasin Basir, Munasri, etc.). Cenozoic radiolaria are most abundant in deep marine- oceanic deposits in areas of high productivity like upwelling zones.

Harting (1864) published some early descriptions of modern radiolaria species from the deep Banda Sea. More recently, modern radiolaria abundances on the Banda Sea floor were studied by Van de Paverd (1989; peak abundances between 2000-4000m).

Radiolaria are a useful group for Mesozoic- Cenozoic biostratigraphy, but mainly of deep marine deposits. The principal biozonations for Cenozoic radiolaria are by Riedel and Sanfilipo (1978), and Sanfilipo et al. (1985).

Bandung-based Indonesian paleontologist Tan Sin Hok did some of the earliest work on radiolarian faunas in the Timor- Roti islands. However, he did not realize that the samples he analyzed for his Ph.D. Thesis (Tan Sin Hok, 1927), were collected by Tan's thesis advisor Brouwer, were a mixture of Late Neogene and Late Jurassic- Early Cretaceous marls. Because of this, he had come to the erroneous conclusion that radiolaria were not suitable as index fossils, which is far from the truth.

Riedel (1953) described 26 Neogene radiolarian species from true 'post-orogenic' Pliocene-Pliocene marly sediments from Roti island (Brouwer Sample 178; Figure X.14), the presence of which Tan had mentioned, but not described in any detail.




Figure X.14. Plio-Pleistocene Radiolaria from near Bebalain Roti Island (Riedel 1953, Plate 85).

.

Cenozoic Palynology (Spores- Pollen)

Palynology is the study of microscopic organic-walled remains of plants and algae that are composed of organic material that does not dissolve in mineral acids. They are collectively are known as palynomorphs and, in addition to pollen and pteridophyte spores, also include afungal spores, algal remains (including dinoflagellate cysts), foraminiferal test linings, plant cuticle and chitinous remains of insect skeletons.

Palynomorphs are generally abundant in Cenozoic sediments of Indonesia, except in limestones and tuffs, and are usually the only microfossils present in non-marine sediments. The principal groups in the Tertiary SE Asia

which are useful in stratigraphy are spores and pollen grains from land plants (reflecting vegetation within a catchment), and also cysts of planktonic marine dinoflagellate algae, fresh-water algae, etc.




Figure X.15. Examples of Neogene pollen and spores from Brunei (a-i) and marine dinoflagellate (j), acritarch (k) and microforaminifer (l).

The flora of SE Asia is extremely diverse, with more than 30,000 higher plant species recorded. These leave a fossil record of about 600 or so pollen and spore types. In most instances, pollen types reflect taxonomic variation at the generic level, but in some plant genera, each species can be identified. This is the case in three genera, *Sonneratia*, (a mangrove), *Alangium* (a swamp forest tree) and *Stenochlaena*. Pollen of these three genera are all stratigraphically useful, with fossil pollen of Sonneratia being described as the pollen taxon *Florschuetzia, Alangium as Lanagiopollis*, and *Stenochlaena* as *Stenochlaeniidites*.

Palynomorphs generally withstand normal sediment diagenesis well, but become modified by katagenesis during deep burial to many thousands of feet (or equivalent heating), which will turn the palynomorphs black, This makes identification more difficult, but it also provides a tool to determine the degree of thermal alteration of a rock sample.

Palynology biozonation

Very few of the common Neogene pollen or spore types have well defined stratigraphic ranges. 'Traditional' palynological zones based on the ranges of pollen and spores, are therefore relatively long- ranging: there are only six or seven zones which can be interpreted from the Neogene (Figure X.2.16), where planktonic forams and nannofossils boast around 20 zones. However, quantitative analyses often allow higher resolution zonations at a basin scale, using abundance fluctuations that reflect semi-regional climate and sea level fluctuations.

The first palynology biozonation for the Tertiary of SE Asia was developed by Shell palynologists Germeraad, Hopping and Muller (1968) and reviewed by Morley (1977). Zonations were further clarified by Morley (1991, 2004 and others). Additional case studies were published by Rahardjo et al. (1994) for Java (Figure x23) and Lelono (2012).




Figure X.16. Eocene- Recent palynological zonation of Java (Rahardjo et al. 1994, in Lelono 2012).

Subdivision of the Oligo-Miocene is mainly based on evolutionary changes in *Florschuetzia*, the pollen of the mangrove genus *Sonneratia*, (Figure X.16, X.17). All of the pollen and spore types used for Neogene zonations reflect pollen variation at the species level. *Phyllocladus hypophyllus* pollen distinguishes the Pleistocene, *Dacrycarpidites australis* pollen first appears in the mid-Pliocene; *Stenochlaeniidites papuanus* pollen is restricted to the late Late Miocene and Pliocene, and species of *Florschuetzia* allow zonation of the Middle and Early Miocene. However, mangrove pollen tend to be absent in supratidal deposits, so the *Florschuetzia* zonation is of limited use in basins dominated by fluvial-alluvial sections, like the onshore Tertiary basins of Thailand (Watanasak, 1990).




Figure X.17. Evolution and stratigraphigraphic ranges of mangrove pollen genus Florschuetzia, leading to the modern Sonneratia (Germeraad et al. 1968 and Muller 1984, modified by Mao and Foong 2013).

Local higher resolution palynology schemes have been developed as a tool for correlations, using:
1. changes in relative abundances that reflect hinterland climate changes (fluctuations in % of rainforest versus grasses or montane palynomorphs);
2. identification of transgressions, which in fluvio-deltaic series are typically reflected by peaks of mangrove pollen (Morley, 1996).

Examples of this type of quantitative biostratigraphy include Yakzan et al. (1996, Malay Basin), Carter and Morley (1996, Early Miocene, Kutei Basin), Morley et al. (2003, Natuna basin), Morley et al. (2004, Makassar Straits) and Lelono and Morley (2011, Oligocene, East Java Sea).

Most Neogene quantitative palynological events reflect events relating to sequence stratigraphy. High resolution biostratigraphic frameworks involving palynology have usually been presented as sequence biostratigraphic studies (e.g. Morley et al., 2011, etc.).

Some of the long term changes in floral compositions reflect plate tectonic movements. Collisions like those between Indian and Eurasia in the Eocene and New Guinea with East Indonesia plates allowed migrations of India and Australian floral elements into SE Asia (Morley, 1998, 2000, Lelono 2007). The position of the Early-Middle Eocene boundary is based on the appearance of elements of the Proto-Indian flora (Morley 1998). The appearance of *Stenochlaena milnei* type and *Dacrycarpus imbricatus* is related to the collision of the Australian Plate with SE Asia.

The development of a spores-pollen zonation for the Paleogene of Indonesia is still work in progress.

Dinoflagellate cysts are often present in palynology preparations, mainly from marine sediments. They are the primary tool for high-resolution biostratigraphy zonation in mid- and high-latitude Cenozoic basins like the Paleogene of the North Sea, but they are relatively rare and poorly studied in the Cenozoic of SE Asia. A few preliminary case studies by Matsuoka (1981, 1983, 1984), Morgenroth (2000, 2008, 2011) and Besems (1993) do suggest potential utility.

Table 5- OTHER CENOZOIC MICROFAUNA/MICROFLORA

FAUNA/FLORA	AREA	REFERENCES
Calcareous nannofossils	Indonesia	Panuju 2010
	Kalimantan, N Borneo	Keij 1964, Lambert & Laporte-Galaa 2005, Panuju et al. 2007
	Java	Okada 1981, A. Kadar 1990,1991, Kapid et al. 1991-2003, Siesser et al. 1984, Theodoridis 1984, Nur Hasjim 1988, Rubiyanto & Harsolumakso 1996, Choiriah & Kapid 1999, Panuju & Kapid 2007
	Sulawesi	Kadar & Sudijono 1993, Priyadi & Sudijono 1993
	Roti	Tan Sin Hok 1927, Kamptner 1955, Jafar 1975
	Timor, Tanimbar	Hadiwisastra 1987, Hadiwisastra 1993
	Recent- Banda Sea	Kleine 1990
Ostracodes	multiple areas	Brady 1880, LeRoy 1939, 1941, Kingma 1948, Van den Bold 1950, Keij 1953-1979, Hanai et al. 1980, Keij 1964,
	Java	McKenzie & Sudijono 198x, Solihin et al. 2015,
	Timor	Fyan 1916
	E Kalimantan	Doeglas 1931
	South China Sea	Zhao 2005
Radiolaria	Roti –Mio-Pliocene	Tan Sin Hok 1927, Riedel 1953
	Sumatra forearc	Ling & Samuel 1998
	Celebes, Sulu Seas	Scherer 1991a,b
	Waigeo	Ling et al. 1991
Diatoms	Java	Grandjean & Reinhold 1933, Reinhold 1937, Ninkovich & Burckle 1979, P. Siregar 1981, Burckle 1982
	Buton	Reinhold 1952
	Indian Ocean	Jouse & Kazarina 1974, De Decker and Gingele 2002
Spores-Pollen palynology	Zonations	Germeraad et al. 1968, Morley 1977, 1991, 1996, Yakzan et al. 1996, Mao & Foong 2013
	Dispersal, climate	Morley 1998, 2000, 2003, 2007, 2011, 2012, Lelono 2007, 2012
	E Kalimantan, Makassar Straits	Barre de Cruz 1982, Carter & Morley 1995, Polhaupessy 1998, 2007, Morley et al. 2004, 2006, Morley & Morley 2010
	East Java Sea	Lelono & Morley 2011
	Java	Polhaupessy 1980, 2009, Takahashi 1982, Rahardjo et al. 1994, Van der Kaars & Dam 1995, 1997, Lelono 2000, 2001, 2007, 2012
	Natuna Basin	Morley et al. 2003, 2007)
	West Papua, PNG	Lelono 2008, Playford 1982
Dinoflagellate cysts	Java	Matsuoka 1981, 1983, 1984, Morgenroth et al. 2000, 2008, 2011
	NW Borneo	Besems 1993
	Sunda Shelf	Kawamura 2002

Cenozoic Macrofossils

Numerous papers have been published on Cenozoic macrofossils of Indonesia (Table 6). Much of this paleontological literature may be >100 years old, but since these old publications are mainly descriptions of fossils and fossil localities and often contain superb illustrations, they are still valuable today.

In Indonesia Cenozoic macrofossils are mainly represented by molluscs and corals in marine deposits and by plants, petrified wood and fresh-water molluscs in terrestrial deposits. Fossil groups that were important in pre-Cenozoic time like cephalopods and brachiopods are greatly diminished after the end-Cretaceous extinction event. For more detail see Van Gorsel (2014) 'An introduction to Cenozoic macrofossils of Indonesia' in Berita Sedimentologi 30.

A key publication on Indonesian fossil genera and species was the Professor Martin memorial volume, edited by Escher et al. (1931). It contains comprehensive listings of all Cenozoic fossil species described from Indonesia, including chapters on Cenozoic molluscs (Van der Vlerk), corals (Gerth), echinoids (Wanner) and plant fossils (Posthumus).

Although theoretically many macrofossil taxa have biostratigraphic value, they have lost their significance as biostratigraphic tools today, primarily because it is much easier to collect an adequate number of microfossils (even in small drill samples) than it is to collect macrofossil assemblages . As a result there are very few active experts in the science of macrofossils and our knowledge on stratigraphic ranges of macrofossil species and their calibrations to modern time scales tend to be remain poorly known.

Molluscs

Molluscs (bivalves and gastropods) are the most common macrofossils in the Cenozoic of Indonesia. Their diversity is overwhelmingly high with about 4000 known Recent species in Indonesia. They are found in depositional environments ranging from fresh water to deep marine. Biogeographically Cenozoic molluscs belong to the Indo-Pacific bioprovince, which is very different from the Mediterranean Province.

Most of the early paleontological studies were dedicated to molluscs and most of these were by Professor Karl Martin of Leiden University, who was nicknamed the 'Linnaeus of Java'. Martin mainly studied collections assembled by geologists/ mining engineers of the 'Dienst Mijnwezen of the East Indies'. He published numerous monographs and papers between 1879 and 1937, especially from Java, from where he identified a total of 1412 species of molluscs. From the Pliocene of North Sumatra 347 species were identified (Martin, 1928), and 232 species from the Pliocene of Timor (Tesch 1915, 1920).

Molluscs were the principal fossil group used for age dating of Cenozoic sediments, until microfossils became the preferred biostratigraphic tools. Martin developed a 'Lyell-type' method of relative age determination, based on the decreasing percentage of living species in increasingly older fossil populations. Percentages of living species by stage were (Van der Vlerk, 1931):

- Eocene: 0 %;
- Early Miocene: 8-20%;
- Late Miocene: 20-50%;
- Pliocene: 50-70%;
- Quaternary: >70%.

This Lyellian percentage method appeared to work reasonably well, but there were anomalies, especially in cases of deep marine mollusc assemblages. For instance, faunas from asphalt-bearing Sampalakosa beds of Buton were initially interpreted as Oligocene by Martin (1933, 1935), because none of the 35 mollusc species was known from the Recent. However, associated foraminifera and diatoms suggested a Late Miocene-Pliocene age, and the molluscs were subsequently recognized as deeper marine fauna, whose modern equivalents wee poorly known (Beets 1952).

Oostingh (1938) and Shuto (1978) started building more conventional mollusc biozonations of the Miocene-Pleistocene of Java, based on vertical ranges of species. Many of their zones are based on species of the gastropod genus *Turritella*. Shuto (1978) and Baggio and Sartori (1996) also proposed updates of the Lyellian percentages proposed by Martin. As noted above, neither this mollusc zonation, nor the percentage method are still used for dating Cenozoic sediments today.

Corals

Corals are common in most Oligocene to Recent limestones across Indonesia. The fossil coral collections at Naturalis Museum, Leiden, contain 271 species from 210 localities in SE Asia (mainly Indonesia; Leloux and Renema 2007). However, relatively little work has been done on their biostratigraphic zonations, biofacies and paleobiogeography and most of the taxonomic work is relatively old.

Modern coral reefs in Indonesia have been described fairly extensively, particularly in a series of studies by Umbgrove between 1928 and 1947. Umbgrove identified and described about 150 species of Recent corals and coral reefs around Java, Sumatra and Sulawesi. Key review papers on modern Indonesian coral reefs include Umbgrove (1946, 1947).

Gerth (1921) recognized the Neogene-Recent corals of Indonesia as typical Indo-Pacific assemblages. The Indo-Australian Archipelago today has the highest coral diversity in the world (Renema et al. 2008). Veron et al. (2015) identified the Birds Head- Sulu Sea region as the global center of peak coral diversity.

Most coral species have fairly narrow temperature ranges in which they thrive, and are therefore sensitive to climate changes. Both periods global cooling and extreme warming can negatively affect coral diversity and abundance. Cenozoic corals and coral reefs in Indonesia therefore show distinct periods of diversification and decline (Wilson 2002 and others).

Corals are generally rare in Eocene (too warm?) and Early Oligocene (too cold?) carbonates of SE Asia, which are dominated by larger forams and coralline algae (Wilson and Rosen 1998). The Late Oligocene- Early Miocene was a period of increased coral diversification and relatively wide distribution of coral reefal limestones (Wilson 2002, 2011, Johnson et al. 2011). This was then followed by several steps of declining diversity and carbonate abundance in Middle Miocene and later time.

An elegant review paper on Indonesian Cenozoic corals is Osberger (1956), which includes listings of principal deposits and range charts of species. Notable case studies on Cenozoic fossil corals from Indonesia include:
- Von Fritsch (1877): Eocene corals of SE Kalimantan
- Felix (1913-1921): Miocene-Pleistocene corals from Java, Kalimantan, Timor,
- Gerth (1923): 120 coral species from the Late Tertiary of East Kalimantan
- Gerth (1921, 1933): corals from Eocene- Miocene of Java;
- Umbgrove (1924-1950): corals from Miocene-Pleistocene of Kalimantan, Java, Sumatra, Buton, etc.
- Osberger (1954, 1955): Late Tertiary corals from Java.
- Santodomingo (2014, 2015, 2016): Miocene corals from patch reefs in East Kalimantan.

In an attempt to use corals for relative age dating Umbgrove (1946) and Osberger (1956) used the Lyellian method of increase in percentage of living coral species through time, similar to what Martin used for mollusks: Eocene-Oligocene 0%; Early Miocene, 6-9%; Middle Miocene 15-30%, Late Miocene-Pliocene 30-60% and Pleistocene ~80%.

As argued by Osberger (1956), whilst the overall trend is real, there is too much variability for this method to be reliable for age dating. Also, as with mollusks, these percentages did not hold up well when dealing with less well-known deeper marine assemblages. For instance, corals from the Lower Pleistocene Pucangan Formation of the Kendeng zone of East Java only had 50% known Recent species, which was explained by Umbgrove (1946) as due to the relatively deep water facies with common poorly known solitary species.

Plant fossils

Plant fossils are locally common in Eocene and Mio-Pliocene non-marine deposits of Western Indonesia. Fossilized wood and plant leaves may be found in two settings: (1) in claystones associated with coal beds, fossilized under poorly oxygenated swamp conditions, or (2) in tuff deposits, where entire forests were killed and preserved after major volcanic eruptions, and where silicified tree trunks may still be found in growth position. The latter are particularly common in various silica-rich Late Miocene-Pliocene volcanoclastic deposits of West Java and South Sumatra.

There is an urgent need to renew the study of fossil woods, since mining of Mio-Pliocene fossil wood for ornamental purposes is currently a major extractive industry in West Java, etc., This is a major opportunity to study the taxonomy and taphonomy of these ancient forests, but there is no evidence that this work is being done. These fossils need to be better understood before their localities are destroyed.

Early descriptions of plant fossils include:
- Mio-Pliocene floras from West Java collected by Junghuhn (Goppert, 1854; Figure X..18);
- Eocene plant fossils collected by Verbeek from the Ombilin Basin, C Sumatra (Heer 1874, 1879);
- Eocene plant fossils collected by Verbeek from SE Kalimantan (Geyler, 1877).

Significant papers on petrified wood from Java and Sumatra include (additional references in Table 6:
- papers by Musper (1938, 1939) on age and localities of fossil wood on Sumatra and West Java;
- monographs on Cenozoic woods from SE Asia by Kramer (1974a, b).

Table 6 - CENOZOIC MACROFOSSILS INDONESIA

FAUNA/FLORA	AREA	REFERENCES
General	Indonesia	Escher et al. 1931
Molluscs (bivalves and gastropods)	Indonesia	Van der Vlerk 1931, Beets 1950, Shuto 1977, 1978, Skwarko et al. 1994, Baggio & Sartori 1996, Beu 2005, Robba 2013
	Java	Jenkins 1864, Martin 1879-1932, Boettger 1883, Haanstra & Spiker 1932, Oostingh 1933-1941, Wanner & Hahn 1935, Pannekoek 1936, Schilder 1937, 1941, Van Regteren Altena 1938-1950, Shuto 1974, 1978, 1980, Premonowati 1990, Zacchello 1984, Premonowat 1990, Robba 1996, Scolari 1999, 2001, Bazzacco 2001, Piccoli & Premonowati 2001, Rolando 2001, Van den Hoek Ostende et al. 2002, Hasibuan 2004, Leloux and Wesselingh 2009
	Kalimantan, Borneo	Boettger 1875, Martin 1914, Beets 1941-1986, Cox 1948
	Sumatra	Woodward 1879, Boettger 1880, 1883, Icke & Martin 1907, Martin 1928, Oostingh 1941, Haanstra & Spiker 1932, Wissema 1947
	Sulawesi	Schepman 1907, Beets 1950
	Buton	Martin 1933, 1935, 1937, Beets 1952, Janssen 1999
	Seram, Timor	Fischer 1927, Tesch 1916, 1920, Robba et al. (1989
	West Papua	Beets 1986
Tertiary Corals	Indonesia	Osberger 1956, Leloux & Renema 2007, Wilson & Rosen 1998, Wilson 2002
	Java	Reuss 1867, Martin 1879, 1880, Felix 1913, Gerth 1921, 1933, Yabe & Eguchi 1941, Umbgrove 1945, 1946a,b, 1950, Osberger 1954, 1955, Premonowati 1990, 1996
	Kalimantan	Von Fritsch 1877, Felix 1921, Gerth 1923, Umbgrove 1929
	Sumatra	Gerth 1925, Umbgrove 1926
	Sulawesi	Von Kutassy 1934
	Buton	Umbgrove 1943
	Timor, Seram	Felix 1915, 1920, Umbgrove 1924
	Flores, Sumba	Umbgrove 1939, Umbgrove 1946
	New Guinea	Felix 1912, Gregory & Trench 1916, Yabe & Sugiyama 1942a,b
Calcareous Algae	Indonesia	Lignac-Grutterink 1943, Johnson & Ferris 1949, Ishijima et al. 1978
	N Borneo	Johnson 1966
Echinoids	Multi-region	Lambert & Jeannet 1935, Jeannet & Martin 1937
	Kalimantan, Java	Von Fritsch 1877, Gerth 1922
	Timor, Aru Islands	Gerth 1927, Currie 1924
Plants, Wood	Indonesia, SE Asia	Krausel 1925, Posthumus 1931, Kramer 1974, Bande & Prakash 1986, Van Konijnenburg et al. 2004
	Java	Goppert 1854, Crie 1888, Krausel 1923, 1926, Den Berger 1927, Musper 1938, 1939, Mandang et al. 1996, 2004, Srivastava & Kagemori 2001

	Kalimantan	Geyler 1875
	Sumatra	Heer 1874, 1879, Krausel 1922, 1929, Den Berger 1923, Musper 1938, 1939, Schweitzer 1958
Fish (pre-Pleistocene)	S Sulawesi	Brouwer & De Beaufort 1923, De Beaufort 1934
	C Sumatra-Eocene	Rutimeyer 1874, Gunther 1876, Von der Marck 1876, Sanders 1934, Musper 1936
	C Java	De Beaufort 1928, Vorstman 1929 (otoliths)
Vertebrates (pre-Pleistocene)	C Sumatra-	Eocene bird skeleton: Lambrecht 1931, Van Tets et al. 1989
	C Sumatra	Oligocene bird tracks: Zaim et al. 2011, Zonnneveld et al. 2011, 2012
	Java – U Miocene	Sea cow: Von Koenigswald 1952
	E Kalimantan	Eocene Anthracocerid and Suidae teeth: Stromer 1931
	Timor- Eocene	Anthracocere skull: Von Koenigswald 1967, Ducrocq 1996
	PNG Miocene	Turtle: Glaessner 1942, Sea cow: Fitzgerald at al. 2013




Figure X.18. Mio-Pliocene plant fossils from Java, collected by Junghuhn.

X.3. Jurassic- Cretaceous

A significant amount of literature has been published on Mesozoic faunas of Indonesia. This sub-chapter X.3 of Bibliography 8.0 contains 11 pages with 90 papers on Jurassic- Cretaceous faunas and floras of Indonesia and surrounding regions. These tend to be papers of a more general nature, and the majority of references on this topic is in the area chapters if the fossils described were from a specific area only. Most of the key papers on Jurassic and Cretaceous fossil groups papers were flagged in Tables 7 and 8. For a more extensive introduction to Jurassic- Cretaceous faunas and floras in Indonesia see Van Gorsel (2014).

Early reviews of Mesozoic geology and stratigraphy of Indonesia include Wanner (1925, 1931) and Umbgrove (1935, 1938). Mesozoic fossil localities on Sumatra were discussed by Tobler (1923) and and Fontaine and Gafoer (1989). Another useful collection of Pretertiary paleontologic studies in SE Asia is Fontaine (1990).

Other 'classic' reviews of Jurassic stratigraphy and key faunas of SE Asia include the papers by Fontaine et al. (1983), Sukamto and Westermann (1992) and Sato (1995).

Many of the paleontological papers from Indonesia are quite old (most of them from the early 1900's), but are still valuable. These include monographs on Jurassic- Cretaceous ammonites, belemnites, brachiopods, bivalve molluscs, etc. More recent work is mainly by Fauzie Hasibuan, Bandung, but mainly on fossils from Misool island.

The Jurassic in East Indonesia is dominated by open marine pelagic facies (Misool, Sula, East Sulawesi, Buton, Timor, etc.) or by marine continental margin deposits, locally rich in ammonites and belemnites (West Papua). In West Indonesia (Sumatra, SW Sarawak), Jurassic shallow marine carbonate and clastic facies are more common.

A characteristic deep marine Late Jurassic pelagic limestone facies with abundant radiolarians and calcispheres is known from Buton, East Sulawesi, Timor, Seram and Misool (Wanner 1940). In these areas the Jurassic is overlain by relatively thin Cretaceous pelagic limestones, often with radiolarian cherts and with common *Globotruncana* planktonic foraminifera in the upper parts.

As noted above, early work on Mesozoic radiolaria was by Tan Sin Hok (1927), from Roti island, although he assumed they were of Late Neogene age. Tan described 141 species and varieties (138 new) from four deep marine radiolarian-rich limestones from the Bebalain area, most of them of Late Jurassic- Early Cretaceous age. Subsequent authors recognized that most of these species were of Mesozoic age (Riedel 1952, 1953), and the samples must have come from the deep marine Jurassic-Cretaceous sediments that unconformably underlie the Plio-Pleistocene marls on Roti and Timor. Many of Tan Sin Hok's new species have since been reported from Late Jurassic-Early Cretaceous deep water sediments, such as the Argo Abyssal Plain (Baumgartner 1993), Tanimbar (Jasin and Haile 1996) and the SW Sulawesi Barru Complex (Munasri 2013).

For age dating of Mesozoic rocks microfossils tend to be more significant than macrofossils. The preferred microfossil groups are:

1. Radiolaria in deep marine deposits;
2. Conodonts in Triassic and older shallow marine carbonates;
3. Dinoflagellate cysts in Late Triassic- Early Cretaceous shallow marine continental margin clastics (widely used in oil exploration wells in NW Australia and New Guinea, but, little of this work has been published for the Indonesian region.
4. Planktonic foraminifera and clcareous nannofossils in Cretaceous and younger open marine deposits;
5. Spores-pollen in non-marine- marginal marine deposits.

Unlike NE Thailand, Jurassic-Cretaceous vertebrate faunas are extremely rare in Indonesia, and limited to marine fish and reptile *(Ichthyosaurus)* fossils.

Jurassic- Cretaceous reefal limestones

Unlike Late Triassic limestones, which are relatively widespread across the entire the Indonesian region, Jurassic and Cretaceous shallow marine reefal limestones are known mainly from the Sundaland region of western Indonesia (Sumatra, Kalimantan- Sarawak). With the exception of some Early Jurassic 'near-reefal' limestones on Timor and in the Kolonodale area of East Sulawesi, no Jurassic- Cretaceous shallow marine reefal limestones are known from the NW Australia- New Guinea domain (or terranes derived from it after Jurassic).

Late Jurassic reefal limestones are known from many localities in West and South Sumatra (all part of 'Woyla terranes'?) (Yabe 1946, Fontaine et al. 1983, Beauvais et al. 1988, 1989, Fontaine and Gafoer 1989). Many of these appear to contain the hydrozoan *Cladocoropsis miriabilis* and the foram *Pseudocyclammina lituus*. Another key Late Jurassic reefal limestone occurrence is the relatively thick Bau limestone of the West Sarawak- NW Kalimantan border area is (Wolfenden 1965, Beauvais and Fontaine 1990).

These, and similar Jurassic limestones in West Thailand and the Palawan Block (Philippines), are rich in coral and calcisponges and stromatolites, but also have a relatively high mud matrix content, so these are sometimes characterized as 'lime mud mounds' (Beauvais et al. 1985).

Cretaceous Orbitolina

Early to Mid-Cretaceous shallow marine limestones with the larger foram *Orbitolina* and rudist-type molluscs are generally interpreted as low-latitude Tethyan forms. They are also known from Sundaland, The Philippines and Japan, but not from Australia- New Guinea or Eastern Indonesia. In West Indonesia they appear to define a belt of 'Tethyan' low-latitude shallow marine deposits along the Early Cretaceous margin of Sundaland:

- numerous localities across Kalimantan and West Sarawak (Martin 1888, 1889, Hofker 1963, Hashimoto and Matsumaru 1977), also Sabah (Leong 1972);
- South Sumatra (Lampung and Gumai Mts= Woyla Terranes?) (Zwierzycki 1931, Musper 1934, Yabe 1946);
- Central Java Lok Ulo accretionary complex (Verbeek 1891, Harloff 1933);
- West Latimojong Mountains, SW Sulawesi (Brouwer 1934).




Figure X.19. The first illustration of Early Cretaceous larger foram Orbitolina from West Kalimantan by Von Fritsch (1879), originally described as Patellina scutum).

Table 7 - CRETACEOUS FOSSILS INDONESIA

FAUNA/FLORA	AREA	REFERENCES
Faunas- General	SE Asia	Hashimoto et al. 1975
	Sumatra	Beauvais et al. 1989
Upper Cretaceous *Pseudorbitoides, Omphalocyclus*	Birds Head	Visser and Hermes 1962
	Papua New Guinea	Crespin and Belford 1957, Glaessner 1962
	Kalimantan	Van Gorsel 1978
Upper Cretaceous planktonic foraminifera (*Globotruncana*)	Timor, Leti	Schubert 1915, De Roever 1940, Sartono 1975
	Sula islands	Kholiq et al. 2011
	Misool	Vogler 1941
	Sabah	Adams and Kirk 1962
	Central Java	Asikin et al. 1992
	W Kalimantan	Tan Sin Hok 1936
	SE Sulawesi, Buton	Van der Vlerk 1925, Koolhoven 1932, Keijzer 1945
	PNG	Owen 1973, Haig 1981
	Halmahera	Brouwer 1923
Calc. Nannoplankton	West Papua	Panuju et al. 2010
Molluscs	Sumatra	Baumberger 1923, 1925, Musper 1934
	Kalimantan	Martin 1889, Vogel 1904

	South Sulawesi	Hasibuan and Limbong 2009
	West Papua , PNG	Heinz 1928, Skwarko 1967, Skwarko et al. 1983
Early Cretaceous *Orbitolina*	Kalimantan	Von Fritsch 1879, Geinitz 1883, Martin 1888, 1889, Hofker 1966, Hashimoto and Matsumaru 1974, Djamal et al. 1995, Bassi et al. 2009
	West Sulawesi	Brouwer 1934
	W Sarawak	Hashimoto and Matsumaru 1977
	Sabah	Leong 1972
	Central Java	Verbeek 1891, Verbeek and Fennema 1896, Harloff 1929
	S Sumatra	Zwierzycki 1931, Musper 1934, Yabe 1946, Beauvais et al. 1989
Radiolaria	Sarawak, Sabah	Tan 1978, Jasin 1985-2000, Aitchison 1994, Asis and Jasin 2012, 2013, Jasin and Tongkul 2013
	C Kalimantan	Hinde 1900, Grunau 1965
	SE Kalimantan	Wakita et al. 1998
	SW Sulawesi	Wakita et al 1994, 2000, Munasri 1995, 2013
	South Central Java	Okamoto et al. 1994, Wakita et al. 1994
	Buton	Soeka 1991, Ling and Smith 1995
	Roti	Tan Sin Hok 1927, Riedel 1953
	Timor	Tan Sin Hok 1927, Clowes 1997, Munasri 1998, Munasri and Sashida 2018
	Ungar (Tanimbar)	Jasin and Haile 1996
	N Indian Ocean	Renz 1974, Riedel and Sanfilippo 1974, Baumgartner 1993
Lower Cretaceous calpionellids	Buton, Timor	Wanner 1940
	NW Australia	Brunnschweiler 1960
Ammonites	Sumatra	Baumberger 1925
	SW Sarawak	Ishibasi 1982
	W Kalimantan	Krause 1904, 1911, Von Koenigswald 1939
	PNG	Matsumoto and Skwarko 1991, 1993
Lower Cretaceous belemnites	West Papua	Challinor 1989
	NW Australia-SE Asia	Mutterlose 1992
Fish (shark teeth)	Timor	Weiler 1932, De Beaufort 1932
Rudists	NW Borneo	Lau 1973, Fontaine and Ho 1989, Skelton et al. 2011
	SE Kalimantan	Martin 1888
	Misool	Boehm 1924

Table 8- JURASSIC FOSSILS INDONESIA		
FAUNA/FLORA	**AREA**	**REFERENCES**
Jurassic Stratigraphy	Sumatra	Fontaine et al. 1983
	Indonesia	Wanner 1931, Sato 1975, Sukamto and Westermann 1992
U Jurassic calcispheres (*Pithonella*)	Seram, Buton, Misool, Timor	Brouwer 1919, Tan Sin Hok 1927, Wanner 1940, Vogler 1940, Bolli 1974
	Buton	Bothe 1927
	Roti	Tan Sin Hok 1927
Late Jurassic limestones with *Cladocoropsis, Pseudocyclammina*	NW Kalimantan/ Sarawak (Bau Lst)	Wilford and Kho 1965, Wolfenden 1965, Beauvais and Fontaine 1990
	Sumatra	Silvestri 1925, 1932, Yabe 1946, Hanzawa 1947, Bennett et al. 1981, Beauvais 1983, 1989, Bassoulet 1989
	NE Palawan	Fontaine et al. 1983, Bassoulet 1983
U Jurassic corals	Sarawak	Beauvais and Fontaine 1990
	Sumatra	Beauvais 1983, 1989
Late Jurassic shallow marine foraminifera	W Sarawak (Bau Lst)	Bayliss 1966
	Sumatra	Silvestri 1925, 1932, Yabe 1946, Bassoulet 1989,
Middle- Late Jurassic Belemnites	Misool	Stolley 1929, 1935, Challinor 1989, 1991
	Sula Islands	Boehm 1907, 1912, Kruizinga 1921, Stolley 1929, Challinor and Skwarko 1982, Challinor 1989, 1991
	Timor, Roti, Babar	Rothpletz 1892, Stolley 1929, Stevens 1964
	Yamdena	Stolley 1929
	W Papua/ PNG	Challinor 1990
	Central Sulawesi	Stolley 1943
	Indo-Pacific	Stevens 1965
E-M Jurassic bivalves	NW Kalimantan, Sarawak	Martin 1899, Vogel 1896, 1900, Newton 1897, 1903, Tamura and Hon 1977, Hayami 1984
	Timor	Krumbeck 1923
	Misool	Soergel 1913, Wandel 1934, Hasibuan 2004
	PNG	Skwarko 1973, Grant-Mackie et al. 2006
U Jurassic bivalves (*Malayomaorica, Inoceramus haasti*) ('anti-tropical')	Sula, Buru, Seram,	Krumbeck 1923, Wandel 1936
	Timor, Roti	Krumbeck 1922, 1923
	Misool	Krumbeck 1934
	E Sulawesi	Hasibuan and Kosworo 2008
	Papua New Guinea	Glaessner 1945, Edwards and Glaessner 1953
	Australia NW Shelf	Brunnschweiler 1960
M-U Jurassic ammonites	Sula Islands	Kruizinga 1926, Westermann and Callomon 1988
	W Papua- PNG	Boehm 1913, Gerth 1927, 1965, Schluter 1928, Westermann and Getty 1970, Westermann and Callomon 1988, Westermann 1995
	Timor, Roti	Boehm 1908
	Babar	Callomon and Rose 2000
	W Kalimantan	Schairer and Zeiss 1992
E Jurassic ammonites	Yamdena, E Sulawesi	Wanner and Jaworski 1931, Jaworski 1933
	Roti, Timor	Krumbeck 1922
	W Kalimantan	Krause 1911, Hirano et al. 1981
M-U Jurassic Radiolaria	Sumatra	McCarthy et al. 2001

	SE Kalimantan	Wakita et al. 1998
	W Sarawak	Jasin et al. 1996, Jasin and Said 1999
	East Sulawesi	Hojnos 1934
	Sula Islands	Pessagno and Meyerhoff Hull 2002
	Timor, Roti	Sashida et al. 1999, Haig and Bandini 2013
U Jurassic nannofossils	Sula islands	Panuju 2011
	Timor	Kristan-Tollmann 1988a,b
	PNG	Haig 1979
Jurassic- E Cretaceous Dinoflagellate zonations	NW Australia margin	Cookson and Eisenack 1958, 1960, 1974, Helby Morgan and Partridge 1987, 2004, Partridge 2006, Mantle 2009, Mantle and Riding 2010, Riding, Helby et al. 2012
	Papua New Guinea	Davey 1988, 1999, Welsh 1990
	Misool	Helby and Hasibuan 1988, Sarjeant et al. 1992
	Sula Islands	Lelono and Nugrahaningsih 2012
Spores-Pollen	NW Australia margin	Burger 1996
Jurassic Coccoliths	Timor	Kristan-Tollmann 1988
E Jurassic brachiopods	Seram	Wanner and Knipscheer 1951
***Lithiotis* Limestone**	Timor Fatu Limestone	Krumbeck 1923, Geyer 1977, Hayami 1984




Figure X.20. Early Jurassic ammonites. 1. Lytoceras rotticum, 2. Arietites (Euechioceras) wichmanni and 3. Aegoceras subtaylori 3 from Roti (Krumbeck 1922); 4. Arietites geometricus from Batu Baraketak mud volcano, Roti (Rothpletz 1892); 5. Aegoceras borneense from West Kalimantan (Krause 1911).

X.4. Triassic

This sub-chapter X.4 of Bibliography 8.0 contains 9 pages with 65 papers on Triassic faunas and floras of the Indonesia- SE Asia region. These tend to be papers of a more general nature, and many more titles on this topic are in the various area chapters if the fossils described are from a specific area only. Many of those papers were flagged in Table 9, which lists most of the key papers on the Triassic fossil groups. Much of this introduction is from Van Gorsel (2014).

A large amount of literature exists on Triassic faunas of Indonesia.. Early reviews of Triassic faunas and facies in the Indonesian region include Zwierzycki (1925), Wanner (1931) and Umbgrove (1935). A more recent, brief review of Triassic biostratigraphy and correlations of East Indonesia is by Hasibuan (2010).

Triassic sedimentary and igneous rocks are relatively widespread in Indonesia, both in East and West, especially Late Triassic rocks (Early Triassic is only known from Timor). They represent a wide variety of shallow and deep marine facies, in limestones, clastics and pelagic cherts. Triassic fossils have been described from Timor- Roti, Sumatra, W Kalimantan, E Sulawesi, Buton, Buru, Seram, Ambon, Misool, etc.

Most marine Triassic deposits of SE Asia- Indonesia appear to represent relatively low-latitude depositional settings in and around two branches of the Tethys Ocean (Paleo-Tethys, Mesotethys). Many of the authors of paleontological papers on Triassic faunas from East Indonesia since the early 1900's commented on the Alpine-Tethyan affinity of these Triassic faunas, with remarkable similarities in many of the species and rock facies between Indonesia and the eastern Alps, Himalayas, etc.

Triassic is a period characterized by major extinction events at both its base and at its top.

At many localities in Eastern Indonesia the Middle- Late Triassic is developed in a 'flysch-type' clastic facies, locally overlain by Norian-Rhaetian limestones (Timor, Savu/ Roti, Leti/ Babar, East Sulawesi, Seram, Ambon, Misool, Buru, Buton, etc.).

Triassic shallow marine bivalves

The composition of Triassic bivalve assemblages in the Indonesian region varies depending on facies and paleogeographic setting: (1) shallow marine '*Myophoria* faunas' and (2) deep marine, pelagic bivalve faunas characterized by *Daonella, Halobia or Monotis* (see below).

Shallow marine Middle- Late Triassic bivalve-dominated limestones and sandstones with common *Myophoria, Cardita, Gervillia, Costatoria, Paleocardita, Indopecten verbeeki, Pinna blanfordi, Krumbeckiella* = *Timoria timorensis* Krumbeck), etc., have been reported from both West Indonesia (Sumatra) and East Indonesia (Misool, Buru) (Figure X.21).

Among the richest assemblages are the Nucula marls from Misool (Jaworski 1915). These are generally viewed as 'Tethyan' in nature, but may be assumed to be part of the southern, Gondwana margin of the (Meso-?)Tethys in Triassic time. The bivalve-rich 'Padang Fauna' of West Sumatra, collected by Verbeek NE of Lake Singkarak, was initially described by Boettger (1881) as an Early Eocene age assemblage. It contains *Myophoria, Paleocardita globiformis, Pinna blanfordi* and *Pecten (Indopecten) verbeeki* and was re-described and re-interpreted by Krumbeck (1914) as Late Triassic in age, with strong affinities to Circum-Mediterranean Carnian faunas,

'*Myophoria* sandstone' is also known from various localities in the central belt of the Malay Peninsula and Singapore (authors in Table 1), in what is now called the Semantan Formation and which are viewed as marine deposits in a foreland basin tied to the closure of the Paleo-Tethys during final E-ward subduction of W Malaya lithosphere beneath Eastern Malaya (Ismail et al. 2007).




Figure X.21. Typical Late Triassic shallow marine bivalves from Sumatra and Misool. 1. Paleocardita globiformis, 2-3. Pecten (Indopecten) verbeeki and 4. Myophoria myophoria from West Sumatra (from Boettger 1881, Krumbeck 1914).. 5-6. Myophoria vestita from 'Nucula Marls' of Misool (Jaworski 1915).

Triassic (hemi-)pelagic bivalves

Middle and Late Triassic deep marine shales and pelagic deposits in the Tethyan and Circum-Pacific regions often contain beds with common flat, thin-walled bivalves (also called 'flat clams' or 'paper shells'; Figures X.22, X.23). Pelagic bivalve assemblages are dominated by *Claraia* in Early Triassic, *Daonella* in Middle Triassic (Early Carnian), *Halobia* in Late Triassic (Carnian- Norian) and *Monotis* in latest Triassic (M Norian-Rhaetian; McRoberts 2010). Where found, they are often abundant and rock-forming. They are important biostratigraphic index fossils.

In Indonesia Triassic pelagic bivalves are found in East Indonesia, in the deep marine deposits of Timor (Krumbeck 1924), Roti, Buton (Sikumbang et al. 1995), Seram (Wanner 1907), Buru, Misool, Babar, etc. In West Indonesia they are known from Sumatra. They are not known from Australia, but this could be due to a scarcity of open marine facies of that age.



Figure X.22 .Late Triassic hemi-pelagic bivalves Monotis salinaria from Seram (Wanner, 1907)




Figure X.23. Middle Triassic pelagic thin-shelled molluscs Daonella indica from the Baung area, Timor (Krumbeck 1924). Massive hemi-pelagic shell accumulations like these are common in Triassic deep marine deposits across the (Meso-)Tethys Ocean.

These pseudo-pelagic, thin-shelled bivalves may have an 'anti-tropical' geographic distribution, like morphologically and ecologically comparable bivalves in the Late Jurassic (*Buchia, Aucella, Malayomaorica*) and Cretaceous (*Inoceramus*). Some provinciality between species has been suggested for Late Triassic *Monotis* (Westermann 1973, Thenius 1980, Silberling 1985), with Tethyan Norian assemblages characterized by *Monotis salinaria*.

Triassic ammonoids

Early, Middle and Late Triassic ammonoid assemblages are the most diverse of all ammonoid assemblages in Indonesia, and most of the species are from thin but complete condensed pelagic cephalopod limestones of Timor.

They were described in voluminous monographs by Welter (1922, 1915 and 1914 respectively) and Diener (1923). The Triassic 'Cephalopod Limestone' of Timor is a condensed, pelagic facies commonly called 'Hallstatt-type', and is the only place in Indonesia with a complete marine Triassic succession. Unfortunately the formation is mainly known from isolated blocks in in melange formation, but at Kapan, West Timor, Wanner (1913) observed the transition between Permian crinoid limestone into Early Triassic cephalopod limestone. Restored thickness suggests the entire Triassic in cephalopod limestone facies is very thin (<10m). Many of the fossils in this limestone are coated with a thin manganese layer, reflecting long periods of non-deposition, on a deep sea floor. Timor is the only place in Indonesia with known Early and Middle Triassic ammonoids.

In the Early Triassic ammonoid faunas had to recover from the mass extinction at the end of the Permian, after which new, diverse ammonoid assemblages developed rather rapidly. In Indonesia Early Triassic ammonoid faunas are known only from the cephalopod facies of Timor, from where Welter (1922) described 71 species of genera *Meekoceras, Flemingites*, etc.

Middle Triassic ammonoids from Timor were first documented by Welter (1915). As noted by Welter (1922) Early Triassic ammonoid assemblages from West Timor share many similarities with Himalayan-Mediterranean Triassic faunas.

Late Triassic ammonoid faunas of Timor are also extremely rich and diverse and are characterized a.o. by haloritids *Halorites* and *Juvavites, Anatomites, Amarassites* and many others (Figure X.24). Von Arthaber (1926) distinguished 110 species in the Carnian-Norian. Tatzreiter (1981) counted 90 species of trachyostracous ammonoids (not including the more numerous leiostracous ammonoids) in a 1m thick block of condensed Middle Norian limestone at Baun. Wanner (1931) reported 462 species from a 2m thick block of Carnian Cephalopod Limestone at Bihati.




Figure X.24. Example of some Late Triassic ammonoids from Timor. 1. Juvavites sarasinii from Bihati (Diener 1923), 2. Neotibetites (Krumbeck 1913), 3. Cladiscites crassestriatus from Bihati (Von Arthaber 1927), 4. Agathiceras.

Triassic brachiopods

Late Triassic brachiopod assemblages in East Indonesia often contain the rhynchonellid brachiopod species *Misolia misolia* (Figure X.25). The species is typical of Carnian-Norian (and younger?) shelfal marine deposits of the Gondwana margin of the southern/eastern Mesotethys (Dagys, 1993), and has been reported from Oman through the Spiti area of the Himalayas to East Indonesia and the NW Australian margin. In eastern Indonesia it is present in Misool, Timor, East Sulawesi, Buru, Seram, Ambon and Buton (references in Table 9 and the Bibliography).




Figure X.25. Triassic rhynchonellid brachiopod Misolia misolia from Buru (1) and from 'Athyrid Limestone' of Misool (Von Seidlitz, 1913).

Triassic corals

Following the major end-Permian extinction event, corals are relatively rare globally through much of the Triassic. The Late Triassic is a time in which coral reef limestones become common again along the Tethys margins, although calcareous sponges and algae remain important components of these reefal limestones.

In Indonesia Late Triassic corals are known from 'Fau Limestone' of Timor (Vinassa de Regny 1915), the Manusela Limestone of Seram (Wanner 1907, Wilckens 1937) and from Bangka (De Neve and De Roever 1947). Characteristic genera include *Thecosmilia* spp., *Isastraea*, *Retiophyllia* and *Montlivaltia* (Figure X.25). Many of the species from Timor are the same as those from the Austrian Alps (Vinassa de Regny 1915).

Late Triassic hydrozoan or stromatoporoid Lovcenipora

Upper Triassic reefal limestones in SE Asia often contain a hydrozoan or stromotoporoid named *Lovcenipora vinassai* (Seram, Timor; Wanner 1952; Figure 6). This is a Tethyan taxon, common in the Late Triassic of the Mediterranean. It was erroneously equated with a somewhat similar-looking Late Jurassic Tethyan hydrozoan *Cladacoropsis mirabilis* (Renz 1926), but these are separate taxa (Yabe 1946). It had led workers like Van der Sluis (1949) to erroneously assign a Jurassic age to the Manusela Limestone of Seram, an opinion followed by Van Bemmelen (1949), but fiercely protested by Wanner et al. (1952), but is still somewhat entrenched in Seram geological literature (Kemp 1992, etc.; see also Charlton and Van Gorsel (2014).

In Indonesia Triassic *Lovcenipora* is known from Seram, Buru (Gerth 1910) and from the Fatu Limestones of Timor (Vinassa de Regny 1915, Pia 1924). *Lovcenipora* has also been described from Jurassic- Cretaceous of Sumatra, but in most cases these are misidentifications and should be assigned to *Cladocoropsis miriabilis* (e.g. Yabe 1946).

Heterastridium

Hetrastridium is a remarkable Late Triassic hydrozoan fossil that is widespread in the Norian of the Tethys realm. It is a globular fossil, with a diameter typically 1-5cm and is the only hydrozoan colony with a probably (hemi-) pelagic lifestyle. It is locally abundant in the Upper Norian of the Hallstatt limestone in the Carnian Alps. From here it is distributed through the Tethyan belt towards East Indonesia, The Philippines and New Caledonia (Campbell 1974) to the Panthalassan terranes in Japan, New Zealand and North America.

In Indonesia *Heterastridium conglobatum* is locally common on Timor (Gerth 1915, 1942; Figure X.26.). It is found mainly in the thin, condensed pelagic 'Cephalopod Limestone', which is mainly composed of ammonites, and which is commonly compared to the 'Hallstatt-facies' of the Northern Calcareous Alps. Other occurrences include East Seram (Gerth 1909, 1942) and Misool (Krumbeck 1913).




Figure X.26. Late Triassic pelagic hydrozoan Heterastridium conglobatum from Timor. 1. From Bihati (size ~3.2cm); 2-3 From Oe Batok, Amarassi (size ~2 cm) (Gerth, 1915, 1942).

.

Late Triassic Krusin/ Tonkin floras

A characteristic Late Triassic floral assemblage is present in Indochina and South China, termed the *Dictyophyllum-Chlathropteris* floristic province and is viewed as a Late Carnian-Norian coastal floristic assemblage. Representatives of this flora are also found in SW Sarawak ('Krusin Flora'; Kon'no 1968, Vozenin-Serra 1983, Kimura 1984), in dredge samples from Reed Bank in the South China Sea (Kudrass et al. 1986) and probably also on Bintan Island, Riau (Jongmans 1951). The floral similarities suggest NW Borneo was closely connected to Indochina in Late Triassic time.

Table 9- TRIASSIC FOSSILS INDONESIA

FAUNA/FLORA	AREA	REFERENCES
Trias faunas, biostratigraphy	General	Wanner 1907, 1931, Zwierzycki 1925, Hasibuan 2008, Charlton et al. 2009 (Timor)
	Timor	Charlton et al. 2009
Late Trias Bintan Flora	Riau Archipelago,	Jongmans 1951, Wade-Murphy et al. 2008
Late Trias Tonkin/ Krusin Flora	N Vietnam, Thailand	Kon'no and Asama 1973, Vozenin-Serra and Franschesci 1999
	Reed Bank, S China Sea	Kudrass et al. 1986
	SW Sarawak	Kon'no 1968, 1972, Kimura 1984
Late Trias Palynofloras	Seram	Martini et al. 2004
Dinoflagellates	Seram	Helby et al. 1987, Martini et al. 2004
Late Trias corals	Buru	Gerth 1910
	Timor	Vinassa de Regny 1915, Roniewicz et al. 2005
	Seram	Wilckens 1937
***Heterastridium* (hydrozoan)**	Timor, Seram	Gerth 1909, 1915, 1927
	Misool	Krumbeck 1913
Late Trias brachiopods (incl. Rhaetian *Misolia*)	Misool	Von Seidlitz 1913, Hasibuan 1990, 2012, MacFarlan et al. 2011
	Seram	Deninger 1918, Krumbeck 1922, Wanner 1923, 1952
	Buru	Von Seidlitz 1913, Krumbeck 1913
	Ambon	Jaworski 1927
	East Sulawesi	Von Loczy 1934, Von Kutassy 1934
	Buton	Hasibuan 2010
	Timor	Krumbeck 1922, 1924, Grunau 1957

	NW Australia margin	Campbell 1994
M-L Trias pelagic bivalves *(Daonella, Halobia, Monotis)*	Thailand	Kobayashi and Tokuyama 1959
	W Kalimantan, Sarawak	Vogel 1904, Tamura and Hon 1977, Silberling 1985
	Seram	Krumbeck 1922
	Timor, Roti	Rothpletz 1892, Wanner 1907, Krumbeck 1924, Kutassy 1931, Ichikawa 1958, Gruber in Kristan-Tollmann 1987
	Sumatra	Volz 1899, Krumbeck 1914
	PNG	Skwarko 1967, 1973, Skwarko and Kummel 1974
M-L Trias shallow marine bivalves (*Myophoria* assemblages)	W Sumatra	Boettger 1981, Krumbeck 1914, Musper 1930
	Buru	Krumbeck 1913
	Misool	Jaworski 1915
	Papua New Guinea	Skwarko 1963, 1973
	Malay Peninsula, Singapore	Newton 1903, 1925, Cox 1936, Tokuyama 1961, Kobayashi and Tamura 1968, Tamura 1970, 1973
Gastropods	Timor	Tichy 1979
Lovcenipora	Seram, Burur	Wanner 1907, 1952, Gerth 1910, Pia 1924
	Timor	Vinassa de Regny 1915, Krumbeck 1921, 1924?
	Sumatra	Vinassa de Regny 1915
U Trias foraminifera	Timor	Kristan-Tollmann 1988
	Sumatra	Vachard 1989
	PNG	Kristan-Tollmann 1986, 1990
U Trias belemnites	Timor	Wanner 1911, Von Bulow 1915, Gheyselinck 1934
U Trias ammonites	Timor	Wanner 1911, Welter 1914, 1915, Diener 1923, Kieslinger 1924, Von Arthaber 1926, Tatzreiter 1980,1981, 1983
	Buru, Seram	Krumbeck 1913, Wanner 1928
E-M Trias ammonites	Timor	Welter 1915, 1922, Kummel 1968, Nakazawa and Bando 1968, Brayard et al. 2009
Trias conodonts	Sumatra	Metcalfe et al. 1979, 1989
	Timor	Nogami 1968, Koike 1984, Berry et al. 1984, Nicoll and Foster 1998
M-U Trias radiolaria	Timor	Hinde 1908, Rose 1994, Sashida et al. 1996, 1999
	South Sumatra (Garba)	Putra and Munasri 2016
U Trias dasyclad algae	Seram, Buru	Pia 1924
Norian-Rhaetian reefal limestones	SE Asia	Vachard and Fontaine 1988
	Sumatra	Metcalfe et al. 1979, 1989
	Seram, Buru	Wanner 1907, Martini et al. 2004, Charlton and Van Gorsel 2014
	East Sulawesi	Martini et al. 1997
	Banda Sea (Sinta Ridge)	Villeneuve et al. 1993, 1994
	Timor	Welter 1914, 1915, Haig et al. 2007
	Bangka	De Neve and De Roever 1948
	Papua New Guinea	Skwarko et al. 1976, Kristan-Tollman 1986, 1990

X.5. Paleozoic

This sub-chapter X.5 of Bibliography 8.0 contains 13 pages with 105 papers on Paleozoic faunas and floras of the Indonesian region, as well as related forms from mainland SE Asia. These tend to be papers of a more general nature, and many more titles on this topic are in the various area chapters if the fossils described were from a specific area only. Many of those papers were flagged in Tables 10-12, which list most of the key papers on Paleozoic fossils.

A fair body of literature exists on Paleozoic faunas of Indonesia, but much of it is in older and hard-to-find books and papers, and many of the papers are not in English or Indonesian. An early review of Paleozoic stratigraphy is by Brouwer (1931). Much of this introduction is from Van Gorsel (2014) 'Introduction to Paleozoic fossils' in Berita Sedimentologi 31.

In the Indonesian region the most complete Paleozoic sedimentary section is in West Papua, South of the Central Range, where older parts of the Australian continental margin sequence are exposed. The oldest fossils are Ordovician-Silurian age corals and graptolites.

In West Indonesia the only Early Paleozoic fossils are the enigmatic occurrence of a Devonian coral and stromatoporoid in limestone blocks in a melange section of uncertain age in NE Kalimantan. Relatively complete Paleozoic successions are known from the 'Sibumasu terrane' along the West part of the Malay Peninsula (Langkawi Islands, etc.), Peninsular Thailand, NW Thailand and East Myanmar, but somehow pre-Permo-Carboniferous rocks have not been reported from Sumatra, much of which is supposed to part of the Sibumasu terrane.

Late Paleozoic (Permian) faunas and floras are more widespread than Early Paleozoic in the Indonesian region. Well-known localities include West Papua (Visser and Hermes 1962), West Sumatra (Fontaine and Gafoer 1989), NW Kalimantan- SW Sarawak (Terbat Limestone; Fontaine 1990, Vachard 1990) and Timor (numerous papers).




Figure X.27.. Distribution of Paleozoic rocks/ fossils in Indonesia, showing the main Paleozoic outcrop areas on North and West Sumatra, NW Borneo, Timor and West Papua. (Umbgrove 1938).

Early Paleozoic (Ordovician- Devonian)

Early Paleozoic deposits are very rare in Indonesia, where the oldest fossils are Ordovician-Silurian nautiloids, conodonts and graptolites from West Papua. The oldest fossils from Western Indonesia are a Devonian *Heliolites* coral and a stromatoporoid from limestone blocks in a melange deposit in NE Kalimantan (Rutten 1940, 1947, Sugiaman and Andria 1999).

A Late Devonian brachiopod and a Permian ammonite were reported from the Kalosi region of C Sulawesi by Brouwer (1919), but their localities were never confirmed. They were believed to probably come from a local Chinese pharmacy by Von Koeningswald (1933).

The oldest fossils described from Indonesian territory are from the Ordovician-Silurian of West Papua. Cambrian and Late Precambrian sediments are probably present in this as well, but no diagnostic fossils have yet been recovered. Studies of Paleozoic fossils from West Papua are few, probably partly because faunas are not abundant and partly because outcrops of Early Paleozoic are in areas with difficult physical and political access. Most of the fossils described are from float samples from rivers draining the southern slopes of the Central Range.

Ordovician fossils reported from West Papua include:

1. Conodonts from 'basement limestone' in oil exploration wells Noordwest 1 and Cross Catalina 1 in the Central Range, including Ordovician *Serratognathus bilobatus* (Nicoll 2006). These limestones are part of the extensive Middle Cambrian- Early Ordovician Goulburn Group of carbonate-dominated shelf sediments, which underlie most of the Arafura Sea and West Papua South of the Central Ranges (Zhen et al. 2012);
2. Llanvirnian graptolites from shale from the Heluk River in the eastern foothills of the Central Range (Fortey and Cocks 1986; not described or illustrated);
3. Possible occurrences of Ordovician-age orthoconic nautiloids of the *Orthoceras*-group, described as *Irianoceras antiquum* by Kobayashi and Burton (1971), but this was deemed to be a junior synonym of *Bactroceras latisiphonatum* Glenister 1952 by Crick and Quarles van Ufford (1995). These nautiloids are from black shale nodules in river float within and south of the Central Range of West Papua. The problem is that (1) the nodules look very similar to those from Lower Kembelangan Formation black shales, which yielded common Middle-Late Jurassic ammonites, and (2) the fossils appear to have been collected in areas with nearby outcrops of Jurassic rocks, but with no evidence of any Paleozoic sediments. These observations suggest a more likely Jurassic age for these nautiloids, but this type of straight nautiloids is not known from post-Triassic rocks anywhere in the world. It is hard to decide whether these 'Ordovician' nautiloids represent (1) material from as yet unidentified outcrops of Ordovician shales in the Central Range; (2) an as yet undescribed nautiloid species of Jurassic age, or (3) reworked Ordovician fauna into Middle-Late Jurassic sediments.
4. Another occurrence of molds of possible Ordovician *Orthoceras* is in phyllitic shale (presumably Kemum Formation), just N of the mouth of the Wesan River in the NW part of the Birds Head (Kruizinga 1957).



Figure X.28. The characteristic Silurian tabulate 'chain coral' Halysites wallichi from a tributariy of the Noordoost/ Lorentz River, West Papua (Musper 1938).

Silurian-age fossils are known only from West Papua:

1. Graptolites *Monograptus turriculatus* and *M. marri* from the highly-deformed deep water sediments of the Kemum Formation in the north-central Birds Head (Llandoverian; Visser and Hermes 1962);
2. Small trilobites and brachiopods from float samples in rivers draining the southern slopes of the Central Range (Martin, 1911), associated with Silurian conodonts (Ludlowian; Van den Boogaard 1990);
3. Conodonts from Modio Dolomite in Charles Louis Range, SW West Papua, with *Panderodus* cf. *simplex*, indicate a Silurian age (Nicoll and Bladon 1991);
3. Silurian cosmopolitan coral *Halysites wallichi* was also found in river float in a tributary of the Noordoost/Lorentz River (Musper, 1938; Figure Figure X.28);
4. Late Silurian (M Ludlow) thelodont and acanthodian fish scales from Lorenz River in eastern W Papua and Kemum Fm of north part of Birds Head (Turner et al. 1995).

Devonian

Devonian fossils are relatively rare in Indonesia, and are known only from West Papua and from blocks in melange in NE Kalimantan.

Middle or Late Devonian corals, including *Heliolites* and *Favosites*, and stromatoporoids, have been reported from the dark grey 'Modio Dolomite Formation', which outcrops south of the Central Range of West Papua (Gerth 1927, Keijzer 1941, Oliver et al. 1995). These carbonates may be remnants of a widespread Middle Devonian reef system that continues for about 2000 km along the East Australia and New Guinea margin (Copper and Scotese 2003, Torsvik and Cocks 2013). Pebbles of M-U Devonian sandstones with the brachiopod genus *Spirifer* were reported from the same region by Teichert (1928).

In NE Kalimantan Devonian corals (*Heliolites*) and the stromatoporoid *Clathrodictyon* cf. *spatiosum* are present in limestone blocks in the 'Danau Formation' melange complex at the Telen River (Rutten 1940, 1947). *Heliolites* is a genus that is geographically widespread, also known from Indochina, NE Thailand, Laos, East Australia and Europe. Age of the melange complex has not been properly documented, but is likely of Early Cretaceous age (Tate 1992).

TABLE 10 - ORDOVICIAN- DEVONIAN

FAUNA/FLORA	AREA	REFERENCES
Devonian corals	West Papua	Gerth 1927, Keijzer 1941, Oliver et al. 1995
	NE Kalimantan	Rutten 1940, 1947, Sugiaman and Andria 1999
Devonian brachiopods	West Papua	Stehn 1927, Feuilleteau de Bruyn 1921
Late Silurian-Devonian fish	West Papua	Turner et al. 1995
	SE Asia	Wang et al. 2010
Silurian corals	W Papua- S of C Range	Gerth 1927, Teichert 1928, Musper 1938
Silurian conodonts	W Papua- S of C Range	Van den Boogaard 1990, Nicoll and Bladon 1991
Ordovician-Silurian graptolites	W Papua, Birds Head	Visser and Hermes 1962 (Silurian)
	W Papua, Heluk River	Fortey and Cocks 1986 (M Ordovician)
Ordovian(?) orthoconic nautiloids	W Papua Star Mountains	Kobayashi. and Burton 1971
	W Papua Central Range	Crick and Quarles van Ufford 1995
	Birds Head	Kruizinga 1957
E Ordovician conodonts	W Papua, S Central Range	Nicoll 2002, 2006

Carboniferous

Carboniferous sediments and fossils are comparatively rare in the Indonesian region, and are limited to North Sumatra, West-Central Sumatra, West Papua and possibly also NW Kalimantan. In the late 1800's many of the dark-colored Permian limestones from Sumatra were erroneously assigned to the Carboniferous, due to their superficial resemblance to the Visean 'Kohlenkalk' of NW Europe.

Sumatra

Early Carboniferous sediments are the oldest sediments identified in Sumatra and may be from two different tectonic terranes (Fontaine and Gafoer, 1989, Barber et al. 2005):

- temperate late Visean Alas Fm limestones in North Sumatra, age-datd with conodonts (Metcalfe 1983). These are probably part of the Sibumasu Terrane, which at this time was still part of the Australian margin.
- shallower marine and warmer-climate Kuantan Fm limestone with corals (*Syringopora, Siphonodendron*), calcareous algae (*Koninckopora*) and cosmopolitan foraminiferal assemblages from West Sumatra (Agam River, NE of Padang; Fontaine and Gafoer, 1989, Kato et al. 1999). This is part of the West Sumatra Block, with likely affinities to the low-latitude Indochina Block.

The unfossiliferous pebbly mudstones of the Bohorok Formation of West and North Sumatra are probably glacial deposits from the widespread Late Carboniferous- earliest Permian glaciation of Gondwana, but no fossils have been reported (similar rocks from Langkawi, NW Malaysia, do have cold-water brachiopod, etc. faunas).

NW Kalimantan- West Sarawak

In NW Borneo, in the border area between West Sarawak and NW Kalimantan, the oldest fossil-bearing rocks are tightly folded, steeply dipping sediments with chert and grey limestones of the Terbat Formation. These contain diverse latest Carboniferous and earliest Permian fusulinid assemblages with *Pseudoschwagerina, Paraschwagerina*, etc. (Krekeler 1932, 1933, Cummings 1962, Sanderson 1966, Vachard 1990, etc.). Correlative deposits are present in NW Kalimantan (Zeijlmans van Emmichoven, 1939). The fusulinid assemblages suggest affinity with low-latitude Cathaysian regions, not with the Sibumasu terrane.

West Papua

Conodonts from the Aimau Fm in the SW Tamrau Mountains of the Birds Head contain conodonts typical of Late Carboniferous (*Hindeodus minutus, Neognathus*; Nicoll and Bladon 1991).

TABLE 11 - CARBONIFEROUS

FAUNA/FLORA	AREA	REFERENCES
Corals	W Sumatra	Fontaine 1983, 1989
	West Papua	Kato et al. 1999
	Thailand	Fontaine et al. 1991
Foraminifera	W Sumatra	Metcalfe 1983, 1989, Vachard 1989
	NW Borneo	Cummings 1962, Sanderson 1966, Vachard 1990
Conodonts	Sumatra	Metcalfe 1983, 1986
	West Papua	Nicoll and Bladon 1991

Permian

Relatively rich Permian faunas and floras are known from many localities in mainland SE Asia, Indonesia and West Papua. In Indonesia Permian faunas and floras are common on Timor, West-Central Sumatra and West Papua (not in Papua New Guinea) and, to a lesser degree, from Borneo. For reviews of the shallow marine and non-marine Permian faunas and floras of SE Asia see Fontaine (1986, 2002). Key references are listed in Table 12 and in the Bibliography. Most of this introduction is from Van Gorsel (2014).

For biostratigraphic correlations of marine sequences brachiopods and mollusks have been the main tool in the Gondwana realm, while fusulinid foraminifera are the principal group used for correlations along the lower latitude Tethyan margin.

The Late Paleozoic faunas from Sumatra are generally comparable with those described from Thailand and Peninsular Malaysia, while the Permian of West Papua is Gondwanan.

The classic, highly diverse marine Permian faunas from Timor are famous for being the richest marine Permian faunas in the world, with over 600 species described by 1926 (Wanner 1926). These have been the subject of numerous papers in the early 1900's. A comprehensive recent review of Permian marine faunas of Timor is by Charlton et al. (2002). Unfortunately, most of the Permian fossiliferous sediments on Timor are not in any stratigraphic order, but occur as blocks in melange, olistostrome or broken formations. It is generally accepted that material from the famous fossil localities of Somohole and Bitauni areas are older (E Permian, ~Sakmarian- Artinskian) than those from the Basleo and Amarassi areas (late Middle Permian, ~Capitanian)

Early Permian cold-climate bivalves and brachiopods

Early Permian sediments associated with glacial marine deposits across northern Gondwana (Australia, India, etc.) often contain thick-shelled bivalves of the genera *Atomodesma* (Figure X.29) and *Eurydesma* and the cool-climate brachiopod *Globiella foordi* (now also called *Cimmeriella foordi*). Comparable bivalve assemblages may be present in the Early Permian of the Sibumasu- Cimmerian terranes now in Sumatra, NW Malaysia, W Thailand and SW China (Sun 1993).

In Indonesia assemblages with these genera were found in the Early Permian Maubisse Formation of Timor (Beyrich 1865, Rothpletz 1892, Wanner 1922, 1926, 1940, Belford 1960, Bird et al. 1989, Hasibuan 1994), but they are associated with relatively diverse marine faunas and glacio-marine deposits are not known from Timor. These faunas may suggest a proximity to glacial Gondwana of this part of Timor in earliest Permian time, but are not necessarily part of the glaciated terranes.

The presence in Timor Leste of a diverse fusulinid assemblage interpreted as of latest Carboniferous- earliest Permian age and reflecting a relatively warm climate (Davydov et al. 2013) is puzzling in the context of widespread glaciations on Gondwana at this time.




Figure X.29.. Permian bivalve Atomodesma exarata from the Kupang area, West Timor (Beyrich 1865).

Permian corals

Permian corals are relatively widespread in SE Asia They generally occur in carbonate lithologies and may be associated with fusulinid larger foraminifera,. Assemblage compositions differ with age, water depth and with paleogeographic position:

- Early Permian limestones from the Indochina terrane (East Thailand, etc.) contain typical 'Cathaysian', tropical, high-diversity coral and fusulinid assemblages, dominated by compound corals;
- Early Permian deposits of the Sibumasu Terrane have either no corals or small, solitary rugose corals, reflecting cooler and/or deeper waters (e.g. Peninsular Thailand; Fontaine et al. 1994, Yunnan, SW China; Wang and Sugiyama 2002). The low diversity assemblages of solitary rugose coral species, have been called '*Lytvolasma* faunas' or '*Cyathaxonia* faunas', and are generally viewed as 'anti-tropical', cooler climate coral assemblages (Kossovaya 2009).
- by late Middle and Late Permian time the Sibumasu terranes had moved towards lower latitudes and started to have similar high-diversity coral and fusulinid faunas as the Indochina terranes.




Figure X.30. Examples of Permian solitary and compound corals from Timor (Gerth 1921) (Carcinophyllum, Dibunophyllum (Verbeekiella), Pterophyllum, Favosites and Michelinia)

In Indonesia Permian coral faunas are known mainly from.

1. Timor Permian corals are locally abundant in the Maubisse Formation/ Basleo beds. They are mainly '*Cythaxonia*-faunas' with solitary corals like *Lytvolasma, Timorphyllum, Lophophyllidium, Verbeekiella* (Figure X.30), *Zaphrentis, Amplexus* and *Wannerophyllum*. Colonial rugose corals like *Michelinia, Favosites, Lonsdaleia timorica* and *L. molengraaffi* are present as well, but are relatively rare (Gerth 1921, Koker 1924, Wang 1947, Von Schouppe and Stacul 1955). The Timor Permian coral assemblages are very similar to those reported from the Baoshan Block, SW China (Zhao and Zhou 1987);
2. Sumatra. Permian corals have been reported from several localities in West Sumatra. Some of the Middle Permian limestones from West Sumatra contain high diversity corals that look similar to 'Cathaysian' assemblages of Central Thailand (Guguk Bulat; Fontaine 1983,1989);
3. West Papua. Permian corals are widely distributed in the Aifam Fm (Visser and Hermes 1962, p. 54), including solitary *Amplexus* on the Birds Head (Broili 1924). However, typical low-latitude compound corals appear to be absent here (Fontaine et al. 1994, p. 39).

Gerth (1926) already noted that the Permian coral fauna of Timor indicated a relatively warm paleoclimate, while Permian deposits on adjacent Australia contained glacial deposits, suggesting that Timor and Australia must have been farther apart in Permian time. However, if the Permian corals on Timor are younger than the earliest Permian glacial deposits on Gondwana, which they probably are, the contrast may not be as significant.

Permian ammonoids

Permian ammonoids are generally rare in Indonesia/ SE Asia, but the ammonoid assemblages of Timor are among the richest in the world (Smith, 1927, Wanner 1932). Wanner (1926) counted 37 species of ammonoids and 21 nautiloids. Most numerous genera are *Agathiceras* and *Paralegoceras* (= *Metalegoceras*; Figure X.31). Another Permian ammonoid locality in Indonesia isa rare *Agathiceras* from the folded series of Belitung (Kruizinga, 1950).

Blendinger et al. (1992) noted the remarkable similarity between the Middle Permian ammonoids from the cephalopod limestone of Timor with those from the West Mediterranean (Sosio Limestone of Siclily) and Oman, suggesting unrestricted faunal exchange in a Middle Permian Paleotethyan seaway. Ehiro (1997, 1998) classified the Middle Permian ammonoid faunas from 'allochthous Timor' in his' Equatorial Tethyan province', based on the presence of taxa like *Timorites* and *Waagenoceras*, which are not known from Australia.



Figure X.31. One of the most common Permian ammonoids from West Timor (Bitauni): Metalegoceras sundaicum (formerly Paralegoceras; Haniel 1915).

Permian Trilobites

Trilobites are rare in Indonesia, but have been reported from Permian sediments of three areas: Sumatra (Roemer 1880), Timor (Tesch 1923, Gheyselinck, 1937) and float in the Noord River in West Papua (Martin 1911).

They are mainly of the genus *Pseudophillipsia: P. timorensis* Roemer from Basleo, West Timor and *P. sumatrensis* from the Padang Highlands of West Sumatra (Figure X.32). Leman and Sone (2002) described similar *Pseudophillipsia* from the early Capitanian (Middle Permian) from Pahang, in the Central Belt of the Malay Peninsula (= west margin of East Malaya/ Indochina terrane).




Figure X.32. Permian trilobites. 1. Phillipsia sumatrensis from Permian of West Sumatra (Roemer 1880); 2 and 3. Neoproetus indicus from Bitauni, West Timor (Tesch 1923).

Permian Fusulinid foraminifera

Fusulinid larger foraminifera are tropical-subtropical shallow marine carbonate taxa (estimated paleolatitude range between 0 and 40° N and S), with a reputation of being excellent guide fossils in Carboniferous-Permian time. Fusulinids are widespread in Permian shallow marine limestones across SE Asia and areas further west, generally on terranes that border the Paleotethys suture. Hundreds of papers have been written on this group in SE Asia. For more details see references in Table 12 and the Bibliography.

Interpretation of fusulinid foram faunas can be very difficult. The taxonomy is overwhelming, with over 100 genus names and 1000's of species names. Fusulinids reached a maximum in diversity and sizes in the Middle Permian, as did other reefal fauna (corals, large molluscs, etc.). A significant extinction event of large fusulinids took place at the end of the Middle Permian (end or late Capitanian; e.g. Hada et al. 2014). The Late Permian is characterized by fusulinid assemblages that are reduced in size and diversity. Fusulinids went completely extinct at the mass extinction event at the end of the Permian.

In Indonesia Permian fusulinid foraminifera have been reported from 6-7 main areas, mainly on Sumatra, NW Borneo, Timor and the Birds Head of West Papua:

1. NW Kalimantan- Sarawak border area. Late Carboniferous- earliest Permian 'Terbat Limestone' of the NW Kalimantan- Sarawak border area were first reported by Krekeler (1932, 1933), and by several generations of subsequent authors (Table 12). Fusulinid assemblages are quite diverse and similar to 'Tethyan' faunas from Eastern Thailand and South China (Cummings 1962, Vachard 1990, Fontaine 1990, Sakamoto and Ishibashi, 2002). Tan Sin Hok (in Krekeler 1933) examined the fusulinid beds from Sadong valley and believed them to be same species (and in same volcanoclastic facies) as the Early Permian assemblages of Jambi. The age and nature of the Terbat Limestone assemblages suggest affinities to the Indochina Block, not Gondwana or Sibumasu (as do associated Triassic-Jurassic faunas and floras).

2. Padang Highlands, West Sumatra (Figure X.33). Middle Permian fusulinids have long been known from West Sumatra, mainly from the famous Guguk Bulat locality. Several of the large Middle Permian fusulinid index species of the Tethyan province were first described from Sumatra, like *Verbeekina verbeeki* (Geinitz, 1876), *Sumatrina annae* (Volz, 1904) and *Schwagerina padangensis* (Lange, 1925). Tien (1988) also recorded *Colania douvillei*. Many of the fusulinid species described from this part of West Sumatra are also common on the 'Cathaysian' Indochina Block of NE Thailand, but some have also been reported from the Sibumasu terrane, which by the end of the Middle Permian had moved into lower latitudes (e.g. Ueno et al. 2003).
3. Jambi, SW Sumatra. Early Permian fusulinids from the thin '*Productus* Limestone' horizon in the Mengkareng Formation at Telok Gedang along the Merangin River, Jambi, underlying the beds with the famous 'Cathaysian' Jambi Flora (Ozawa 1929, Thompson 1936, Vachard 1990, Ueno et al. 2007). Most abundant is *Pseudofusulina rutschi* (= *Rugofusulina rutschi*; similar to a more widely known species *Rugofusulina alpina* Schellwien). Also present is *Pseudoschwagerina meranginensis* (assigned to *Sphaeroschwagerina* by Davydov et al. 2013). It is a low-diversity assemblage, believed to be of Early Permian age, but more precise age interpretations vary from Upper Asselian (Tien 1989, Vachard 1990) to 'most likely Sakmarian' (Ueno 2007). None of the species described can be tied directly to assemblages elsewhere, so conclusions on precise age and paleobiogeographic affinity lack a real firm basis. Additional fusulinid occurrences are the Artinskian- Kungurian assemblage from nearby Batu Impi (Tien (1989, Ueno et al. 2007). Two additional small occurrences on Sumatra worth noting are in West Sumatra (Batang Siputar; Hahn and Weber 1981) and in South Sumatra (Bukit Pendopo; Palembang; De Neve 1949).




Figure X.33. Middle Permian fusulinid foraminifera from West Sumatra. 1. Fusulina granum-avenae (Verbeek 1896). 2. Sumatrina annae from NE of Lake Singkarak (Volz, 1904). 3. Verbeekina verbeeki from Guguk Bulat, Padang Highlands (originally described as Fusulina princeps by Brady, 1875).

4. Timor and adjacent islands Leti and Roti. Fusulinids are also known from various localities on Timor, and adjacent Roti and Leti islands (Schubert 1915a, Thompson 1949, Davydov et al. 2013). Many of the Timor assemblages are of low-diversity, but high abundance, and are dominated by a species initially described as

Fusulina wanneri by Schubert (1915), the type species of the 'anti-tropical' genus *Monodiexodina*. The small fauna of verbeekinids described from Leti Island by Schubert (1915b) with *Doliolina lepida* var. *lettensis* differs from assemblages known from Timor (Thompson 1949). Another different latest Carboniferous-earliest Permian assemblage from Timor Leste was described recently by Davydov et al. (2013).

5. Birds Head of West Papua. Rare fusulinids have been reported from West Papua. These the only fusulinid occurrences on undisputed Permian Gondwanaland, but are poorly documented. One occurrence in the Birds Head was figured by Visser and Hermes (1962, p. 54). Another possible fusulinid occurrence was reported, but not figured, from Permian limestone in a consultant biostratigraphy report of oil exploration well TBF 1X (3947m; NE of Misool in Bintuni Bay, south of Birds Head).
6. Bangka- Belitung. Less-known and rare fusulinids were reported from the intensely folded Permian beds of North Bangka (De Roever 1951) and Belitung (Strimple and Yancey 1974).

Permian brachiopods

In Indonesia Permian brachiopods are known from Sumatra, Timor and West Papua (Table 12). The principal monographs on Indonesian brachiopods are by Broili (1915, 1916), Hamlet (1928) and Wanner and Sieverts (1935), all from Timor. Permian brachiopods were described from Sumatra by Meyer (1922) and West Papua by Archbold (1981).

Productus and *Spirifer* groups dominate the Timor and West Papua assemblages (Figure X.34). The brachiopod faunas from Timor are relatively rich (49 species). However, unlike many other fossil groups from Timor like crinoids and blastoids, no new species were identified in the first monograph on this group by Broili (1916), attesting to the relatively cosmopolitan nature of these brachiopod taxa (mainly of Boreal and Palaeoequatorial genera; Crippa et al. 2014). Studies on paleobiogeographic patterns in Permian brachiopod assemblages therefore appear to have been somewhat non-diagnostic.




Figure X.34. Permian brachiopods. 1. Spirifer from Timor; 2. Stereochia (Productus) semireticulatus from Timor and 3. same species from Sibelabu, SE of Padang, West Sumatra.

The genus *Stereochia* (Figure X.5.7, 2,3 is of interest because it is commonly regarded as an anti-tropical genus (Shi et al. 1995, Crippa et al. 2014). In mainland SE Asia *Stereochia-Meekella* brachiopod fauna characterizes the Sibumasu terrane in Peninsular Thailand and the NW Malay Peninsula (Fang 1994). In Indonesia *Stereochia* was reported as '*Productus semireticulatus*' from Timor (Beyrich 1865, Broili 1916) and

from the Padang Highlands, West Sumatra (Woodward 1879). It is also the dominant brachiopod genus associated with the Early Permian Jambi flora of SW Sumatra (*S. semireticulatus* or *S. irianensis*; Hasibuan et al. 2000, Crippa et al. 2014).

Permian Crinoids and Blastoids

Timor Island has long been famous for its unique Permian deposits with abundant, diverse and well-preserved crinoid and blastoid faunas. Wanner (1923) identified 239 crinoid species in 75 genera. Two-thirds of these species are not known outside Timor (Wanner 1924, Webster 1998). Half of all crinoid species are poteriocrinids, with dominant genera *Timorocrinus, Ceriocrinus, Paraburs acrinus*, etc.

Most of the Timor crinoids and blastoids are from red-brown marls and tuffs with interbedded limestones, a formation named Maubisse Formation in Timor Leste or Sonnebait Series in older literature on West Timor (Figure X.35). They were believed to be relatively warm, shallow marine deposits, but they may actually be mostly hemi-pelagic organisms that ended up in clastics-free deep water carbonates that are often associated with basic volcanics (seamounts?).

The richest occurrences of crinoids are in the Basleo area near Niki-Niki, and are probably from exotic blocks in Neogene melange deposits, and are associated with cephalopods of Middle Permian age (Haniel 1915). Crinoid assemblages from the Amarassi region of SW Timor are less diverse and probably of Late Permian age (Wanner 1923).




Figure X.35. Example of well-preserved crinoids from the Basleo area, Timor (Wanner 1923).

Permian blastoid assemblages from Timor have the highest abundances and diversity in the world. The main monographs on blastoids are Wanner (1924, 1940). Of the 13 genera known from Timor only three or four also occur outside Timor. (Figure X.35).




Figure X.36. Typical example of blastoid from Koeafeoe, West Timor, Schizoblastus (= Deltoblastus) delta (Wanner 1924).

The only other place in SE Asia where some of the Timor species of crinoids and blastoids were found is in the late Early- Middle Permian Ratburi Limestone of Peninsular Thailand (Racey et al. 1994; Sibumasu Terrane). Species of 'Basleo fauna' include the crinoids *Timorocrinus pumulus* Wanner 1924, *Paraburs acrinus* and *Timorocidaris sphaeracantha* Wanner 1920, and the blastoid *Deltoblastus permicus* (Wanner 1910). Although they are present in much smaller numbers in Peninsular Thailand than at the Timor localities, their presence does suggest they were in the same faunal province around Artinskian time.

Outside SE Asia rare *Deltoblastus* have been reported from Oman and Sicily, both also on Cimmerian terranes.

Permian floras

Permian plant fossils, often associated with thin coal beds, are known from West Sumatra and West Papua, with less well-documented occurrences in NW Kalimantan and Belitung island.

The Early Permian floras from Sumatra, NW Kalimantan and Belitung are generally perceived to be warm-Cathaysian floras, while cooler-Gondwanan *Glossopteris* floras (but mixed with some Cathaysian elements) are present in West Papua. The Cathaysian nature of the Jambi flora plays an important role in the plate reconstruction history of Sumatra (Barber et al. 2005).

The famous Early Permian 'Jambi flora' from the Merangin River area in SW Sumatra was first discovered by Tobler, and described by Jongmans and Gothan (1925) (Figure X.37). The flora was initially viewed as of Euramerican affinity, without Gondwanan or Cathaysian *Gigantopteris* flora elements. However, after the Jambi paleobotanical expedition of 1925-1927 collected additional material, Jongmans and Gothan (1935) also recognized some North Cathaysian species.

The Jambi flora was recently re-sampled and studied by a group from Naturalis Museum, Leiden, and the Geological Survey of Indonesia (Van Waveren et al., 2007, Booi et al., 2009). They also recognize affinities to Cathaysian flora, but argue that is not a fully Cathaysian flora, but its greatest similarity is with floras from North China, either the Artinskian Shansi Series (Asama et al. 1975) or the Kungurian Lower Shihhotse beds (Van Waveren et al. 2007). The Jambi Flora is probably best characterized as a late Early Permian temperate subgroup of the true low-latitude Cathaysian floral province.

The age of the Jambi flora is Early Permian, probably Late Asselian or Sakmarian, but exactly what stage within the Early Permian remains to definitively established (see duscussion of fusulinids above).



Figure X.37. Permian flora from the Merangin River area, West Jambi, West Sumatra: 1. Lepidodendron mesostigma, Mengkarang River; 2. Pecopteris arborescens, Sungei Garing (Jongmans and Gothan 1925, 1935).

Permian plant assemblages are also known from West Papua, both the Birds Head and areas south of the Central Range. They were first described by Jongmans (1940, 1941), who documented only Cathaysian and Euramerican species (*Taeniopteris, Pecopteris, Sphenophyllum*). Hopping and Wagner (in Visser and Hermes, 1962) also identified' Gondwanan' *Glossopteris* and *Vertebraria*. The West Papua floras are generally viewed as mixed floras, dominated by Gondwanan elements, but with common Cathaysian elements (Asama et al. 1975, Li and Wu 1994, Rigby 1998, 2001).

A poorly known Permian plant assemblage was also reported from SE Belitung island by Van Overeem (1960). It was provisionally identified by Jongmans as a Permian Cathaysian (*Gigantopteris*) flora, but has never been described. No plant fossils are known from Timor, where Permian sediments are in marine facies.

The existence of mixed Gondwanan- Cathaysian floras in West Papua (and in parts of mainland SE Asia like Thailand and Laos is significant for Permian plate reconstructions. Because *Glossopteris* and many Cathaysian plants like *Gigantopteris* have relatively large seeds, which are unlikely be dispersed across wide oceans. These mixed Permian floras therefore suggest some configuration of land connections (with only very narrow seaways) between the 'Cathaysian' and Gondwana provinces in Permian time, not a wide Paleotethys Ocean.

TABLE 12 - PERMIAN

FAUNA/FLORA	AREA	REFERENCES
Permian General	Timor	Wanner 1926, Charlton et al. 2002
	Sumatra	Fliegel 1901, Fontaine and Gafoer 1989
Mixed *Glossopteris*-Cathaysian flora	West Papua,	Jongmans 1940, 1941, Hopping and Wagner in Visser and Hermes 1962, Rigby 1997, 1998, 2001, Playford and Rigby(2008, Srivastava and Agnihotri 2010
'Cathaysian' flora	West Sumatra	Posthumus 1927, Jongmans and Gothan 1935, Asama 1976, Li and Wu

		1994, Van Waveren et al. 2005, 2007, Booi et al. 2008, 2009
	Belitung	Jongmans in Van Overeem 1960
	NW Kalimantan	Jongmans in Zeijlmans 1939
Palynoflora	West Papua	Playford and Rigby 2008
	Australia	Kemp et al. 1977
Permian- E Triassic Radiolaria	Malay Peninsula	Sashida et al. 1993, 1995, Spiller and Metcalfe 1994, 1995, Jasin 1997
	Thailand	Kamato et al. 2008, 2013
Crinoids	Timor	Wanner 1916,1923, 1929- 1951, De Maresz Oyens 1940
	Belitung	Strimple and Yancey 1976
Blastoids	Timor	Wanner 1924, 1940, Breimer and Macurda 1965, 1972, Webster 1998, Sprinkle and Waters 2013
Ammonoids	Timor, Leti	Wanner 1915, 1932, Haniel 1915a,b, Smith 1927, De Roever 1940, Gerth 1950, Furnish and Glenister 1971, Glenister et al. 1973
	Bangka/Belitung	Kruizinga 1950,
	West Papua	Glenister et al. 1983
Molluscs	Timor	C. Wanner 1922, 1940, 1942, J. Wanner 1940, Hasibuan 1994
	Sumatra	Fliegel 1901
	West Papua	Dickins and Skwarko 1981
Brachiopods	Sumatra	Fliegel 1901, Meyer 1922, Tan Sin Hok 1933, Hasibuan et al. 2000, Crippa et al. 2014
	Timor, Leti, Roti	Rothpletz 1892, Broili 1915, 1916, 1922, Krumbeck 1924, Hamlet 1928, Wanner and Sieverts 1935, Shimizu 1966, Archbold and Barkham 1989, Archbold and Bird 1989, Kato et al. 1999, Winkler Prins 2008
	West Papua	Visser and Hermes 1962, Archbold 1981, 1991, Archbold et al. 1982
Permian Fusulinid foraminifera	West Sumatra	Geinitz 1876, Volz 1904, Von Staff 1909, Lange 1925, Ozawa 1929, Tan Sin Hok 1933, Thompson 1936a,b, Hahn and Weber 1981, Fontaine 1983, Vachard 1989, Ueno 2003, 2006
	South Sumatra	De Neve 1949, 1961
	Bangka, Belitung	De Roever 1951, Van Overeem 1960
	NW Kalimantan/ W Sarawak	Krekeler 1932, 1933, Tan Sin Hok (in Krekeler 1933), Cummings 1962, Sanderson 1966, Fontaine 1990, Vachard 1990, Sakamoto and Ishibashi 2002
	NE Kalimantan (Upper Kutai)	Tan Sin Hok 1930, Sugiaman and Andria 1999
	Timor, Roti, Leti	Schubert 1915a,b, Thompson 1949, Nogami 1963, Charlton et al. 2002, Davydov et al. 2013
	West Papua	Visser and Hermes 1962
Tubiphytes	Sumatra	Tien 1986
	Timor	Riding and Barkham 1999 (= *Shamovella*)
Corals	Timor	Gerth 1921, Koker 1924, Wang 1947, Von Schouppe and Stacul 1955, 1959, Minato and Kato 1965, Niermann 1975, Sorauf 1984, 2004
	West Sumatra	Volz 1904, Minato and Kato 1965, Nguyen Duc Tien 1989a,b
	Thailand	Fontaine et al. 1994
Smaller foraminifera	Sumatra	Nguyen Duc Tien 1989a,b
	Sibumasu	Zhao and Zhou 1987, Wang et al. 2013
Conodonts	Timor	Van den Boogaard 1987
	SE Asia	Mei and Henderson 2001, 2002
Ostracodes	Timor	Grundel and Kozur 1975, Bless 1987

X.6. Quaternary Hominids, Mammals

This sub-chapter X.6 of Bibliography 8.0 contains 146 pages with 1267 papers on Quaternary mammals, including hominids, from the Indonesian region, as well as related forms from mainland SE Asia. Additional titles on this topic may be in area chapters.

Indonesia has long been an area of great interest for the study of the Late Pliocene- Pleistocene vertebrate faunas, including hominids, and their evolution and dispersal from mainland Asia across the islands of Java, Sulawesi and farther East. There are many additional papers on this large field of research, but not all could be included in this bibliography. Early 1900's milestone work has been by Dubois and Von Koenigswald.

Relatively rich Eocene and Miocene land mammal assemblages are known from Thailand, Myanmar and other areas of mainland SE Asia. Remarkably, no equivalent deposits are known from Indonesia, although parts of Sumatra and Kalimantan must have been land areas at that time, with likely land bridges to mainland SE Asia. Only two presumably Eocene mammal occurences are known from Indonesia: remnants of acanthrotheres from Kalimantan (Stromer 1931) and an enigmatic occurrence on Timor (Von Koenigswald 1967, Ducrocq 1996).




Figure X.38. Main Quaternary vertebrates localities in Indonesia (Aziz et al. 1995). 1. Padang Highlands (Lidah Ayer, Jambu and Sibrambang caves), 2. West Java (Cijulang, Cijurey, Citarum, etc), 3. Central Java (Bumiayu, Sangiran, and Patiayam), 4. East Java (Trinil, Kedungbrubus, Ngandong, Widodaren, Sampung, Wajak, Mojokerto), 5. North Borneo (Niah Cave), 6. Brunei, 7. East Kalimantan (Samarinda), 8. Sangihe Island : (Pintareng), 9. South Sulawesi (Walanae), 10. Sumba (Kawangu), 11. Flores (Ola Bula, Ola Kile, Mata Menge (Bajawa Area), 12. Timor (Atambua).

Land mammals started to appear on Java, Sulawesi and Flores relatively late in geologic history, i.e. at around 2.5 Ma (or later?), as these areas emerged to form more significant land areas during Late Pliocene-Pleistocene tectonic uplift processes and volcanism.

Migration of land mammals from the SE Asia mainland to Java was also facilitated during Pleistocene glacial periods of sealevel lowstands, when the Sunda shelf was largely exposed and became the most likely migration route (e.g Sartono 1987).

Pre-Pleistocene mammal fossils are extremely rare in Indonesia, although relatively diverse mammal assemblages are known from the Eocene- Pliocene of mainland SE Asia. This is probably mainly a reflection of the rarity of pre-Pleistocene terrestrial deposits in the archipelago. Eocene acanthocerids (*Hippopotamus* family), common in Asia at that time, were reported from West Kalimantan and from Timor. The latter find is another indication that parts of what is now Timor Island ('Banda Terrane') may have been attached to Sundaland in Eocene time.

Present-day distributions of faunas and floras still reflect plate tectonic past history. Well-known biogeographic boundaries like the Wallace Line (1869), separating balanced 'Asian' Sundaland faunas from unbalanced island faunas on Sulawesi, Flores and islands farther East, and Lydekkers Line, separating the Australian faunas of Australia and New Guinea in the SE from the impoverished island faunas to its West.

The history of latest Pliocene- Recent terrestrial vertebrate evolution and dispersal is best documented on Java, where there is a succession of faunas reflecting island conditions in the (latest Pliocene-) Early Pleistocene (Satir Fauna), followed by the arrivals of more diverse Asian mammal populations (including *Homo erectus*) in the Middle Pleistocene around 1 Ma (Figure X.39.).




Figure X.39. Succession of Pleistocene mammal assemblages of Java (De Vos 1995).

Pleistocene vertebrate faunas from SW Sulawesi have also been interpreted as impoverished island faunas of Asian origin.

Island faunas

Early island populations are typically composed of a limited number of migrant species of Asian origin that are relatively good swimmers (elephants, hippos, deer; not humans or other predators). In the absence of carnivorous predators, mammal species commonly developed into either dwarfed species (pygmy *Stegodon* elephants) or giants (giant tortoises, rats, Komodo lizards, etc.) in the Pleistocene of, Timor, SW Sulawesi and Sumba. Similar forms from East Java and West Java suggest these parts of Java were also islands in the Early Pleistocene.

Pleistocene hominids

The first fossil evidence for human evolution and migration came with the discovery of 'Java Man' (*Pithecanthropus erectus*; now called *Homo erectus*) at Trinil by Dubois in 1891. Finding this rare material was a remarkable story of perseverance and luck. Additional *Homo erectus* fragments were reported in the 1930's and later.

Younger, but still primitive Upper Pleistocene hominid fossils were found in a Solo River terrace at Ngandong in East Java, named *Homo soloensis* ('Solo Man') and represent another much-debated group of hominids transitional between *H. erectus* and *H. sapiens*. Latest age dating of these skulls is in the 40-70 ka range.

The incomplete fossil record, the often poorly documented stratigraphic positions of much of the fossil hominid material (much of it collected by local farmers), and the notoriously inconsistent results of various radiometric direct dating methods of hominid fossils still leaves much room for debate on timing of arrivals, evolution and extinctions of hominids in the SE Asia region.

The most widely accepted interpretation of *Homo erectus* history has been an evolutionary transition from *Homo habilis* in East Africa around 1.8-1.6 Ma and a migration into SE Asia/ Java around 1.0 Ma. However, Swisher et al. (1994) reported a radiometric age of 1.81 Ma for a *Homo erectus* skull from Mojokerto, East Java, the oldest date for any *Homo erectus*, and suggested it therefore probably originated in Asia instead of Africa. However, their dating is on tuff samples that may not be from the same horizon as the *Homo erectus* skull (De Vos and Sondaar 1994). Some recent Ar/Ar dating of pumice layers at Sangiran dome yielded an age range of the *Homo erectus*-bearing interval between 1.51- 1.01 Ma.




Figure X.40. Examples of Homo erectus fossils from Java (from T. Ingicco presentation). There is an apparent half-million year gap in the fossil record of Java between older Homo erectus of Sangiran- Trinil and younger populations from Ngandong- Sambungmacan (actual gap may actually be wider than indicated here)

Homo floresiensis

The Liang Bua cave on Flores island was the site of the latest significant discovery of a cave-dwelling, dwarf population of hominids of Late Pleistocene age (95-13 ka). It was named *Homo floresiensis*, adults are only one meter tall, and it may overlap in time with 'Solo Man' and with modern man *(Homo sapiens)*. One of the more recent age estimates places *H. floresiensis* between ~100- 60 ka (Sutikna et al. 2016)

This discovery has been the subject of much debate. Most authors view it as a late descendants of the *Homo erectus* group, that underwent dramatic island dwarfism, although many of its characteristics are more like *Homo erectus* that is older than the Central Java *H. erectus* (e.g. Kaifu et al. 2011, 2015). Others viewed it as aberrant forms of early *Homo sapiens*, affected by diseases like Down syndrome or microcephalism (Henneberg et al. 2008, 2014 and others). Jacob et al. (2006) and others have argued that these Flores hominids should be regarded as a dwarf population of *Homo sapiens*. Brown & Maeda (2009) and Argue at al (2017) argued that *H. floresiensis* are not descendants of either *H. erectus* or *H. sapiens.*

Older *Homo erectus* may have reached the island of Flores by 0.8 Ma, as suggested by relatively well-dated stone tools from the Mata Menge site, associated with a *Stegodon* mammal assemblage and tektites of the Australiasian tektite event of ~780 ka (Maringer & Verhoeven 1970, Sondaar et al. 1994, Morwood et al. 1997, Van den Bergh et al. 2009). The stone tools resemble those from Sangiran, C Java (Van Heteren and De Vos 2012), which are associated with classic *H. erectus*. Initially no hominid fossils were found at this site, but recent excavations yielded *H. erectus* or *H. floresiensis*-like mandible and teeth. (Van de Bergh et al. 2016).

As in Central Java, there is a significant gap in the fossil record of hominids and other mammals in Flores between the ~800ka and the younger than 100 ka hominid assemblages.

Arrival of modern humans (Homo sapiens)

Anatomically modern humans (*Homo sapiens*) are commonly believed to have migrated from Africa, and arrived into island SE Asia by 75- 80 ka and reached Australia ~50,000 years ago (Oppenheimer 2009). Recent dating of deposits and fossils associated with Late Pleistocene modern human teeth excavated in a cave in West Sumatra by Dubois (1891) appears to confirm an arrival date before ~70 ka (Westaway et al. 2017).




Figure X.41. Paleogeography of Southeast Asia during Pleistocene, showing land areas during sea level lowstands and likely general mammal migration routes (Sartono 1973)

Mammal migration routes

Several authors have speculated on the possible migration routes and times of migration of mammals (including hominids) from SE Asia (e.g. Sartono 1973, Fig. X.41).

Obvious times of migration would be during Pleistocene glacially-driven sealevel lowstands, like the -120m sealevel drop at the Last Glacial Maximum at ~18,000 years ago or the -160m drop at ~160,000 years ago (e.g. Molengraaff , Voris 2000). These lowered sea levels would have exposed all of the Sunda Shelf, making for easy migrations into Sumatra and Borneo and into S Sulawesi from the south, but also would have narrowed some of the seaways to islands further East.

Possible land bridges and likely migration routes into Eastern Indonesia and the Philippines were discussed by Heaney 1985, Sartono 1973, 1987, Tjia 2006, De Vos 1995, etc.

X. REFERENCES INDONESIA- SE ASIA: PALEONTOLOGY, BIOSTRATIGRAPHY

The following chapters on paleontology-biostratigraphy papers tend to be of a more general nature. Numerous additional papers with biostratigraphy content that are case studies with geological implications for specific areas, are listed under those areas;and are not repeated here. These chapters also include titles from Circum-Indonesian areas that may be relevant for interpretations of Indonesian materials.

X.1. Quaternary/Recent faunas-microfloras and distribution, Biogeography

Adisaputra, Mimin Karmini (1985)- Paleontological analyses of the Sawu Basin, Lombok basin and Argo abyssal plain. Proc. 14th Annual Conv. Indonesian Association Geologists (IAGI), Jakarta, p. 205-221.
(online at: https://www.iagi.or.id/web/digital/40/PIT_IAGI_1985_paper19.pdf)
(Recent foraminifera distribution in seafloor samples collected during Snellius-II Expedition)

Adisaputra, Mimin K. (1988)- Late Quaternary calcareous nannoplankton in the surface sediment of Makasar and Flores basin, Indonesia. Bull. Marine Geological Institute 3, 1, p. 25-36.

Adisaputra, Mimin K. (1989)- Planktonic foraminifera in recent bottom sediments of the Flores, Lombok and Savu Basins, eastern Indonesia. In: J.E. van Hinte et al. (eds.) Proc. Snellius II Symposium, Jakarta 1987, Netherlands J. of Sea Research 24, 4, p. 465-475.
(Planktonic foraminiferal assemblages differ between Flores, Lombok and Savu Basins. In Flores Basin Ng. dutertrei dominant, followed by Gr. menardii, Pulleniatina obliquiloculata, Hastigerina siphonifera and Globigerina bulloides. Lombok and Savu Basins dominated by Gr. menardii, with Pulleniatina obliquiloculata, Gs. ruber and Gr. tumida)

Adisaputra, Mimin K. (1991)- Mikrofauna dan potensi wisata perairan Benoa, Bali. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 1, 2, p. 2-6.
('Microfauna and recreational potential of the water of Benoa, Bali'. Beach sand at Benoa, S of Denpasar, SE Bali, contains abundant foraminifera of genus Schlumbergerella, also Amphistegina lessoni, Calcarina calcar, Tinoporus spengleri, Baculogypsisna sphaerulata, Operculina, etc.)

Adisaputra, Mimin K. (1992)- Mikrofauna perairan Muria. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 2, 5, p. 11-16.
('Microfauna in the waters of Muria'. Shallow marine small benthic forams of Java Sea N of Muriah volcano. At less than 20m common Ammonia beccarii, Eponides praecinctus and Asterorotalia trispinosa. In deeper parts more Ammonia annectens, Pseudorotalia schroeteriana and Quinqueloculina. Not much detail)

Adisaputra, Mimin K. (1992)- Late Neogene planktonic foraminifera of the Makasar Basin. Bull. Marine Geological Institute 7, 1, p. 15-21.
(Abundant planktonic foraminifera in Makassar Straits bottom samples between 42-2300m, collected during Snellius II expedition. Globigerinoides ruber dominant in North, Neogloboquadrina dutertrei in S part)

Adisaputra, Mimin K. (1996)- Planktonic foraminifera and oxygen isotope records in two cores from the Banda Sea and Indian Ocean. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 6, 57, p. 10-17.
(Study of Quaternary planktonic foraminifera from two ~7-8m long French- Indonesian cores collected in 1990, one SW of Timor (2313m), one in Banda Sea depth (3163m). Neogloboqudrina dutertrei is indicator of rel. low salinity, Globorotalia menardii for rel. high salinity)

Adisaputra, Mimin K. (1996)- Biostratigrafi kuarter sedimen dasar laut perairan Indonesia bagian Timur dan Samudera Hindia. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 6, 59, p. 2-6.
('Stratigraphy of Quaternary seafloor sediments in waters of eastern Indonesia and the Indian Ocean'. Study of Quaternary planktonic foraminifera from 46 seven meter long seafloor cores in deep waters of E Indonesia and adjacent Indonesian Ocean. Four subzones in Quaternary Globorotalia truncatulinoides zone (N22-N23), from

top: (1) Globorotalia fimbriata- Bolliella adamsi, (2) Globigerina calida, (3) Globorotalia crassaformis hessi and (4) Globigerinoides cyclostomus))

Adisaputra, Mimin K. (1997)- Foraminifera sedimen permukaa perairan Selat Bangka-Belitung. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 7, 70, p. 2-10.
('Foraminifera from seafloor sediments of Bangka-Belitung Straits'. 38 seafloor samples from straits S of Bangka and Belitung in water depths 10-50m. Dominant foraminifera Operculina ammonoides and other spp., followed by Quinqueloculina, Amphistegina lessonii, Elphidium, Cellanthus, Pseudorotalia schroeteriana, etc.)

Adisaputra, Mimin K. (1998)- *Schlumbergerella floresiana* accumulation in coastal zone of Bali and Nusatenggara, Indonesia: implementation for tourism. Proc. 33rd Session Coord. Comm. Coastal and Offshore Programmes E and SE Asia (CCOP), Shanghai 1996, p. 310-316.
('White sands' along coasts of E Bali, W Lombok, N Sumbawa and S Flores composed mainly of rounded foraminifera Schlumbergerella floresiana (formerly also called Tinoporus, Baculogypsina, Baculogypsinoides; JTvG). Forams derived from adjacent coral reefs)

Adisaputra, Mimin K. (1998)- Foraminifera bentos pantai Senggigi, Lombok Barat dan asosiasinya; faktor penunjang pariwisata. Proc. 27th Annual Conv. Indonesian Association Geologists (IAGI), 2 (Sedimentology Paleontology Stratigraphy), Yogyakarta, p. 53-65.
(online at: https://www.iagi.or.id/web/digital/56/32.pdf)
('Benthic foraminifera of Senggigi Beach, W Lombok, and its associations; factors supporting tourism'. Senggigi beach with abundant foraminifera derived from adjacent reef flat, dominated by Schlumbergerella floresiana. Also Baculogypsina, Baculogysinoides spinosus, Amphistegina lessonii, Calcarina calcar))

Adisaputra, Mimin K. (2000)- Recent foraminifera on the coast and offshore of East Lombok, Eastern Indonesia. Proc. 36th Session Coord. Comm. Coastal and Offshore Programmes E and SE Asia (CCOP), Hanoi 1999, p. 181-200.
(Benthic foraminifera from East Lombok coast and Alas Strait shallow waters down to 90m. In N and central parts Amphistegina lessonii dominant and associated with Calcarina. In S Asterorotalia ('Rotalinoides') gaimardii dominant, still with Amphistegina. Beach samples in N with common Schlumbergerella and Baculogypsinoides, derived from coral reef. Planktonic foraminifera rare)

Adisaputra, Mimin K. (2000)- Late Neogene planktonic foraminiferal biostratigraphy of two cores in Timor waters, Indonesia. Majalah Geologi Indonesia (IAGI) 24, 1, p. 39-50.

Adisaputra, Mimin K. & M. Hendrizan (2011)- Foraminifera perairan Balikpapan, Kalimantan Timur: lingkungan pengendapan dan pengaruhnya. J. Geologi Kelautan 9, 2, p. 119-133.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/205/195)
('Foraminifera from waters off Balikpapan, East Kalimantan: depositional environment and its effects'. 195 benthic and 34 planktonic foraminifera species in Makassar Straits seafloor samples off Balikpapan, between 18-562m depth. Asterorotalia trispinosa dominant around 20m, Heterolepa praecincta most abundant from ~50-300m, Karreriella brady and Uvigerina spp. common only >300m, etc. Cycloclypeus only at 71 and 83m. Abundant planktonic foraminifera Neogloboquadrina dutertrei below 100m, indicating rel. low salinity.)

Adisaputra, Mimin K., M. Hendrizan & A. Kholiq (2010)- Katalog foraminifera perairan Indonesia. Pusat Puslitbang Geologi Kelautan, Bandung, p. 1-198.
('Catalog of Foraminifera collected from Indonesian seas')

Adisaputra, Mimin K. & D. Rostyati (2000)- Recorded Recent foraminifera in the surface sediment of Sunda Strait water. Proc. 29th Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 4, p. 117-132.
(Foraminifera from 12 seafloor samples in Sunda Straits and adjacent Indian Ocean between 52- 2180m. Rel. common planktonic foram Neogloboquadrina dutertrei, possibly related to rel. low salinity. Most common benthics Bulimina and Bolivina (mostly 490-1580m). Hyalinea balthica between 489-1078m)

Adisaputra, Mimin K. & D. Rostyati (2003)- Foraminifera sedimen dasar Laut Delta Mahakam, Kalimantan Timur. J. Geologi Kelautan 1, 3, p. 1-10.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/98/88)
('Foraminifera in sediments offshore Mahakam Delta, E Kalimantan'. Foram distribution in samples mainly from 10-100m water depth. 39 species of planktonic, 149 benthic foraminifera. Common benthics Amphistegina lessonii, Heterolepa, Operculina and Pseudorotalia schroeteriana. Max. diversity at depth of 63,5 m)

Amijaya, H., Ngisomuddin & Akmaluddin (2010)- Characterization of July 17, 2006 tsunamiite at South coast of West Java. J. Southeast Asian Applied Geology (UGM) 2, 1, p. 35-39.
(online at: https://journal.ugm.ac.id/index.php/jag/article/viewFile/7232/5672)
(Deposits of July 26 tsunami at Pangandaran Beach, W Java. Mainly f-m sand, ~10-12cm thick, separated from older beach sediment by erosional surface. Sedimentary structures parallel lamination and current ripples. No vertical fining trends. With transported shallow and deeper marine benthic foraminifera, incl. Ammonia, Elphidium, Amphistegina, Cibicides., Biginerina, Bolivina, Bathysiphon, Nodosaria and Quinqueloculina)

Anderson, J.A.R. (1963)- The flora of the peat swamp forests of Sarawak and Brunei, including a catalogue of all recorded species of flowering plants, ferns and fern allies. Singapore Gardens Bull. 20, p. 131-228.
(All modern coastal and deltaic peat swamps of N Borneo raised bog type. 243 plant/tree species, in 6 communities: 1) Mixed swamp forest, 2) Alan forest, 3) Alan bunga forest, 4) High pole forest, 5) Low pole forest, 6) Padang keruntum)

Ardi, R.D.W, Aswan, K.A. Maryunani, E. Yulianto, P.S Putra & S.H. Nugroho (2023)- First occurrence of planktonic foraminiferal species *Boliella adamsii* as a marker for the Pleistocene-Holocene boundary in the sea around Sumba Island. Proc. Second Int. Seminar on Earth Sciences and Technology (ISEST-2023), Bandung, IOP Conference Series: Earth and Environmental Science 1245, 012025, p. 1-12.
(online at: https://iopscience.iop.org/article/10.1088/1755-1315/1245/1/012025/pdf)
(Pleistocene- Holocene boundary depth in cores off Sumba Island nearly coeval with first occurrence of planktonic foraminifer Boliella adamsii (N.B.:both genus and species names missspelled; should be Bolliella adamsi; JTvG))

Ardi, R.D.W., Aswan, K.A. Maryunani, E. Yulianto, P.S Putra & S.H. Nugroho (2023)- Sea surface paleoproductivity reconstruction based on foraminiferal accumulation rate in the western Savu Strait since the Last Glacial Maximum (~23 ka BP). Rudarsko-Geolosko-Naftni Zbornik (Mining-Geology-Petroleum Engineering Bulletin), Zagreb, 38, 1, 62, p. 167-178.
(online at: https://hrcak.srce.hr/ojs/index.php/rgn/article/view/23308/13606)

Ardi, R.D.W., K.A. Maryunani, E. Yulianto, P.S Putra & S.H. Nugroho (2019)- Biostratigrafi dan analisis kedalaman termoklin di Lepas Pantai Barat Daya Sumba sejak Pleistosen Akhir berdasarkan kumpulan foraminifera planktonik. Bulletin of Geology (ITB) 3, 2, p. 355-362.
(online at: https://buletingeologi.com/index.php/buletin-geologi/issue/download/8/Paper%203%20-%20Ryan%20Dwi%20Wahyu%20Ardi)
('Biostratigraphy and thermocline depth analysis in the Southwest Sumba Offshore since the Late Pleistocene based on planktonic foraminifera assemblages'. Late Pleistocene- Holocene 235cm long deep marine core S8, off SW Sumba. Relatively abundant thermocline dwellers Neogloboquadrina dutertrei, Pulleniatina obliquiloculata and Globorotalia menardii during Pleistocene, indicating shallower thermocline (eutrophic condition), and tied to SE monsoon domination during Pleistocene (stronger Indonesian Throughflow))

Aswan, Y. Zaim & Y. Rizal (2006)- Distribution of Quarternary freshwater molluscs fossils in Jawa. In: Y. Zaim et al. (eds.) S. Sartono: dari hominid ke delapsi dengan kontroversi, Penerbit ITB, Bandung, Chapter 9, p. 109-120.

Auliaherliaty, L., K.T. Dewi & Y.A. Priohandono (2004)- Foraminifera di Teluk Sepi- Blongas, Lombok selatan, Nusa Tenggara Barat dan kaitannya dengan faktor lingkungan. J. Geologi Kelautan 2, 3, p. 1-8.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/115/105)

(Seafloor sediment samples of 1-78m in Sepi-Blongas Bay, S Lombok, with 133 foraminiferal species, mainly Ammonia beccarii, Elphidium crispum, Pattelina, Amphistegina lessonii, Calcarina, Pyrgo, Quinqueloculina, etc. Genus Calcarina with 15 species (?))

Baccaert, J. (1976)- Soritidae of the Lizard Island reef complex: a preliminary report. Annales Soc. Geologique Belgique 99, p. 237-262.
(online at: https://popups.uliege.be/0037-9395/index.php?id=5366)
(Eight species of soritids, incl. Marginopora vertebralis)

Baccaert, J. (1986)- Foraminiferal bio- and thanatocoenoses of reef flats, Lizard Island, Great Barrier Reef, Australia: nature of substrate. Annales Soc. Royale Zoologique Belgique 116, 1, p. 3-14.
(online at: https://www.vliz.be/imisdocs/publications/ocrd/278944.pdf)
(Foraminifera from three reef flats of of Lizard Island archipelago in N Great Barrier Reef. Highest abundances reached by Soritidae (Peneroplis, Sorites, Marginopora, Spiroloculina, etc.), Amphistegina lobifera, Calcarinidae (Calcarina spengleri, Baculogypsina sphaerulata) and Elphidiidae (E. crispum). 104 species identified. Distributions closely related to distribution of algal substrate)

Baccaert, J. (1987)- Distribution patterns and taxonomy of benthic foraminifera in the Lizard Island reef complex, northern Great Barrier Reef, Australia. Ph.D. Thesis, Universite de Liege, 4 vols., 146p, 290p., 109p.
(online at: https://vliz.be/en/imis?module=person&persid=10828)
(vol. 1- Text online at: https://www.vliz.be/imisdocs/publications/ocrd/240380.pdf)
(vol. 3- 109-plate Atlas online at: https://www.vliz.be/imisdocs/publications/ocrd/240384.pdf

Barbin, V., J.C. Cailliez & D. Decrouez (1987)- Sable a *Schlumbergerella floresiana* (foraminifere) et *Conus mobilis skinneri* (gasteropode) de Kesuma Sari (SSE Bali, Indonesie). Revue de Paleobiologie, Geneve, 6, 1, p. 159-164.
(Carbonate sands composed of large globular foram Schlumbergerella floresiana and Conus gastropods in SE Bali)

Barker, R.W. (1960)- Taxonomic notes on the species figured by H.B Brady in his report on the foraminifera dredged by the H.M.S. 'Challenger' during the years 1873-1876. Special Publ. SEPM 9, p. 1-238.
(Useful taxonomic revision of beautifully illustrated modern deep water foraminifera book of Brady (1884) (see also Jones (1994, 2014))

Barmawidjaja, D.M. (1991)- Studies in living and fossil foraminifers from seasonally productive regions. Ph.D. Thesis University of Utrecht, Geologica Ultraiectiana 82, p. 1-221.
(online at: http://dspace.library.uu.nl/handle/1874/238680)
(Collection of papers on subrecent foraminifera in Adriatic Sea, semi-enclosed Kau Basin (Halmahera; sill depth 40m) and Molucca Sea. Three categories of species: epifaunal, predominantly infaunal and infaunal. Kau Basin, Halmahera was freshwater lake during Last Glacial Maximum, became reconnected with open ocean about 10 ka due to sea level rise at Pleistocene- Holocene transition. Dysoxic bottom conditions prevailed throughout Holocene. Piston core off Halmahera in N Molucca Sea suggest Pleistocene glacial climate was drier than today and surface water T 2.5° C lower than today)

Barmawidjaja, D.M. (1993)- Sebaran mikrohabitat Holosen bentos foraminifera di Teluk Kau. Proc. 22nd Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 2, p. 1114-1129.
('Distribution of Holocene benthic foraminifera microhabitats in Kau Bay'. In English. Three piston cores of 8.5m long in Late Quaternary sediments from 20, 260 and 460m water depth in Kau Bay, NE Halmahera. Kau Bay was freshwater lake in Weichselian (Late Pleistocene Last Glacial Maximum. Foram assemblages dominated by opportunistic infauna (Bolivina spp., Bulimina marginata, Hopkinsina) and opportunistic epifauna, mainly responding to relatively low oxygen levels in basin after marine flooding. Rel. common Cassidulina crassa and Cancris auriculus)

Barmawidjaja, D.M. (1994)- Pengaruh lingkungan terhadap sebaran foraminifera plangton di Teluk Kau, Halmahera. Proc. 23rd Annual Conv. Indonesian Association Geologists (IAGI), Jakarta, 1, p. 173-183.
('Influence of environment on planktonic foraminifera distribution in Kau Bay, Halmahera')

Barmawidjaja, D.M., R. Kapid & B. Dwiyanto (1996)- Environmental factors controlling the distribution of benthonic foraminifera of Jakarta Bay. Proc. 25th Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 2, p. 1-15.

Barton, E.S. (1901)- The genus *Halimeda*. In: M. Weber (ed.) Siboga Expeditie, Uitkomsten op zoologisch, botanisch, oceanographisch en geologisch gebied verzameld in Nederlandsch Oost-Indie 1899-1900, Monographie LX, Brill, Leiden, p. 1-32.
(online at: http://images.algaebase.org/pdf/5964B9370f1d92EC18jmK2CED443/16210.pdf)
(Recent calcareous algae Halimeda rel. widespread in coastal reefs around of E Indonesia: LesserSunda Islands, Sulawesi, Aru and Kai islands and in Banda-Archipelago. Seven species distinguished)

Bassi, D., J.C. Braga, G. Di Domenico, J. Pignatti, S. Abramovich, P. Hallock, J. Konen, Z. Kovacs, M.R. Langer et al. (2021)- Palaeobiogeography and evolutionary patterns of the larger foraminifer *Borelis* de Montfort (Borelidae). Papers in Palaeontology 7, 1, p. 377-403.
(Read-online at: https://onlinelibrary.wiley.com/doi/epdf/10.1002/spp2.1273)
(Paleobiogeography and evolutionary lineages of Late Eocene-Recent alveolinid Borelis species, incl. Indo-Pacific Borelis pygmaea Hanzawa 1930 from Late Oligocene Rajamandala Limestone in W Java))

Bassi, D., Y. Iryu, S. Kinoshita, K. Fujita & J. Pignatti (2023)- A new species of the larger porcelaneous foraminifer *Borelis* provides novel insights into Neogene to Recent western Pacific palaeobiogeographical dispersal patterns. Palaeogeogr. Palaeoclim. Palaeoecology 628, 111764, p. 1-14.
(online at: https://www.sciencedirect.com/science/article/pii/S0031018223003826)
(New Recent porcelaneous larger foraminiferal species: Borelis matsudai (Alveolinoidea, Borelinae), from Sekisei Lagoon, S Ryukyu Islands (Japan; in W Indo-Pacific Warm Pool). Related to Borelis pulchra group)

Belasky, P. (1996)- Biogeography of Indo-Pacific larger foraminifera and scleractinian corals: a probabilistic approach to estimating taxonomic diversity, faunal similarity, and sampling bias. Palaeogeogr. Palaeoclim. Palaeoecology 122, p. 119-141.
(On modern corals and larger foraminifera distribution in Indo-Pacific. Eastward decline in diversity due primarily to shallowing of thermocline and significant cooling of Equatorial Undercurrent in E Pacific)

Biekart, J.W. (1989)- The distribution of calcareous nannoplankton in Late Quaternary sediments collected by the Snellius II Expedition in some southeast Indonesian basins. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B92, 2, p. 77-141.
(Descriptions of Late Quaternary calcareous nannofossils in three Snellius II Expedition (1984-1985) cores from Savu Sea, Timor Trough and Lombok Ridge. Dominant species Florisphaera profunda)

Biekart, J.W. (1989)- Coccolithophores in the Upper Quaternary of some Indonesian basins. In: J.E. van Hinte et al. (eds.) Proc. Snellius II Symposium, Jakarta 1987, Netherlands J. of Sea Research 24, 4, p. 523-530.
(Quaternary calcareous nannofossils from E Indonesia piston cores dominated by Florisphaera profunda)

Biswas, B. (1976)- Bathymetry of Holocene foraminifera and Quarternary sea-level changes on the Sunda shelf. J. Foraminiferal Research 6, 2, p. 107-133.
(Benthic foraminifera distribution on Sunda Shelf. Only 3 species in littoral zone of S China Sea, 107 species at shelf edge. Planktonic species 4% of total fauna at 20-30m, increasing to 65-80% at shelf edge)

Boettger, A. (1890)- Ad. Strubell's Konchylien aus Java I. Bericht Senckenbergische Naturforschenden Gesellschaft Frankfurt 1890, p. 137-173.
(online at: https://www.zobodat.at/pdf/Berichte-der-Senckenberg-naturf-Ges-Frankfurt_1890_0137-0173.pdf)

('Adolf Strubell's mollusc shells from Java-1'. Early description of fresh and brackish water gastropods and bivalves from West Java)

Boettger, A. (1890)- Ad. Strubell's Konchylien aus Java II und von den Molukken. Bericht Senckenbergische Naturforschenden Gesellschaft Frankfurt 1891, p. 241-318.
('Adolf Strubell's mollusc shells from Java II and from the Moluccas')

Brady, H.B. (1884)- Report on the foraminifera dredged by H.M.S. Challenger during the years 1873-1876. Scientific results of the voyage of H.M.S. Challenger during the years 1873-76, Section V, Zoology 9, XXII, London, p. 1-814.
(Classic description of Recent and subfossil benthic foraminifera from deep waters of SE Asia. For revisions of taxonomy see also Barker 1960 and Jones 1994)

Brady, G.S. (1880)- Report on the Ostracoda dredged by H.M.S. Challenger during 1873-1876. Scientific results of the voyage of H.M.S. Challenger during the years 1873-76, Reports 1, Zoology III, p. 1-184.
(Early descriptions of modern ostracods from Challenger seafloor samples, mainly from deep water SE Asia)

Broecker, W.S., E. Clark, J. Lynch-Stieglitz, W. Beck, L.D. Stott, I. Hajdas & G. Bonani (2000)- Late Glacial diatom accumulation at 9°S in the Indian Ocean. Paleoceanography 15, 3, p. 348-352.
(10m-long section in core from 3800m depth at 9°S on Ninety-East Ridge, Indian Ocean, consists of 2/3 of diatom Ethmodiscus rex and 1/3 planktonic foraminifera. Age ~30 ka- 11 ka. During glacial time, Pacific Ocean thermocline waters may have moved above site, providing silica and nutrients required by diatoms)

Bronnimann, P. & A.J. Keij (1986)- Agglutinated foraminifera (Lituolacea and Trochamminacea) from brackish waters of the state of Brunei and of Sabah, Malaysia, Northwest Borneo. Revue de Paleobiologie, Geneve, 5, 1, p. 11-31.

Bronnimann, P., A.J. Keij & L. Zaninetti (1983)- *Bruneica clypea* n.gen. n.sp., a Recent remaneicid (Foraminiferida; Trochamminacea) from brackish waters of Brunei, Northwest Borneo. Revue de Paleobiologie, Geneve, 2, 1, p. 35-41.

Bronnimann, P., J.E. Whittaker & L. Zaninetti (1992)- Brackish water foraminifera from mangrove sediments of southwestern Viti Levu, Fiji Islands, Southwest Pacific. Revue de Paleobiologie, Geneve, 11, p. 13-65.

Bronnimann, P. & J.E. Whittaker (1993)- Taxonomic revision of some Recent agglutinated foraminifera from the Malay Archipelago, in the Millett Collection, the Natural Museum of History, London. Bull. British Museum (Natural History), Zoology, London 59, 2, p. 107-124.

Budiarto, E.Y., M. Hendrizan, R. Setijadi, R.A. Troa, R. Zuraida & E. Triarso (2023)- Foraminiferal paleoproductivity during Holocene in Sunda Strait. Proc. International Conference Radioscience, Equatorial Atmospheric Science and Environment and Humanosphere Science (INCREASE 2022; Indonesia, online), Springer Proceedings in Physics 290, p. 745-755.
(Paleoproductivity in Sunda Strait study area driven by salinity, sea surface temperature, and thermocline depth. Three facies: (I) increase in Globigerina bulloides, indicating paleoproductivity influenced by upwelling and SE monsoon; (II) relatively shallow thermocline reflected by extinction of Globorotalia crassaformis and decreasing Globorotalia hirsuta (increased run-off and terrigenous influx controlled by NW monsoon; (III) decreasing oligotrophic Globigerinoides, indicating eutrophic water during increased run-off)

Burke, S.C. (1981)- Recent benthonic foraminifera of the Ontong Java Plateau. J. Foraminiferal Research 11, p. 1-19.

Burke, S.C., W.H. Berger, W.T. Coulbourn & E. Vincent (1993)- Benthic foraminifera in box core ERDC 112, Ontong Java Plateau. J. Foraminiferal Research 23, p. 19-39.

Carbonel, P. & T. Hoibian (1988)- The impact of organic matter on ostracods from an equatorial deltaic area, the Mahakam Delta- Southeastern Kalimantan. In: T. Hanai (eds.) Evolutionary biology of ostracoda, its fundamentals and applications, Developments in palaeontology and stratigraphy 11, Elsevier, p. 353-366.
(On ostracod fauna in Mahakam delta area. In front of delta mouth number of species decreases, Hemicytheridea reticulata relatively common, and ornamentation of polymorphic species decreases. Between delta mouths ornamentation increases, probably due to less degradation of organic matter here)

Carpenter, W.B. (1883)- Report on the specimens of the Genus *Orbitolites* collected by H.M.S. Challenger during the years 1873-1876. H.M.S. Challenger Reports 7, Zool. XXI, p. 1-47.
(online at: www.19thcenturyscience.org/HMSC/HMSC-Reports/Zool-21/htm/doc.html)
(Descriptions of Recent larger foram Orbitolites complanata from coast of Australia and Fiji (= Marginopora vertebralis Quoy and Gaimard; JTvG))

Chaproniere, G.C.H. (1991)- Pleistocene to Holocene planktonic foraminiferal biostratigraphy of the Coral Sea offshore Queensland, Australia. BMR J. Australian Geology Geophysics 12, 3, p. 195-221.
(online at: www.ga.gov.au/corporate_data/49552/Jou1991_v12_n3.pdf)
(Well-illustrated latest Pliocene- Holocene planktonic foraminifera biostratigraphy (N19-N23) in cores from Queensland and Townsville Troughs)

Chen, C. & H.L. Lin (2017)- Applying benthic foraminiferal assemblage to evaluate the coral reef condition in Dongsha Atoll lagoon. Zoological Studies 56, 20, p. 1-16.
(online at: http://zoolstud.sinica.edu.tw/Journals/56/56-20.pdf)
(Recent foraminifera distribution in Dongsha Atoll, northern S China Sea. Porcelaneous foraminifera dominant (76%, 48 species, miliolids). Fourteen hyaline species (incl. common Calcarina))

Chen, C.W., K.Y. Wei, H.S. Mii & T.N. Yang (2008)- A Late Quaternary planktonic foraminiferal oxygen isotope record of the Banda Sea: chronostratigraphy, orbital forcing, and paleoceanographic implications. Terrestrial Atmospheric Oceanic Sciences (TAO) 19, 4, p. 331-339.
(online at: http://tao.cgu.org.tw/index.php/articles/archive/geology/item/766-2008194331images)
(Oxygen isotope time-scale based on planktonic foram Globigerinoides sacculifer at piston core site MD012380 in water depth 3232m in Banda Sea was established for past 820 ky. Spectral analysis of 18 O time-series reveals distinct periodicities of 100, 41, and 23 ky, indicating strong orbital forcing)

Cheng, Z., C. Weng, S.Y. Foong & Lu Dai (2020)- A study on modern pollen rain and pollen morphology in the tropical western Malay Peninsula and its implications for paleoenvironmental reconstructions in the Sunda region. Review Palaeobotany Palynology 279, 104236, p. 1-44.
(Modern pollen rain samples from western Malay Peninsula. With 29 plates)

Choiriah, S.U., M. Maha. D.N. Aji & K.T. Dewi (2011)- Study of Ostracoda in the Territorial Nusawere, Bay of Pangandaran, West Java. Proc. 7th Malaysia Geoheritage Conference & 4th Indonesia-Malaysia Joint Geoheritage Conference, Kuala Trengganu 2011, p. 1-9. *(online at: http://eprints.upnyk.ac.id/22316/1/5.%20Study%20of%20Ostracoda%20%28Geoheritage%20Malaysia%29.pdf*
(Presentation online at: http://www.ukm.my/wgm/wgm2011pdf/6.%20Siti%20Umiyatun%20Choiriah.pdf)
(70 species of Recent ostracodes in seafloor samples in Pangandaran Bay, off S coast of W Java near Pangandaran. Conclusion: age of ostracodes is Holocene and bathymetry is shallow marine (0-50m) (!))

Cleary, D.F.R., L.E. Becking, N.J. de Voogd, W. Renema, M. de Beer, R.W.M. van Soest & B.W. Hoeksema (2005)- Variation in the diversity and composition of benthic taxa as a function of distance offshore, depth and exposure in the Spermonde Archipelago, Indonesia. Estuarine Coastal Shelf Science 65, p. 557-570.
(online at: https://www.academia.edu/12325805/Variation_in_the_diversity_and_composition_of_benthic_taxa_as_a_function_of_distance_offshore_depth_and_exposure_in_the_Spermonde_Archipelago_Indonesia)

Cleary, D.F.R. & W. Renema (2007)- Relating species traits of foraminifera to environmental variables in the Spermonde Archipelago, Indonesia. Marine Ecology Progress Series 334, p. 73-82.

(Dinoflagellate symbionts and an orbitoidal chamber arrangement in foraminifera from SW Sulawesi waters linked to exposed reefs and hard substrate, whereas rhodophyte symbionts linked to sheltered reefs and sandy substrate. Etc.)

Cleary, D.F.R., Suharsono & B.W. Hoeksema (2006)- Coral diversity across a disturbance gradient in the Pulau Seribu reef complex off Jakarta, Indonesia. Biodiversity and Conservation, Springer, 15, p. 3653–3674.
(online at: www.researchgate.net/publication/302382186_Coral_diversity_across_a_disturbance Etc.)
(Study of coral richness, cover, and composition in Jakarta Bay and Pulau Seribu reef complex off Jakarta. Coral cover very low and composition differed markedly in near-shore zone 1 (Jakarta Bay), where human-induced disturbance is most intense, and highest in outlying reefs of Seribu islands)

Cole, A.R., P.T. Harris & J.B. Keene (1995)- Foraminifers as facies indicators in a tropical macrotidal environment; Torres Strait-Fly River delta, Papua New Guinea. In: B.W. Flemming & A. Bartholomae (eds.) Tidal signatures in modern and ancient sediments, Int. Association of Sedimentologists (IAS), Special Publ. 24, p. 213-223.

Collen, J.D. & D.W. Garton (2004)- Larger foraminifera and sedimentation around Fongafale Island, Funafuti Atoll, Tuvalu. Coral Reefs 23, 3, p. 445-454.
(Larger foraminifera common around Fongafale Island, Tuvalu. In shallow lagoon mainly larger foraminifera (Amphistegina lessonii, A. lobifera, Baculogypsina sphaerulata, Calcarina spengleri, Marginopora vertebralis, Sorites marginalis). In deeper water Halimeda replaces foraminifera)

Collins, A.C. (1958)- Foraminifera. In: Great Barrier Reef Expedition 1928-29, Scientific Reports, British Museum (Natural History), London, 6, 6, p. 335-437.

Coustillas, F. (1983)- Les facies recents de la plate-forme orientale de Kalimantan (Indonesie) et leur contenu micropaleontologique (foraminiferes benthiques). Doct. Thesis Universite de Bordeaux, France, p. 1-188.
('Recent facies of the eastern platform of Kalimantan and their micropaleontological content (benthic foraminifera)')

Culver, S. J., E. Leorri, D.R. Corbett, D.J. Mallinson, N.A.M. Shazili et al. (2013)- Infaunal mangrove swamp foraminifera in the Setiu wetland, Terengganu, Malaysia. J. Foraminiferal Research 43, 3, p. 262-279.
(Mangrove swamp samples from Setiu wetlands of NE Peninsular Malaysia with 13 infauna taxa. Six taxa live in all cores: Ammobaculites exiguus, Bruneica clypea, Caronia exilis, Haplophragmoides, Siphotrochammina and Trochammina inflata. Upper mangrove swamp populations with Arenoparella mexicana (=Trochammina inflata mexicana), Haplophragmoides wilberti, Miliammina fusca, Miliammina obliqua, Trochammina inflata and calcareous Helenina anderseni. Low-mangrove-swamp and adjacent mudflat populations dominated by calcareous species such as Ammonia aoteana, Rosalina sp., Elphidium oceanicum, and Triloculina oblonga)

Culver, S.J., D.J. Mallinson, D.R. Corbett, E. Leorri, A.A. Rouf, N.A.M. Shazili, R. Yaacob, J.E. Whittaker, M.A. Buzas & P.R. Parham (2012)- Distribution of foraminifera in the Setiu estuary and lagoon, Terengganu, Malaysia. J. Foraminiferal Research 42, p. 109-133.
(Four benthic foram thanatofacies in Setiu wetlands of NE Peninsular Malaysia, related to variations in salinity and hydrodynamics: (1) low salinity estuarine: low diversity assemblage dominated by Ammotium directum, Trochammina amnicola, Miliammina fusca and Ammobaculites exiguus; (2) medium salinity lagoon: dominated by Ammobaculites exiguus; (3) high salinity estuary and lagoon: high diversity, dominated by A. exiguus and Ammonia aff. A. aoteana; (4) normal marine salinity inlet and adjacent lagoon: high diversity dominated by Amphistegina lessonii, Ammonia aff. A. aoteana)

Cushman, J.A. (1917)- New species and varieties of foraminifera from the Philippines and adjacent waters: (Scientific Results of the Philippine Cruise of the Fisheries Steamer "Albatross" 1907-1910, No. 35). Proc. United States National Museum 51, p. 651-662.
(online at: https://repository.si.edu/handle/10088/15002)

(Brief descriptions of new species of deep water foraminifera from Albatross Expedition in Philippines. No figures)

Cushman, J.A. (1919)- The relationships of the genera *Calcarina, Tinoporus* and *Baculogypsina* as indicated by recent Philippine material. United States National Museum Bull. 100, 1, 4, p. 363-368.
(online at: https://repository.si.edu/handle/10088/21244)

Cushman, J.A. (1921)- Foraminifera of the Philippine and adjacent seas. United States National Museum Bull. 100, 4, p. 1-589.
(online at: https://repository.si.edu/handle/10088/21264)
(Extensive descriptions of Recent benthic and planktonic foraminifera from 600 shallow and deep water dredge samples collected during 'Albatross Expedition' around Philippines)

Cushman, J.A. (1924)- Samoan foraminifera. Publ. Carnegie Inst. Washington 342, 21, p. 1-75.

Cushman, J.A. (1932)- The foraminifera of the Tropical Pacific collections of the Albatross, 1899-1900, Part 1- Astrorhizidae to Trochamminidae. United States National Museum Bull. 161, p. 1-84.
(online at: https://repository.si.edu/handle/10088/10059)
(Part 1 of descriptions of Recent foraminifera in deep water samples from around Equatorial Pacific islands)

Cushman, J.A. (1933)- The foraminifera of the Tropical Pacific collections of the Albatross, 1899-1900, Part 2- Lagenidae to Alveolinellidae. United States National Museum Bull. 161, p. 1-79.
(online at: https://repository.si.edu/handle/10088/10058)

Cushman, J.A. (1942)- The foraminifera of the Tropical Pacific collections of the Albatross, 1899-1900, Part 3- Heterolicidae and Buliminidae. United States National Museum Bull. 161, p. 1-67.
(online at: https://repository.si.edu/handle/10088/10057)
(Descriptions of smaller bulimimid benthic foraminifera from tropical Pacific Ocean. For Part 4 of series see Todd 1965)

Cushman, J.A., R. Todd & R.J. Post (1954)- Recent foraminifera of the Marshall Islands: Bikini and nearby atolls, part 2, oceanography (biologic). U.S. Geological Survey (USGS) Prof Paper 260-H, p. 319-384.
(online at: http://pubs.usgs.gov/pp/0260h/report.pdf)
(331 species from 195 samples from lagoons and outer slopes of Marshall Islands: Rongerik, Rongelap, Bikini and Eniwetok, range in depth from beach to 835 fathoms. Reef flat fauna characterized by common Calcarina spengleri, Marginopora vertebralis, Homotrema rubrum, Miniacina miniacea, Carpenteria proteiformis and Amphistegina madagascariensis. Lagoon fauna dominated by A. madagascariensis and Heterostegina suborbicularis)

Dai, L., Q. Zhang, S.Y. Foong, Z. Cheng, C. Weng & L. Mao (2023)- Pollen morphology of selected tropical plants in Peninsular Malaysia and its implication in the paleoecological reconstruction of Southeast Asia. Review Palaeobotany Palynology 316, 104935, p. 1-53.
(Pollen morphology of 124 Recent species (111 genera) from tropical Malaysia. 19 pollen morphotypes, of which the tricolporate type occurs in half of samples. With ample illustrations)

Dalby, A.P., A. Kumar, J.M. Moore & R.T. Patterson (2000)- Preliminary survey of arcellaceans (Thecamoebians) as limnological indicators in tropical Lake Sentani, Irian Jaya, Indonesia. J. Foraminiferal Research 30, p. 135-142.
(On low diversity fresh water Thecamoebian amoebae assemblage in Lake Sentani, NE Papua)

Dawson, J.L., S.G. Smithers & Q. Hua (2014)- The importance of large benthic foraminifera to reef island sediment budget and dynamics at Raine Island, northern Great Barrier Reef. Geomorphology 222, p. 68-81.
(Larger foraminifera Baculogypsina sphaerulata, Marginopora and Amphistegina contribute 55% of calcareous sediment produced on Raine Island reef)

Dawson, S. (2007)- Diatom biostratigraphy of tsunami deposits: examples from the 1998 Papua New Guinea tsunami. Sedimentary Geol 200, 3-4, p. 328-335.
(Variable and often chaotic diatom assemblages can be attributed to tsunami waves incorporating and depositing diatoms from intertidal and offshore habitats during runup and subsequent backwash. Tsunami sand deposits have high % of broken diatom valves and dominance of centric (circular) species)

De, S. & A.K. Gupta (2010)- Deep-sea faunal provinces and their inferred environments in the Indian Ocean based on distribution of Recent benthic foraminifera. Palaeogeogr. Palaeoclim. Palaeoecology 291, p. 429-442.
(Distributions of 46 species of deep-sea benthic foraminifera from 131 core-top samples (322-5013 m) from across Indian Ocean. Two faunal provinces: (1) NW (Arabian Sea): with high organic flux and pronounced oxygen minimum zone (dominated by Uvigerina peregrina, Robulus nicobarensis, Bolivinita spp., Bulimina aculeata, Bulimina alazanensis, Ehrenbergina carinata and Cassidulina carinata); (2) S, SE and E Indian Ocean (dominated by Nuttallides umbonifera, Epistominella exigua, Globocassidulina subglobosa, Uvigerina proboscidea, Cibicides wuellerstorfi, Cassidulina laevigata, Pullenia bulloides, Oridorsalis umbonatus, Gyroidinoides soldanii), suggesting well-oxygenated, cold deep water)

Debenay, J.P. (2013)- A guide to 1,000 foraminifera from the Southwestern Pacific New Caledonia. IRD Editions, Montpellier, Publications Scientifiques du Museum National Histoire naturelle, Paris, p. 1-383.
(online at: http://horizon.documentation.ird.fr/exl-doc/pleins_textes/divers14-01/010058351.pdf)
(Descriptions and environmental conditions of 1000 species of Recent foraminifera found around New Caledonia)

Debenay, J.P. & G. Cabioch (2007)- Recent and Quaternary foraminifera collected around New Caledonia. In: Compendium of marine species of New Caledonia, Documments Scientifiques Techniques II-7, Institut de Recherche pour le Developpement (IRD), Noumea, New Caledonia, p. 69-94.
(online at: www.ird.nc/biodec/downloads/Compendium/Version%20v%E9rrouill%E9e/Debenay-Cabioch-v.pdf)
(Listing and some illustrations of modern forams around New Caledonia)

Debenay, J.P. & C.E. Payri (2007)- Epiphytic foraminiferal assemblages on macroalgae in reefal environments of New Caledonia. J. Foraminiferal Research 40, 1, p. 36-60.
(online at: http://jfr.geoscienceworld.org/content/40/1/36.full.pdf)
(152 species of epiphytic foraminifera identified from New Caledonia)

De Deckker, P. & F.X. Gingele (2002)- On the occurrence of the giant diatom *Ethmodiscus rex* in an 80-ka record from a deep-sea core, southeast of Sumatra, Indonesia: implications for tropical palaeoceanography. Marine Geology 183, p. 31-43.
(Deep-sea core from water depth 2542 m off SE Sumatra shows 'blooms' of giant diatom Ethmodiscus rex in Indian Ocean during last glacial period, particularly in Last Glacial Maximum. Blooms caused by increases in salinity and nitrate levels near surface. No major upwelling recorded during glacial times. During glacial period Indonesian Archipelago was much drier, preventing low-salinity 'cap' at surface of oceans)

De Neve, G.A. (1949)- Foraminifera from the shore zone of the Islands of Roeang and Siaoe. Chronica Naturae 105, 4, p. 113-115.
(online at: http://colonial.library.leiden.edu/….)
(Recent benthic foraminifera in shoreline sands of islands between Sulawesi and Philippines. Mainly miliolids, Robulus, Elphidium, Heterostegina, Amphistegina, Calcarina spp., Baculogypsina sphaerulata)

De Neve, G.A. (1949)- Opmerkingen over een foraminifenhoudende kalksteen van Baranti op het schiereiland Sanggar (Soembawa) en de strandzanden van Poeloe Ngali (Salehbaai) en Poeloe Madang. Chronica Naturae 105, 4, p. 116-118.
(online at: http://colonial.library.leiden.edu/….)

('Remarks on a foraminifera-bearing limestone of Baranti on the Sanggar Peninsula (Sumbawa) and the beach sands of Pulau Ngali (Saleh Bay) and Pulau Madang'. Late Neogene limestone similar to described by Tobler (1918), with Schlumbergerina, Alveolinella quoyi, Calcarina spp., Baculogypsina sphaerulata, etc. Baculogypsina tetraedra n.sp. described by Tobler (1918) from this area is probably synonym of Tinoporus baculatus and T. floresianus (Schlumberger))

De Neve, G.A. (1949)- Foraminifera from the shore zone of the islands of Morotai and Ternate. Chronica Naturae 105, 7, p. 196-198.
(online at: http://colonial.library.leiden.edu/....)
(Recent foraminifera from beach sand samples of SW Molotai and Ternate: Quinqueloculina, Robulus, Elphidium, Heterostegina, Orbitolites, Rotalia gaimardii, Amphistegina radiata, Calcarina spp., Baculogypsina sphaerulata. With summary of foraminifera from deep water samples off W Halmahera from 'Albatross-Expedition', as reported by Cushman (1921))

De Neve, G.A. (1949)- Foraminifera from the shore zone of Parigi and Poso (Gulf of Tomini). Chronica Naturae 105, 10, p. 252-254.
(online at: http://colonial.library.leiden.edu/....)
(Recent foraminifera from beach sand samples near Parigi and Poso, C Sulawesi. Up to 13 species, incl. miliolids, Elphidium, Operculina, Verbeekia, Orbitolites, Siderolites tetraeda and Calcarina. With summary of Gulf of Tomini deep water samples (1240-1966m) from 'Albatross-Expedition', as reported by Cushman (1921))

De Neve, G.A. (1949)- Een notitie over de foraminiferen van de oudste diepzee-lodingen in Indonesia. Chronica Naturae 105, 11, p. 291-292.
(online at: http://colonial.library.leiden.edu/....)
('Note on the foraminifera from the oldest deep sea soundings in Indonesia'. Brief note on earliest descriptions of deep sea foraminifera from Indonesia by Ehrenberg (1854) and Harting (1861, 1864))

De Silva, L.P. & P.J. Militante-Matias (1998)- Foraminiferal assemblages of Pagbilao Bay, Philippines. J. Geological Society of the Philippines 53, 1-2, p.
(134 foram species identified, representing 7 foraminiferal assemblages: river mouth, intertidal zone- Patayan Island, intertidal zone- Bocboc point, beach, reef/carbonate platform, inner bay, and inner bay channel. Nature of substrate strongly influences distribution of shallow water benthic foraminifera)

Dewi, K.T. (1993)- Ostracoda from the Java Sea, West of Bawean Island, Indonesia. Masters Thesis, University of Wollongong, Australia, p. 1-165.
(online at: https://ro.uow.edu.au/theses/2832)
(Seafloor samples from Sunda Shelf W of Bawean island contain 113 species of ostracodes, including 7 new)

Dewi, K.T. (1997)- Ostracoda from the Java Sea, West of Bawean Island, Indonesia. Marine Geology Institute (MGI), Bandung, Special Publ. 4, p. 1-86.
(same as Kresna Tri Dewi (1993) above; Masters Thesis, University of Wollongong, Australia)

Dewi, K.T. (2000)- Distribution of ostracoda from South of Tanjung Selatan, South Kalimantan. Bulletin of the Marine Geology 15, 1, p. 1-14.

Dewi, K.T. (2014)- Ostracoda from subsurface sediments of Karimata Strait as indicator of environmental changes. Bulletin of the Marine Geology 29, 1, p. 1-10.
(online at: ejournal.mgi.esdm.go.id/index.php/bomg/article/download/60/61)
(43 species of ostracods from 3 short seafloor cores of Sunda Shelf, in water depth 11-27m. Highest abundance/ diversity in upper 70cm of cores. Main genera Actinocythereis, Hemicytheridea, Loxoconcha, Neocytheretta, Stigmatocythere, Neomonoceratina, Phlyctenophora, Argillloecia, etc. (see also Mostafawi 1992))

Dewi, K.T., I. Adhirana, Y.A. Priohandono & L. Gustiantini (2016)- Ostracoda sebagai indikator perubahan lingkungan perairan sekitar PLTU Tarahan, Teluk Lampung, Sumatera. J. Geologi Kelautan 14, 1, p. 1-12.

(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/335/263)
('Ostracoda as indicators of marine environmental changes off the Tarahan power plant, Lampung Bay, Sumatera'. Ostracods from surface sediments in Lampung Bay quite diverse and abundant, with 27 genera, dominated by Keijella, Hemicytheridea and Cytherella. Also locally abundant Bairdopillata)

Dewi, K.T., L. Arifin, A. Yuningsih & Y. Permanawati (2012)- Meiofauna (Foraminifera) dalam sedimen dan keterkaitannya dengan pantai pasir putih Senggigi serta kondisi perairan Lombok Barat. J. Ilmu Teknologi Kelautan Tropis 4, 1, p. 47-54.
(online at: www.itk.fpik.ipb.ac.id/ej_itkt41/jurnal/...)
('Meiofauna (Foraminifera) in sediments and its association with the white sand beaches and water conditions of Senggigi, West Lombok'. Schlumbergerella floresiana abundant in white sands of Senggigi beach, but not common in offshore samples)

Dewi, K.T., N.C.D. Aryanto & Y. Noviadi (2007)- Land-sea interactions in coastal waters off NE Kalimantan: evidence from microfaunal communities. Bulletin of the Marine Geology 22, 1, p. 1-15.
(online at: ejournal.mgi.esdm.go.id/index.php/bomg/article/download/1/1)
(Microfauna in seafloor samples in 14-43m deep water off Nunukan and Sebatik islands in NE Kalimantan with typical microfauna of shallow marine ostracoda (Hemicytheridea spp., Keijella spp., Cytherella) and foraminifera (Asterorotalia trispinosa, Operculina))

Dewi, K.T., P. Frenzel & A. Muller (2008)- Mikrofauna (ostracoda) di sekitar paparan Sahul dan Laut Banda dalam kaitannya dengan batimetri. Proc. 37th Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 1, p. 127-138.
(online at: https://www.iagi.or.id/web/digital/14/2008_IAGI_Bandung_Mikrofauna-di-Sekitar-Paparan.pdf)
(Study of recent ostracods distribution in Snellius II samples along transects from Timor, Tanimbar, Seram into Banda Sea down to 3070m. Highest number of ostracods between 100-210m water depth, with Bairdopillata, Neonesidea, Paranesidea, Paracytheridea, Hemiparacytheridea, Foveoleberis, Polycope and Loxoconcha)

Dewi, K.T., P. Frenzel, A. Muller & D. van Harten (2004)- Recent ostracoda (microcrustacea) from a Banda-to-Timor Sea traverse: implications for paleobathymetric studies. Proc. 33rd Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 1, p. 42-51. *(online at: www.researchgate.net/profile/Peter-Frenzel/publication/255979013_Recent_Ostracoda_Microcrustac Etc.)*
(146 species of ostracods in surface sediments between 100-3070m along Banda Sea- Leti Strait-Timor Sea transect, E of Timor island. Richest assemblages in <100m water; abundance and diversity decreases with depth. Bathyal zone characterized by Cytheropteron, Saida, Bradleya, Paleocythere, Henryhowella, Krithe and Parakrithe; lower bathyal by Acantocythereis, Agrenocythereis and Krithe- Parakrithe spp.; abyssal depths by Argilloecia, Xestoleberis, Cytheropteron)

Dewi, K.T. & M. Hanafi (2013)- Karakteristik komunitas foraminifera laut dalam di Teluk Tomini, Sulawesi. J. Ilmu Teknologi Kelautan Tropis (JITKT; Bogor), 5, 1, p. 17-25.
(online at: http://journal.ipb.ac.id/index.php/jurnalikt/article/view/7742)
('The characteristics of the deep sea foraminiferal community in Tomini Bay, Sulawesi'. Foraminifera from 14 seafloor sediment samples from 600-1400m in Tomini Bay. Samples dominated by planktonic foraminifera (>90%). Benthic foraminifera represented by Cibicidoides wuellerstorfi, Ceratobulimina pacifica, Pyrgo sp., Bolivinita quadrilatera, Uvigerina peregrina, Planulina, etc., also very rare Laticarinina below 1000m)

Dewi, K.T. & D. Illahude (2005)- Ostracoda from off Derawan island, East Kalimantan (LP-1815) in relation to bathymetric zonation. Bulletin of the Marine Geology 20, 1, p. 1-14.
(Depth distribution of Recent ostracoda from 25 seafloor samples at depths 7-628m off Berau/ Tarakan Rivers, NE Kalimantan. 142 species identified. Common ostracods in area around reef complex, with Bairdopillata paracratericola, B. paraalcyanicola and Macrocypris decora. Also high diversity at water depths >200m, with displaced shallow water ostracods and deep water fauna. Bradleya common deepwater ostracod, found from <50m to >500m)

Dewi, K.T., A. Muller, P. Frenzel, L. Auliaherliaty & L. Gustiantini (2003)- Do Quaternary ostracods reflect sea level changes in the Timor Sea? Proc. 32nd Annual Conv. Indonesian Association Geologists (IAGI) and 28th Annual Conv. HAGI, Jakarta, p. 1-11.
(Samples from E of Timor in water depth 1768m show several small ostracod diversity peaks. Peaks of deep-sea taxa like Krithe, Bradleya, Cytheropteron, Acantocythereis and Ambocythere related to changes in paleoproductivity. Other peak with mixed shallow-water (Paracytheridea, Quadracythere, Loxocorniculum, Neonesidea) and deep-sea taxa suggest downslope transport of sediments, possibly regressive events)

Dewi, K.T., N. Nurdin, Y.A. Priohandono & A. Sinaga (2015)- Benthic foraminifera in marine sediment related to environmental changes off Bangka Island, Indonesia. Berita Sedimentologi 33, p. 47-57.
(online at: www.iagi.or.id/fosi/files/2015/09/BS33-Marine-Geology-of-Indonesia-II-R1.pdf)
(60 species of foraminifera in samples from 6-24m water depth offshore E Bangka island. Dominated by Amphistegina, Elphidium, Quinqueloculina, Operculina)

Dewi, K.T. & Y.A. Prihandono (2023)- Mikrofauna (Ostrakoda) dari Teluk Balikpapan; implikasinya untuk interpretasi lingkungan pengendapan. J. Geologi Kelautan 21, 1, p. 13-22.
(online at: https://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/779/585)
('Microfauna (Ostracoda) from Balikpapan Bay: its implication for interpreting depositional environments'. Ostracoda from water depths 1.5-27m 62 species, dominated by shallow marine ostracods Hemicytheridea reticulata, Cytherella semitalis and Alocopocythere kendengensis)

Dewi, K.T., Y.A. Prihandono & H.H. Prabowo (2000)- Ostracoda perairan Utara P. Kangean: kaitannya dengan transpor sedimen. Proc. 29th Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 4, p. 109-116.
('Ostracodes from waters North of Kangean Island; relations to sediment transport'. 30 sediment samples with 30 species, dominated by Bairdopillata, Cytherella, Foveoleberis. Not much detail)

Dewi, K.T. & E. Saputro (2013)- Sebaran spasial foraminifera dalam kaitannya dengan kedalaman laut dan jenis sedimen di Teluk Bone, Sulawesi Selatan. J. Geologi Kelautan 11, 3, p. 165-173.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/240/230)
('Spatial distribution of foraminifera in relation to water depth and sediment types in Bone Bay, S Sulawesi'. Foraminifera from 23-85m water depth in N Bone Bay mainly Operculina spp., Heterolepa, Pseudorotalia. Absence of 'Sunda species' Asterorotalia trispinosa)

Dewi, K.T., E. Setiyabudi, R. Oktavitania & H. Samodra (2023)- Tinjauan kontribusi fosil dalam penetapan warisan geologi. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 24, 4, p. 225-234.
(online at: https://jgsm.geologi.esdm.go.id/index.php/JGSM/article/view/804/547)
('An overview of fossil contributions to geoheritage determination'. Listing of 28 potential geoheritage sites across indonesia that contain unique Permian-Pleistocene fossil assemblages)

Dharma, B. (2005)- Recent & fossil Indonesian shells. Conch Books, Hackenheim, Germany, p. 1-424.
(Comprehensive monograph/atlas of Recent and fossil molluscs of Indonesia. 1,982 Recent species in 536 genera With 150 plates (plates 111-150 of fossil species)

Dhillon, D.S. (1968)- Notes on the foraminiferal sediments from the Lupar and Labuk estuaries, East Malaysia. In: Geological Papers 1967, Geological Survey Borneo Region, Malaysia, Kuching, Bull. 9, p. 56-73.
(Recent benthic foraminifera distribution in SW Sarawak river estuaries. 46 species. Arenaceous brackish water species found far upriver, while shallow marine hyaline calcareous species dominate near mouth of estuary. Includes new genus with two new species, Gobbettia wilfordi and Gobbettia klompei)

Dhillon, D. S. (1968)- A new genus of the Haplophragmoidinae from Malaysia. Contr. Cushman Foundation Foraminiferal Research 19, 4, p. 140-141.

Dhillon, D.S. (1969)- Three new species of foraminifera from Malaysia. Journal of Paleontology 43, 3, p. 767-770.

(Three new species of arenaceous foraminifera proposed from Recent sediments in Labuk Estuary, SW Sarawak: Arenoparrella malaysiana, Haplophragmoides labukensis, and H. sabahensis (also described in Dhillon (1968))

Ding, X., F. Bassinot, F. Guichard, Q.Y. Li, N.Q. Fang, L. Labeyrie, R.C. Xin, M.K. Adisaputra & K. Hardjawidjaksana (2006)- Distribution and ecology of planktonic foraminifera from the seas around the Indonesian Archipelago. Marine Micropaleontology 58, p. 114-134.
(Planktonic foraminiferal assemblages five provinces: (1) Banda/ Java region; (2) Timor region; (3) Java upwelling region; (4) Indian monsoon Sumatra region, and (5) NW Australia margin region. Assemblages reflect sea-surface temperature, salinity, thermocline depth, and nutrient supply, related to circulation patterns. Strongest dissolution in Java upwelling region, with lysocline rising above 2800m. Increase in Globigerina bulloides at 10-8 ka BP in Java upwelling region corresponds to decrease in Banda/ Java region, indicating intensification of upwelling in relation to strengthened SE monsoon)

Egger, J.G. (1893)- Foraminiferen aus Meeresgrundproben gelothet von 1874 bis 1876 von S.M. Sch. Gazelle. Abhandlungen Bayerischen Akademie Wissenschaften, Mathematisch-Physikalische Klasse 18, 2, p. 193-458.
(online at: https://www.biodiversitylibrary.org/item/110282#page/229/mode/1up)
('Foraminifera from sea floor samples collected between 1874 and 1876 by S.M. Sch. Gazelle'. Classic study of foraminifera from sea floor samples collected by German Gazelle oceanographic expedition. Incl. material from Banda Sea, Raja Ampat islands, etc. (stations 95-99))

Faiz, N.N., K.R. Mohamed & R. Omar (2017)- Species diversity and distribution of ostracoda (Crustacea) in Sungai Kubang Badak, Langkawi, Malaysia. Malayan Nature J. 69, 2, p. 83-92.
(online at: https://www.researchgate.net/publication/325557157_Species_diversity_and_distribution_of_Ostracoda_Crustacea_in_Sungai_Kubang_Badak_Langkawi_Kedah_Malaysia)
(Estuarine-shallow marine ostracode assemblages in Kubang Badak river and river mouth in N Langkawi. 27 species, dominated by Hemicytheridea wangi and Neomonoceratina delicata. Three assemblages. Density and diversity of ostracoda increases from upper estuary to nearshore areas)

Faiz, N.N., K.R. Mohamed, R. Omar & M.N.A. Malek (2017)- Benthic foraminiferal assemblages in the Pahang River Delta (Pahang, Malaysia). Thalassas: An International J. of Marine Sciences 34, 1, p. 65-79.
(Bottom sediments of Pahang River delta (E coast of Malay Peninsula) yielded 88 benthic foraminifera species. Agglutinated foraminifera (Ammobaculites) dominant at river mouth, while calcareous species (Asterorotalia pulchella) dominated in offshore samples)

Faiz, N.N., R. Omar & Basir Jasin (2007)- Taburan ostrakod di dalam sedimen luar pantai di sekitar Pulau Tinggi, Johor. Sains Malaysiana 36, 2, p. 139-148.
(online at: http://journalarticle.ukm.my/112/1/1.pdf)
(Distribution of ostracoda in offshore sediment around Pulau Tinggi, Johor. 11 shallow(?) marine samples off SE Malay Peninsula with 36 genera/ 51 species of ostracods. Dominant species is Loxoconcha malayensis. Sediment mainly m-grained sand)

Faiz, N.N., Ramlan Omar, M.N. Abd Malek, C. Li & Y. Liu (2016)- Taburan dan kepelbagaian Foraminifera bentik di dalam sedimen permukaan sekitar delta Sungai Pahang, Pahang, Malaysia. Sains Malaysiana 45, 5, p. 669-676.
(online at: www.ukm.my/jsm/pdf_files/SM-PDF-45-5-2016/02%20Noraswana%20.pdf)
('The distribution and diversity of benthic Foraminifera in surface sediment of Pahang River Delta, Pahang, Malaysia'. 82 species of Recent benthic foraminifera offshore Pahang River delta. Amphistegina lessonii and A. gibbosa most abundant, followed by Elphidium advenum, Operculina ammonoides and Asterorotalia pulchella. No specifics on water depth of samples or detailed distribution)

Fajemila, O.T., M.R. Langer & J.H. Lipps (2015)- Spatial patterns in the distribution, diversity and abundance of benthic foraminifera around Moorea (Society Archipelago, French Polynesia). PLoS One 10, 12, e0145752, p. 1-25.

(online at: http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0145752)
(380 species of shallow benthic foraminifera from around Moorea)

Fauzielly, L. (2013)- Distribusi vertikal Ostracoda dan hubungannya dengan perubahan lingkungan di perairan Teluk Jakarta. Bull. Scientific Contribution (UNPAD) 11, 2, p. 108-117.
(online at: http://jurnal.unpad.ac.id/bsc/article/view/8292/3839)
('Vertical distribution of ostracods and relationship with environmental changes in Jakarta Bay waters'. Similar to Fauzielly et al. (2012), below)

Fauzielly, L., T. Irizuki & Y. Sampei (2012)- Vertical changes of Recent ostrocode asemblages and environment in the inner part of Jakarta Bay, Indonesia. J. Coastal Development 16, p. 11-24.
(Ostracodes from sediment core from the inner part of Jakarta Bay. 53 species, dominated by Keijella carriei and Loxoconcha wrighti, which are common in areas with high organic carbon and nitrogen contents)

Fauzielly, L., T. Irizuki & Y. Sampei (2013)- Spatial distribution of Recent ostracode assemblages and depositional environments in Jakarta Bay, Indonesia, with relation to environmental factors. Paleontological Research (Palaeontological Society of Japan) 16, 4, p. 267-281.
(online at: https://bioone.org/journalArticle/Download?urlId=10.2517%2F1342-8144-16.4.267
(Recent ostracodes of Jakarta Bay 94 species, dominated by Keijella carriei, Hemicytheridea reticulata, Loxoconcha wrighti and Hemicytheridea ornata. Cytherella, Cytherelloidea, Neomonoceratina, and Pistocythereis also abundant. Three biofacies: (I) muddy bottom inner-middle bay with K. carriei, L. wrighti, and H. reticulata. Biofacies; (II) muddy bottom outer bay with H. reticulata, H. ornata and Cytherella spp. (III) sandy mud bottom outer bay high-diversity assemblages with Atjehella kingmai, Foveoleberis cypraeoides, Neomonoceratina bataviana and Pistocythereis cribriformis)

Fauzielly, L., L. Jurnaliah & A.H. Hamdani (2014)- Distribusi foraminifera bentonik sedimen paleo-tsunami letusan Gunung Krakatau berdasarkan data inti bor U-6 Di daerah Ujungkulon, Banten. Bull. Scientific Contribution (UNPAD) 12, 2, p. 84-91.
(online at: http://repository.unpad.ac.id/21772/1/Distribusi-foraminifera-bentonik-sedimen...)
('The distribution of benthic foraminifera in sedimentary paleo-tsunami eruption of Krakatoa based on data from core U-6 in the Ujungkulon area, Banten'. Marine microfossils in onland sediments may indicate paleo-tsunami deposits. Sand with coral debris at 90-120cm in core from Ujung Kulon recognized as tsunami deposit. With 55 species of benthic foraminifera, dominated by Streblus beccarii, Planulina wuellerstorfi, Hyalinea balthica, Bulimina marginata, Bolivina spathulata, Elphidium, Uvigerina peregrina Cushman. Origin of tsunami sediments is (outer) shelf-bathyal and inner shelf lagoon)

Fernando, A.G.S., A.M. Peleo-Alampay & M.G. Wiesner (2007)- Calcareous nannofossils in surface sediments of the eastern and western South China Sea. Marine Micropaleontology 66, p. 1-26.
(Calcareous nannofossils in surface sediments of S China Sea in water depths of 35–4345 m. In shallow/ nearshore waters dominant taxa are Emiliania huxleyi and Gephyrocapsa oceanica. In deeper portion of SCS, Florisphaera profunda dominates. Three assemblages recognized:(a) upwelling, (b) oceanic and (c) deep basin assemblages. F. profunda is relatively resistant to dissolution)

Fitriany, R., L. Jurnaliah & T.Y.W.M. Iskandarsyah (2019)- Small benthic foraminifera as an indicator of paleotsunami in Ujungkulon, Banten Province, Indonesia. J. Geological Sciences Applied Geology (UNPAD) 3, 1, p. 74-78.
(online at: http://jurnal.unpad.ac.id/gsag/article/view/22241/10591)
(55 species of small benthic foraminifera in paleotsunami deposits of the Laban Isthmus, Ujung Kulon Peninsula, SW Java, ~2m below surface. Dominant species Planulina wuellerstorfi, a deep marine species)

Flenley, J.R. (1979)- The Equatorial rain forest: a geological history. Butterworths London, p. 1-162.
(Includes review of SE Asia Cretaceous- Neogene evolution of pollen assemblages and chapter on Quaternary palynology and vegetation changes of SE Asia. In PNG vegetation zones moved downward ~1600m during

Pleistocene glacial episodes, suggesting 10°C decline in temperatures (later book with similar title by R.J. Morley (2000) Origin and evolution of tropical rain forests))

Flenley, J.R. (1998)- Tropical forests under the climates of the last 30,000 years. In: A. Markham (ed.) Potential impacts of climate change on tropical forest ecosystems, Climatic Change 39, Kluwer, p. 177-197.
(Incl. review of pollen diagrams from lowlands of SE Asia. Depression of altitudinal vegetation zones during Late Pleistocene Last Glacial Maximum suggests temperatures 5-10°C cooler than now)

Flenley, J.R. (1999)- Problems of the Quaternary on mountains of the Sunda-Sahul region. Quaternary Science Reviews 15, p. 549-555.
(In montane areas in Sunda-Sahul region Upper Montane rainforest appears to be absent in Late Pleistocene. Estimates of temperature lowering in Late Pleistocene strikingly greater in mountains than in lowlands)

Forderer, M. & M.R. Langer (2016)- Five new species and one new genus of recent miliolid foraminifera from Raja Ampat (West Papua, Indonesia). PeerJ 4:e2157; DOI 10.7717/peerj.2157, p. 1-20.
(online at: www.ncbi.nlm.nih.gov/pmc/articles/PMC4924127/pdf/peerj-04-2157.pdf)
(Shallow waters W of Waigeo Island with highly diverse assemblages of 455 species of benthic foraminifera (249 miliolid species). With new genus (Dentoplanispirinella) and 5 new species of miliolids (D. occulta, Miliolinella moia, M. undina, Triloculina kawea, Siphonaperta hallocki). New species relatively rare)

Forderer, M. & M.R. Langer (2018)- Atlas of benthic foraminifera from coral reefs of the Raja Ampat Archipelago (Irian Jaya, Indonesia). Micropaleontology 64, 1-2, p. 1-170.
(Exceptionally rich and diverse shallow marine benthic foram assemblages in Raja Ampat islands, W of Waigeo/ Batanta islands: 421 species, with miliolids highest species number. Incl. 35 species of symbiont-bearing larger foraminifera, amongst highest in modern oceans)

Forderer, M. & M.R. Langer (2019)- Exceptionally species-rich assemblages of modern larger benthic foraminifera from nearshore reefs in northern Palawan (Philippines). Revue de Micropaleontologie 65, 9, 100387, p. 1-25.
(44 larger foraminifera morphospecies from shallow nearshore habitats in N Palawan, among the highest reported anywhere (in Coral Triangle hotspot of biodiversity). Heterostegina depressa and Calcarina mayori most widely distributed taxa)

Forderer, M., D. Rodder & M.R. Langer (2018)- Patterns of species richness and the center of diversity in modern Indo-Pacifc larger foraminifera. Nature Scientific Reports 8, 8189, p. 1-9.
(online at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5974165/pdf/41598_2018_Article_26598.pdf)
(Species richness maps for Indo-Pacific larger foraminifera)

Forel, M.B. (2021)- Recent ostracods (Crustacea) from the southwestern coast of Peninsular Thailand (Satun Province), Andaman Sea. Revue de Micropaleontologie 72, 100526, p. 1-15.
(35 species of ostracods from shallow (<1.5 m) embayment near Mu Koh Phetra National Park, SW Peninsular Thailand, Andaman Sea. Most species well-known from Indo-Pacific area)

Frerichs, W.E. (1967)- Distribution and ecology of benthonic foraminifera in the sediments of the Andaman Sea. Ph.D. Thesis University Southern California, Los Angeles, p. 1-269.
(Recent foraminifera from seafloor samples across Andaman Sea, between 13-3778m depth. Number of forams increases with depth to 1800m; below this affected by dissolution of calcareous foraminifera. Planktonic number increases with depth and distance from shore. Planktonic assemblages from shelf sediments have globose chambers (Globigerina, Globigerinoides, Globigerinita); in bathyal deposits common Globorotalia and Sphaeroidinella. Radiolarian number increases with depth; not significant above middle bathyal depths. Benthic foraminifera five faunal provinces. Etc.)

Frerichs, W.E. (1970)- Distribution and ecology of benthonic foraminifera in the sediments of the Andaman Sea. Contr. Cushman Foundation Foraminiferal Research 21, 4, p. 123-147.

(online at: https://cushmanfoundation.allenpress.com/portals/_default/files/pubarchive/CCFFR/21ccffr4.pdf)
(In Andaman Sea shelf faunas characterized by species of Ammonia, Asterorotalia and Pseudorotalia. Upper bathyal faunas characterized by first appearance of Bolivina robusta. Change in ornamentation of Uvigerina at upper-middle bathyal boundary: costate species in outer shelf-upper bathyal; hispid species (U. auberiana) below 600m. Lower bathyal microfaunas calcareous forams dissolved, characterized by arenaceous Glomospira charoides and Karreriella apicularis. Etc.)

Frerichs, W.E. (1971)- Planktonic foraminifera in the sediments of the Andaman Sea. J. Foraminiferal Research 1, p. 1-14.
(Distribution of Recent planktonic foraminifera in sediments of Andaman Sea controlled by surface salinity, water depth, diagenesis and topography. Large volumes of fresh water in N Andaman Sea result in lowering of surface salinity and exclusion of planktonic forams. Depth zonation: Sphaeroidinella and some Globorotalia species indicative of bathyal depths)

Fujita, K. (2006)- Identification of coral reef environments based on foraminiferal death assemblages from Ishigaki Island, Okinawa, Japan. Proc. 10th International Coral Reef Symposium, Okinawa 2004, p. 528-535.
(Reef-flat foraminiferal assemblages characterized by dominant Calcarinidae, whereas fore-reef foraminiferal assemblages are characterized by various dominant species)

Fujita, K. & S. Kato (2011)- Distribution of gravel-sized empty tests of large benthic foraminifers as practical depositional indicators in tropical reef and shelf carbonate environments. Facies 57, 4, p. 525-541.
(Depth and spatial distributions of Large Benthic Forams in 39 surface sediment samples from W coast of Miyako Island (Ryukyu Islands, NW Pacific))

Fujita, K., Y. Osawa, H. Kayanne, Y. Ide & H. Yamano (2009)- Distribution and sediment production of large benthic foraminifers on reef flats of the Majuro Atoll, Marshall Islands. Coral Reefs 28, p. 29-45.
(Estimates of sediment production by Large Benthic Forams, mainly Calcarina and Amphistegina, on reef flats of Pacific atolls. Both live attached to seagrass/algae and most abundant on ocean reef flat (ORF) and in inter-island channel near windward, sparsely populated islands. Calcarina density higher on windward sides)

Fujita, K., M. Otomaru, P. Lopati, T. Hosono & H. Kayanne (2016)- Shell productivity of the large benthic foraminifer *Baculogypsina sphaerulata*, based on the population dynamics in a tropical reef environment. Coral Reefs 35, p. 317–326

Fujita, K., H. Shimoji & K. Nagai (2006)- Paleoenvironmental interpretations of Quaternary reef deposits based on comparisons of 10 selected modern and fossil larger foraminifera from the Ryukyu Islands, Japan. Island Arc 15, 4, p. 420-436.

Fukumoto, Y., X. Li, Y. Yasuda, M. Okamura, K. Yamada & K. Kashima (2015)- The Holocene environmental changes in southern Indonesia reconstructed from highland caldera lake sediment in Bali Island. Quaternary International 374, p. 15-33.
(Diatoms, pollen, geomagnetic and geochemical analyses on 3.6 m long core from Lake Buyan, Bali, representing ~8000 yrs sedimentation. Laminated sediments at 6.5-5.0 and 3.6-3.1 ka reflect drier climate)

Furio, E.F., R.V. Azanza, Y. Fukuyo & K. Matsuoka (2012)- Review of geographical distribution of dinoflagellate cysts in Southeast Asian coasts. Coastal Marine Science 35, 1, p. 20-33.
(online at: http://repository.dl.itc.u-tokyo.ac.jp/dspace/bitstream/2261/51680/1/CMS350104.pdf)
(Dinoflagellates play role in harmful agal blooms. Review of distribution in marine sediments shows generally 13-50 cyst types, Pyridinium bahamense var. compressum most common species in SE Asia waters)

Furio, E.F., K. Matsuoka, K. Mizushima, I. Baula, K.W. Chan, A. Puyong, D. Srivilai, B.R. Sidharta & Y. Fukuyo (2006)- Assemblages and geographical distribution of dinoflagellate cysts in surface sediments of coastal waters of Sabah, Malaysia. Coastal Marine Science 30, 1, p. 62-73.
(online at: http://repository.dl.itc.u-tokyo.ac.jp/dspace/bitstream/2261/5609/1/KJ00004354582.pdf)

(Dinoflagellate cysts relatively rare in sediments from Kota Kinabalu Bay and Tuaran Estuary. >25 species identified, mainly Spiniferites, Alexandrium, Lingulodinium, Polysphaeridium, etc.))

Gastaldo, R.A. (2012)- Taphonomic controls on the distribution of palynomorphs in tidally influenced coastal deltaic settings. Palaios 27, p. 798-810.
(Includes discussion of Recent palynomorph distribution in meso- to macrotidal Rajang River Delta, Sarawak. Mangrove pollen found throughout delta and alluvial plain sediments, as far as 75 km inland from mouth of rivers in frequencies of 5% of pollen spectra)

Gastaldo, R.A., W. Feng & J.R. Staub (1996)- Palynofacies patterns in channel deposits of the Rajang River and delta, Sarawak, East Malaysia. Palaios 11, 3, p. 266-279.
(Three organic matter facies assemblages in Rajang River delta sediments)

Gastaldo, R.A. & J.R. Staub (1997)- Water column and grab sample palynofacies assemblages from the Rajang River delta, Sarawak, East Malaysia. Palynology 21, p. 145-172.

Girard, E.B., Estradivari, S. Ferse, R.A.-Rappe, J. Jompa & W. Renema (2022)- Dynamics of large benthic foraminiferal assemblages: A tool to foreshadow reef degradation? Science of the Total Environment 811, 151396, p. 1-11.
(online at: https://www.sciencedirect.com/science/article/pii/S0048969721064743)
(In Spermonde Archipelago (S Sulawesi) temporal shift was observed in foraminiferal assemblages from coral-dominated (e.g, Amphistegina lobifera, Calcarina spengleri, Heterostegina depressa) to fleshy algae dominated (e.g. Neorotalia gaimardi, C. mayori))

Glenn, E.C. (1989)- Foraminifera and associated sedimentary constituents in Holocene and Miocene reefs of the Philippines and Indonesia. Ph.D. Thesis, University of Houston, p. 1-665. *(Unpublished)*

Glenn, E.C., J.W. McManus, P.M. Alino, L. Talaue, P. Alino & V. Banzon (1981)- Distributions of live foraminifers on a portion of Apo Reef, Mindoro, Philippines. Proc. 4th International Coral Reef Symposium, Manila 1981, 2, p. 775-780.
(Apo reef W of Mindoro in S China Sea with 10 physiographic zones. Sediment pockets in reef wall with Amphistegina lessonii, Marginopora vertebralis. Deep lagoon with Elphidium, Sorites marginalis, Peneroplis. Reef flat with Calcarina spp., etc.)

Glenn-Sullivan, E.C. & I. Evans (2001)- The effects of time-averaging and taphonomy on the identification of reefal sub-environments using larger foraminifera: Apo Reef, Mindoro, Philippines. Palaios 16, 4, p. 399-408.
(Foraminifera ~40% of sediment at small, isolated Apo Reef. Comparisons of live and dead assemblages show time-averaged assemblages, the product of taphonomic processes, more effective in delineating reefal sub-environments than do live assemblages. Robust calcarinids are in shallow seaward zones; free-living miliolids and small rotaliines in leeward zones. Planktonics and large thin rotaliines in fore reef)

Goeting, S., A. Briguglio, W. Eder, J. Hohenegger, A. Roslim & L. Kocsis (2018)- Depth distribution of modern larger benthic foraminifera offshore Brunei Darussalam. Micropaleontology 64, 4, p. 299-316.
(Recent larger foram distribution at 6 sites in 8-35m water off Brunei Darussalam (mainly sandy patches near shipwrecks, in area of mud-dominated seafloor without larger forams). High diversity assemblages, 16 species. Shallowest samples dominated by Calcarina defrancei and Neorotalia calcar, deepest samples by Amphistegina radiata, Operculina ammonoides and Calcarina mayori. Alveolinella quoii around 30m)

Goeting, S., V. Cosovic; A. Benedetti, F. Fiorini, L. Kocsis, A. Roslim & A. Briguglio (2022)- Diversity and depth distribution of modern benthic foraminifera offshore Brunei Darussalam. J. Foraminiferal Research 52, 3, p. 160-178.
(231 species of foraminifera identified from 8-63 m water depth off Brunei. Most abundant foraminiferal groups from reef and wreck samples are larger benthic foraminifera: especially Calcarinidae and Amphisteginidae. In muddier sites mainly smaller benthic forams with Pseudorotalia and Asterorotalia)

Goeting, S., F. Fiorini, A. Benedetti, L. Kocsis, A. Roslim, N. Zaini & A. Briguglio (2021)- Catalogue of modern smaller benthic foraminifera from offshore Brunei Darussalam. Palaeontographica, A. Palaeozoology-Stratigraphy, 318, 5-6, p. 129-223.
(207 species/ 88 genera pf modern smaller benthic foraminifera from 17 sites off Brunei, including proximal reef, muddy to sandy seafloors near sunken wrecks and two transects in muddy environments between 8-63 m)

Goeting, S., H.C. Lee, L. Kocsis, C. Baumgartner-Mora & D.J. Marshall (2023)- Diversity and distribution of the benthic foraminifera on the Brunei Shelf (Northwest Borneo): effect of seawater depth. Diversity 15, 8, 937, p. 1-23.
(online at: https://www.mdpi.com/1424-2818/15/8/937)
(Benthic forminifera from samples on Brunei shelf between 10m and shelf edge at-200m depth, with 99 species, belonging to 56 genera. General increase in species richness with increasing depth along Brunei shelf, with greatest richness observed at depth of 144 m. Excellent plates)

Gorog, A.J., M.H. Sinaga & M.D. Engstrom (2004)- Vicariance or dispersal? Historical biogeography of three Sunda shelf murine rodents (*Maxomys surifer, Leopoldamys sabanus* and *Maxomys whiteheadi*). Biological J. Linnean Society, London, 81, p. 91-109.
(online at: https://academic.oup.com/biolinnean/article/81/1/91/2639894)
(DNA tests of three rain-forest-restricted murine rodents of Borneo, Sumatra, Java, Malay Peninsula and Indochina do not support hypothesis of migrations enabled by Late Pleistocene land bridges/ rainforest corridors, but suggest older history of divergent evolution since Pliocene fragmentation of Sunda block)

Graham, J.J. & P.J. Militante (1959)- Recent foraminifera from the Puerto Galera area, northern Mindoro, Philippines. Stanford University Publications, Geological Sciences, 6, 2, p. 1-171.

Grand Pre, C.A. (2011)- The application of macro- and microfossils to identify paleoearthquakes in Sumatra, Indonesia and to characterize geomorphic and ecological succession on a marsh platform after Hurricane Isabel in North Carolina, USA. Ph.D. Thesis University of Pennsylvania, p. 1-179.
(Study of Early Holocene coseismic subsidence in Aceh. Buried mangrove soil horizons overlain with sharp contact by 5-20 cm thick sand that tapered landward, with intertidal and shallow marine foraminifera (Ammonia, Asterorotalia, Pararotalia, Quinqueloculina, etc.) and probably tsunami deposit. Sands overlain by 1-3m of silty clay with at base common Cerithidea cingulata, an opportunistic intertidal gastropod)

Gremmen, W.H.E. (1987)- Palynological evidence from Quaternary sediments in Southeast Asia, a review. Palaeohistoria 29, p. 77-84.
(online at: http://ugp.rug.nl/Palaeohistoria/article/view/24871/22319)

Gremmen, W.H.E. (1989)- Palynological investigations in the Danau Tempe Depression, Southwest Sulawesi (Celebes), Indonesia. Modern Quaternary Research in Southeast Asia, Balkema, Rotterdam, 11, p. 123-134.
(online at: http://www.oxis.org/articles-c-j/gremmen-1990.pdf)
(Tempe Lake Depression in SW Sulawesi with three lakes and swamp areas probably remnant shallow marine seaway between Makassar Straits and Bone Bay until recently. In tidal mangrove zone between 7000- 2600 BP)

Guballa, J.D.S. & A.M. Peleo-Alampay (2020)- Pleistocene calcareous nannofossil biostratigraphy and Gephyrocapsid occurrence in Site U1431D, IODP 349, South China Sea. Geosciences (MDPI) 10, 388, p. 1-26.
(online at: https://www.mdpi.com/2076-3263/10/10/388)
(Pleistocene calcareous nannofossil biostratigraphy of IODP Site U1431D in eastern (off Philippines) part of S China Sea. Twelve Pleistocene datums identified. Last occurrence of Calcidiscus macintyrei is below first occurrence of large Gephyrocapsa spp. Etc.)

Guptha, M.V.S. (1981)- Nannoplankton from Recent sediments off the Andaman Islands. Indian J. Marine Sciences 10, p. 293-295.
(online at: http://nopr.niscair.res.in/bitstream/123456789/39098/1/IJMS%2010(3)%20293-295.pdf)

(16 deep marine seafloor samples around Little Andaman, with 14 modern and 24 reworked Eocene-Pliocene nannofossil species. Modern species dominated by Gephyrocapsa oceanica)

Gustiantini, L., K.T. Dewi & D. Illahude (2005)- Perbandingan foraminifera bentik dan plangtonik (P/B ratio) di perairan sekitar Pulau Derawan, Kalimantan Timur. Proc. Joint Convention 30th HAGI- 34th IAGI and 14th PERHAPI, Surabaya, p. 341-348.
('Comparison of benthic and planktonic foraminifera (P/B ratio) in the waters around Derawan Island, East Kalimantan')

Gustiantini, L., K.T. Dewi, A. Muller & Praptisih (2003)- The benthic foraminifera *Ammonia beccarii* as indicator of estuarine environments in Indonesia (Segara Anakan lagoon and southern Gombong, Java). Proc. 32nd Annual Conv. IAGI and 28th Annual Conv. HAGI, Jakarta, p. 1-11.
(Fossil estuarine deposits often with common Ammonia beccarii, abundance is low in modern Segara Anakan lagoon N of Nusakambangan, S Java)

Gustiantini, L., K.T. Dewi & E. Usman (2005)- Foraminifera di perairan sekitar Bakauheni, Lampung (Selat Sunda bagian utara). J. Geologi Kelautan 3, 1, p. 10-18.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/120/110)
('Foraminifera in the waters around Bakauheni, Lampung (northern Sunda Strait)' Abundant foraminifera along Lampung coast in N Sunda Strait, dominated by Asterorotalia trispinosa, Operculina, Pseudorotalia and Elphidium)

Gustiantini, L. & D. Ilahude (2015)- Foraminifera bentik dalam sedimen sebagai indikator kondisi lingkunan terumbu karang di perairan Pulau Cemara Besar dan Cemara Kecil, Kepulauan Karimunjawa, Jawa Tengah. J. Geologi Kelautan 10, 1, p. 35-38
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/213/203)
('Benthic foraminifera in sediment as indicators of coral reef environments in the waters of Pulau Cemara Besar and Cemara Kecil, Karimunjawa Islands, Central Java'. Benthic forams dominated by Calcarina, Amphistegina, Streblus and Reusella)

Gustiantini, L., K.A. Maryunani, R. Zuraida, C. Kissel, F. Bassinot & Y. Zaim (2015)- Distribusi foraminifera di Laut Halmahera dari Glasial Akhir sampai Resen. J. Geologi Kelautan 13, 1, p. 25-36.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/259/249)
('The distribution of foraminifera in the Halmahera Sea from the last Glacial to Recent'. Deep marine Late Pleistocene- Holocene planktonic and benthic foraminifera from core MD10-3339, SE of Halmahera)

Gustiantini, L., S.A. Piranti, R. Zuraida, S. Hyun, D.A.S. Ranawijaya & F.X. Harkinz (2018)- Foraminiferal analysis related to paleoceanographic changes of Arafura Sea and surrounding during Holocene. Bulletin of the Marine Geology 33, 2, p. 105-118.
(online at: https://ejournal.mgi.esdm.go.id/index.php/bomg/article/viewFile/571/446)
(Holocene foraminifera in shallow (152 cm) core from 276 m water depth on Arafura Shelf. Planktonics average 53%. Benthic forams dominated Bolivina, Bulimina and Uvigerina)

Gustiantini, L. & E. Usman (2008)- Distribusi foraminifera bentik sebagai indikator kondisi lingkungan di perairan sekitar Pulau Batam- Riau kepulauan. J. Geologi Kelautan 6, 1, p. 43-52.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/149/139)
('Distribution of benthic foraminifera as indicators of environmental conditions in the waters around Batam Island, Riau archipelago'. High abundance of benthic foraminifera in Batam- Bintan waters, dominated by dominated by Asterorotalia trispinosa, Pseudorotalia annectens, Amphistegina radiata, Quinqueloculina cf. philippinensis, and Operculina ammonoides)

Hada, Y. (1943)- The relation between the foraminifera and deposits of the Java Sea. J. Oceanographic Society of Japan 12, 4, p. 27-36. *(in Japanese with English Abstract)*
(online at: https://www.jstage.jst.go.jp/article/kaiyou1942/2/4/2_4_27/_pdf/-char/en)

(Bottom samples from 31 stations between 21-153m in Java Sea rich in foraminifera. 43 species. Some genera more abundant in sandy deposits (incl. Textularia, Operculina, Amphistegina, Siderolites (=Baculogypsina?; JTvG), Planorbulinella, Alveolinella), some more abundant in muddy deposits (Eponides praecinctus, Rotalia schroeteriana, Quinqueloculina))

Hadiwisastra, S. (1978)- Kumpulan Ostrakoda dari delta Cimanuk. J. RISET Geologi dan Pertambangan (LIPI) 1, 2, p. 9-20.
(online at: http://pustaka.geotek.lipi.go.id/wp-content/uploads/2016/02/Riset-Vol.1-No.2-78.pdf
('Ostracodes from the Cimanuk Delta'. 35 Recent genera of ostracods in shallow waters around Cimanuk 'birdfoot delta', Java. Delta fauna dominated by Hemicytheridea, Loxoconcha, Hemikrithe, Cythereis, etc. Open marine genera Cytherella, Cytherelloidea, Bradleya, etc.)

Hadiwisastra, S. & S. Djoehanah (1979)- Penyebaran foraminifera bentos di delta Cimanuk. J. RISET Geologi dan Pertambangan (LIPI) 2, 1, p. 7-21.
(online at: http://pustaka.geotek.lipi.go.id/wp-content/uploads/2016/02/Riset-Vol.2-No.1-2-2-.pdf)
('The distribution of benthic foraminifera in the Cimanuk delta'. 137 species of benthic foraminifera in 33 bottom samples around Cimanuk Delta, N coast of Java. Four main assemblages)

Haig, D.W. (1979)- Foraminiferids from shoreline sediments, Motupore Islands, Papua New Guinea. Science in New Guinea 6, 3, p. 138-143.

Haig, D.W. (1988)- Distribution of miliolid foraminifera in marine sediments around Motupore Island, Papua New Guinea. Science in New Guinea 14, p. 54-94.

Haig, D.W. (1988)- Miliolid foraminifera from inner neritic sand and mud facies of the Papuan lagoon, New Guinea. J. Foraminiferal Research 18, 3, p. 203-236.
(online at: http://jfr.geoscienceworld.org/content/18/3/203.full.pdf)
(101 species of miliolids from five settings in Papuan Lagoon, SE coast of PNG. Miliolids generally 10-40% of total foram assemblage, which is dominated by rotaliids)

Haig, D.W. (1993)- Buliminid foraminifera from inner neritic sand and mud facies of the Papuan Lagoon, New Guinea. J. Foraminiferal Research 23, 3, p. 162-179.
(online at: http://jfr.geoscienceworld.org/content/23/3/162.full.pdf)
(60 small buliminid-bolivinid-uvigerinid species from in 0-50m water depth in lagoon behind Papuan barrier reef, SE coast of PNG)

Haig, D.W. (1997)- Foraminifera from Exmouth Gulf, Western Australia. J. Royal Society Western Australia 80, p. 263-280.
(Recent foraminifera from 5-30m water depth at Exmouth Gulf: 236 benthic and 6 planktonic species)

Haig, D.W. & S. Burgin (1982)- Brackish-water foraminiferids from the Purari River delta, Papua New Guinea. Revista Espanola Micropaleontologia 14, p. 359-366.

Hallock, P. (1981)- Production of carbonate sediments by selected large benthic foraminifera on two Pacific coral reefs. Journal of Sedimentary Research 51, p. 467-474.
(Carbonate production rates by foram families Asterigerinidae, Calcarinidae, and Nummulitidae in Palau, W Caroline Islands: seaward reef flats up to 2.8 kg CaCO3/m2/yr, equivalent to deposition of almost 1 mm/yr. Productivity on lagoonal reef slopes about one-fifth. In Hawaii production rates much lower because of slower growth rates and absence of family Calcarinidae)

Hallock, P. (1984)- Distribution of selected species of living algal symbiont-bearing foraminifera on two Pacific coral reefs. J. Foraminiferal Research 14, p. 250-261.
(Distribution of 15 larger foram species on reefs of Palau (W Caroline Islands) and Oahu (Hawaii). Four clusters (1) Calcarinidae-dominated on seaward reef flats; (2) Marginopora vertebralis, Amphistegina lobifera

and Peneroplis planatus in protected shoals in <5m water, (3) Amphistegina lessonii on reef slopes of 5-20m, (4) Amphistegina radiata and Operculina ammonoides deeper dwelling taxa)

Hallock, P. (1999)- Symbiont-bearing foraminifera. In: B.K. Sen Gupta (ed.) Modern Foraminifera, Kluwer, Amsterdam, p. 123-149.

Hallock, P. & E.C. Glenn (1985)- Numerical analysis of foraminiferal assemblages: a tool for recognizing depositional facies in Lower Miocene reef complexes. Journal of Paleontology 59, 6, p. 1382-1394.
(late Early Miocene larger foram facies assemblages in wells Matinloc 2 and Libro 1, off NW Palawan, Philippines. Assigned to zone Te5, but more likely Lower Tf?; associated with N8 planktonic forams)

Hamilton, R., T. Hall, J. Stevenson & D. Penny (2019)- Distinguishing the pollen of Dipterocarpaceae from the seasonally dry and moist tropics of south-east Asia using light microscopy. Review Palaeobotany Palynology 263, p. 117-133.
(Review of modern pollen of Dipterocarpaceae family of large trees, which are common in forests across Indonesia and surrounding SE Asia (Dipterocarpus, Hopea, Shorea, Vatica, etc.))

Hanai, T., N. Ikeya & M. Yajima (1980)- Checklist of Ostracoda from Southeast Asia. University Museum, University of Tokyo, Bull. 17, p. 1-236.
(online at: www.um.u-tokyo.ac.jp/publish_db/Bulletin/no17/no17000.html)
(Review of studies on extensive listing of Recent and fossil ostracodes described from SE Asia)

Hanzawa, S. (1951)- Recent and fossil *Cycloclypeus* from the Ryukyu Islands and their adjacent seas. Short Papers Inst. Geology Paleontology, Tohoku University, Sendai, 3, p. 1-12.
(14 samples with Recent Cycloclypeus from off Ryukyu Islands from 87-133m water depth, one from 235m)

Hardy, M.J. (2000)- Origin, distribution, and degradation of sedimentary organic matter in a modern tropical deltaic system (Mahakam Delta, Borneo, Indonesia). Ph.D. Thesis Louisiana State University, Baton Rouge, p. 1-368. *(Unpublished)*

Hardy, M.J. & J.H. Wrenn (2009)- Palynomorph distribution in modern tropical deltaic and shelf sediments-Mahakam Delta, Borneo, Indonesia. Palynology 34, p. 19-42.
(Distribution of terrestrial palynomorphs in Mahakam Delta surface sediments from 12 depositional environments from head of delta to shelf edge can be explained by transport and depositional processes. Amounts of marine palynomorphs (foram linings, copepod eggs, dinocysts) increases gradually offshore)

Harting, P. (1863)- Bijdrage tot de kennis der mikroskopische fauna en flora van de Banda-Zee. Verhandelingen Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 10, p. 1-34.
('Contribution to the knowledge of the microscopic fauna and flora from the Banda Sea' Early report on foraminifera, radiolaria, etc. from Recent deep Banda Sea sediment samples between 1200-4000 fathoms)

Hasan, S.S., M. Mohamed, N. Muhsin & S. Jirin (2013)- The distribution of *Miliammina fusca* in three different environmental setting of Peninsular Malaysia, Malaysia. Proc. Petroleum Geoscience Conference Exhib. (PGCE 2013), Kuala Lumpur, P14, p. 1-4. *(Extended Abstract)*
(Study on distribution of small agglutinated benthic foram Miliammina fusca in three modern depositional settings, Klang Delta, Pahang Delta and Sedili Besar River. Invariably associated with brackish conditions. In Malay Basin M. fusca used as indicator of marine incursions into basin, particularly in Lower Oligocene (upper Group L and Group K))

Haseldonckx, P. (1974)- A palynological interpretation of palaeo-environments in S.E. Asia. Sains Malaysiana 3, 2, p. 119-127.

Haseldonckx, P. (1977)- The palynology of a Holocene marginal peat swamp environment in Johore, Malaysia. Review Palaeobotany Palynology 24, 5, p. 227-238.

(Shallow Holocene peat near Pekan Nanas, Johore, with 47 pollen and spore types. Pollen profile shows succession from open swamp vegetation with mangrove influence to marginal peat swamp facies with river bank vegetation. Radiocarbon dating on deepest peat yielded ~4.9 ka)

Haseldonckx, P. (1977)- Palynology and its application to Quaternary geology in the Sunda shelf region. Proc. Symposium on Quaternary Geology of the Malay-Indonesian coastal and offshore areas, Kuala Lumpur, 1976, United Nations ESCAP, CCOP, Technical Publication 5, p. 33-53.
(Brief, general overview of pollen and spores, processing and application in determination of paleoenvironments. 'Palynology in SE Asia still at early stage of development')

Haslett, S.K. (2001)- The palaeoenvironmental implications of the distribution of intertidal foraminifera in a tropical Australian estuary: a reconnaissance study. Australian Geographical Studies 39, 1, p. 67-74.
(Modern intertidal foraminifera in mangrove-lined microtidal distributary of Barron River Delta (Queensland): (1) saltmarsh: Trochammina inflata assemblage; (2) regularly inundated tidal flat: dominated by Ammonia beccarii; (3a) high tidal flat: >70% Ammonia beccarii and low diversity, and (3b) low tidal flat: 55-65% Ammonia beccarii and diverse small allochthonous species transported into estuary from shelf)

Hayward, B.W. & S. Kawagata (2005)- Extinct foraminifera figured in Brady's Challenger Report. J. Micropalaeontology 24, 2, p. 171-175.
(online at: https://www.j-micropalaeontol.net/24/171/2005/jm-24-171-2005.pdf)
(Brady's classic 1884 monograph on Recent foraminifera from Challenger Expedition samples contains 18 species known to become extinct in M Pleistocene, between 1.2- 0.6 Ma (mainly elongate, benthic foraminifera like Pleurostomella, Stilostomella, Orthomorphina, etc.). Majority (14 species) from two stations off Kei Islands, Banda Sea (191A, 192). Station 192 (~250m depth) is considerably shallower than established fossil bathymetric ranges of extinct species, suggesting tectonic uplift and Pleistocene deposits exposed on seafloor)

Hayward, B.W., S. Kawagata, H.R. Grenfell, A.T. Sabaa & T. O'Neill (2007)- Last global extinction in the deep sea during the mid-Pleistocene climate transition. Paleoceanography 22, PA3103, p. 1-14.
(20% of cosmopolitan deep-sea benthic foraminifera extinct during late Pliocene-M Pleistocene, with peak of extinctions during M Pleistocene Climate Transition (1.2-0.55 Ma). Family Stilostomellidae (30 species) wiped out, Pleurostomellidae (9 species) decimated. Pulsed declines in abundance, earlier in deepst water sites. Tied to demise of microbial food source due to increased cold and oxygenation of S-sourced deep water masses during major late Pliocene and E Pleistocene glacials)

Hendrizan, M., R.A. Troa, R. Zuraida & E. Triarso (2016)- Calcareous nannoplankton (marine algae) analysis in subsurface sediments of Andaman Sea. Bulletin of the Marine Geology 31, 2, p. 91-98.
(online at: http://ejournal.mgi.esdm.go.id/index.php/bomg/article/view/313/277)
(4m thick Quaternary clay sediment core BS36 in S Andaman Sea (Mergui-N Sumatra Basin). With 11 genera of calcareous nannofossils, dominated by Gephyrocapsa, Emiliania, and Helicosphaera)

Hendrizan, M., C.A. Widiyanti, R.E. Prabowo, Munasri & N. Nurdin (2019)- Karakter masa air di Laut Sulawesi berdasarkan analisis foraminifera kuantitatif. J. Geologi Kelautan, 17, 1, p. 9-18.\
(online at: https://www.researchgate.net/publication/337979247_KARAKTER_MASA_AIR_DI_LAUT_SULAWESI_BERDASARKAN_ANALISIS_FORAMINIFERA_KUANTITATIF)
('Water mass characteristics of the Sulawesi Sea based on quantitative foraminifera analysis'. Foraminifera from core STA 3 (water depth 294 m) on W side of North Makassar Straits)

Hessler, I., M. Young, U. Holzwarth, M. Mohtadi, A. Luckge & H. Behling (2013)- Imprint of eastern Indian Ocean surface oceanography on modern organic-walled dinoflagellate cyst assemblages. Marine Micropaleontology 101, p. 89-105.
(Recent distribution of dinoflagellate cysts in 116 marine surface samples of E Indian Ocean. Three distinct regions (1) W and E Indonesia, with high T and low nutrient content of surface water; dominated by Spiniferites spp. and S. ramosus; (2) Indonesian Throughflow (ITF) region, dominated by heterotrophic dinocyst species

(Brigantedinium spp.), reflecting high productivity; (3) offshore NW Australia, characterized by water masses of saline and nutrient depleted Leeuwin Current, with rel. high O. centrocarpum)

Hillen, R. (1986)- Palynology as a tool in delineating tropical lowland depositional environments of Late Quaternary age. In: G.H. Teh & S. Paramananthan (eds.) Proc. 5th Regional Congress Geology Mineral Energy Resources of SE Asia (GEOSEA V), Kuala Lumpur 1984, 1, Bull. Geological Society Malaysia 19, p. 495-504.
(online at: https://gsmpubl.files.wordpress.com/2014/09/bgsm1986037.pdf)
(Samples from Lower Perak and Kelantan lowlands (Malay Peninsula) allowed palynological characterizaton of environments:shallow offshore, deltaic/ esturine, mangrove, fresh water swamp, peat swamp)

Hofker, J. (1927)- The foraminifera from the Siboga expedition, I. Families Tinoporidae, Rotaliidae, Nummulitidae, Amphisteginidae. Siboga Expedition 1899-1900, Monograph 4a, 1, E.J. Brill, Leiden, p. 1-78.
(Modern foraminifera in seafloor samples collected by Siboga Expedition in Indonesia, part 1 of 3)

Hofker, J. (1930)- Foraminifera of the Siboga Expedition, Part II, Families Astrorhizidae, Rhizamminidae, Reophacidae, Anomalinidae, Peneroplidae. Siboga-Expeditie, Monograph 4a, 2, E.J. Brill, Leiden, p. 79-170.
(Modern foraminifera in seafloor samples collected by Siboga Expedition in Indonesia, part 2 of 3)

Hofker, J. (1933)- Foraminifera of the Malay Archipelago. Papers from Dr. Th. Mortensen's Pacific Expedition, 1914-16, part 62, Dansk Naturhistorisk Forening, Videnskabelige Meddedelser 93, C.A. Reitzel, p. 71-167.
(online at: https://www.delpher.nl/nl/boeken/view?identifier=MMKB24:064959000:00005)
(Descriptions of 55 species of Recent foraminifera collected by Th. Mortensen during Danish Expedition to Kai islands in 1922 (should be 1914-1916?). No maps, nothing on sample localities, little on depth ranges, etc.)

Hofker, J. (1951)- The Foraminifera of the Siboga Expedition. Part III. Ordo Dentata, Sub-Ordines Protoforaminata, Biforaminata, Deuteroforaminata. Siboga Expeditie Monograph 4a, 3, Brill, Leiden, p. 1-513.
(Modern foraminifera in seafloor samples collected by Siboga Expedition in Indonesia, part 3 of 3)

Hofker, J. (1968)- Foraminifera from the Bay of Jakarta, Java. Bijdragen tot de Dierkunde 37, p. 11-59.
(online at: www.repository.naturalis.nl/document/548150)
(Rel. low diversity recent foraminifera assemblages in shallow marine Jakarta Bay, dominated by Pseudoratalia schroeteriana, Asterorotalia pulchella, Elphidium batavum, Operculina complanata, Quinqueloculina. Highest diversity in NW part of bay. Pseudorotia most common in muddy substrates in SE, Baculogypsinoides and miliolids mainly in NW, away from delta muds)

Hofker, J. (1978)- Biological results of the Snellius Expedition XXX. The foraminifera collected in 1929 and 1930 in the eastern part of the Indonesian Archipelago. Zoologische Verhandelingen, Rijksmuseum Natuurlijke Historie Leiden 161, p. 1-69.
(online at: www.repository.naturalis.nl/document/155300)
(462 species of foraminifera from 78 sea bottom samples at depths 85- 5138m, collected by 1929-1930 Snellius Expedition to East Indonesia, and a few other shallow marine samples)

Hohenegger, J. (1994)- Distribution of living larger foraminifera NW of Sesoko-Jima, Okinawa, Japan. Marine Ecology 15, p. 291-334.
(Living larger foraminifer restricted to photic zone. Peneroplids in shallow-water from intertidal (common Peneroplis) to 40m (Dendritina, sandy substrates). Soritids subtidal, from reef moat down to 60m. Amphisorus and Marginopora common down to 3 m, Parasorites in deeper parts of reef slope. Alveolinella in upper 40m of reef slope. Amphisteginids in entire photic zone, with test flattening with increasing depth. Calcarinids cling to firm substrates. Baculogypsina restricted to high energy reef flat regions. Calcarina similar facies, extending to fore reefs down to 8m. Baculogypsinoides dominant calcarinid genus from 30-70m depth. Heterostegina prefers hard substrates on reef slope. Operculina frequent in deeper part, independent of substrates. Sandy bottoms from 30-60m inhabited by Nummulites (Operculina) venosus. Cycloclypeus restricted to fore reef areas below 50m down to base of photic zone)

Hohenegger, J. (1995)- Depth estimation by proportions of living larger foraminifera. Marine Micropaleontology 26, p. 31-47.

Hohenegger, J. (1996)- Remarks on the distribution of larger foraminifera (Protozoa) from Belau (Western Carolines). Kagoshima University Research Center Pacific Islands, Occasional Papers 30, p. 85-90.
(online at: http://hdl.handle.net/10232/16891)

Hohenegger, J. (1999)- Larger foraminifera-microscopical greenhouses indicating shallow-water tropical and subtropical environments in the present and past. Kagoshima University Research Center Pacific Islands, Occasional Papers 32, p. 19-45.
(online at: http://hdl.handle.net/10232/16923)
(Larger foraminifera (2mm-15cm) characteristic of shallow marine (sub-)tropical environments, in clear, nutrition-depleted water like for coral reefs. All larger foraminifera house symbiotic microalgae and are thus restricted to photic zone (<150m). In intertidal-shallow subtidal environments high irradiatiois blocked by thicker tests or porcelaneous structures; species living near base of photic zone facilitate light penetration by thin transparent test walls and by light-collecting mechanisms such as nodes and pillars)

Hohenegger, J. (2000)- Coenoclines of larger foraminifera. Micropaleontology 46, Suppl. 1, Advances in the biology of foraminifera, p. 127-151.
(online at: https://www.researchgate.net/publication/279548754_Coenoclines_of_Larger_Foraminifera)
(Good review of depth distribution of modern larger foraminifera, mainly in W Pacific)

Hohenegger, J. (2004)- Depth coenoclines and environmental considerations of Western Pacific larger foraminifera. J. Foraminiferal Research 34, p. 9-33.
(Good overview of depth distribution modern larger foraminifera in W Pacific)

Hohenegger, J. (2005)- Estimation of environmental paleogradient values based on presence/absence data: a case study using benthic foraminifera for paleodepth estimation. Palaeogeogr. Palaeoclim. Palaeoecology 17, p. 115-130.

Hohenegger, J. (2006)- The importance of symbiont-bearing benthic foraminifera for West Pacific carbonate beach environments. Marine Micropaleontology 61, 1, p. 4-39.
(online at: https://www.researchgate.net/publication/222019398_The_importance_of_symbiont-bearing_foraminifera_for_West_Pacific_carbonate_beach_environments)
(In tropical W Pacific and East Indian Ocean, symbiont-bearing benthic foraminifera are significant producers of calcium carbonate grains. Prefer firm substrate in high-energy environments close to front of coral reefs)

Hohenegger, J. (2011)- Large foraminifera: greenhouse constructions and gardeners in the oceanic microcosm; Kagoshima University Museum Bull. 5, Kagoshima, Japan, p. 1-81.
(online at: https://www.museum.kagoshima-u.ac.jp/publications/LARGE%20FORAMINIFERA.pdf)
(Well-illustrated general introduction to larger foraminifera and their symbionts)

Hohenegger, J. & E. Yordanova (2001)- Depth-transport functions and erosion-deposition diagrams as indicators of slope inclination and time-averaged traction forces: applications in tropical reef environments. Sedimentology 48, p. 1025-1046.
(Comparisons of distributions of living versus dead tests of larger foraminifera indicate common downslope transport in two NW Pacific off-reef transects)

Hohenegger, J. & E. Yordanova (2001)- Displacement of larger foraminifera at the western slope of Motobu Peninsula (Okinawa, Japan). Palaios 16, p. 53-72.

Hohenegger, J., E. Yordanova & A. Hatta (2000)- Remarks on West Pacific Nummulitidae (Foraminifera). J. Foraminiferal Research 30, p. 3-28.

(Operculina, Planostegina, Cycloclypeus and Heterostegina from W Pacific. Heterostegina depressa broad range in light intensities and is protected against irradiation by thick tests and cryptic life mode near surface. Test construction enables life under strong hydrodynamic regimes. Lives firmly attached to hard substrates, thus counteracting transportation by water movement. Nummulites venosus lives exclusively on coarse sand and avoids high sediment movement, starting distribution beneath fair weather wave base. Cycloclypeus carpenteri easily transported due to thin, plate-like form. Upper distribution limit correlates with storm wave base, below 50m. Lower distribution limit depends on light intensity and is near base photic zone)

Hohenegger, J., E.K. Yordanova, Y. Nakano & F.Tatzreiter (1999)- Habitats of larger foraminifera on the upper reef slope of Sesoko Island, Okinawa, Japan. Mar. Micropal. 36, p. 109-168.
(Peneroplis common on reef flat, hardgrounds down to 30m. Dendritina on sandy bottoms avoids uppermost slope, found down to 50m. Alveolinella similar distribution, common on hard bottom. Parasorites restricted to sandy substrates, 20-80m. Sorites and Amphisorus firm substrates between reef edge and 50m. Amphistegina species prefer hardgrounds, A. radiata also common on sand. Calcarinids withstand high energy, abundant on firm substrates close to reef edge. Baculogypsinoides deeper slope, sandy bottom, avoids shallowest parts. Hard substrates settled by Heterostegina down to 80m, occasionally on sandy bottoms. Nummulites on sands between 20- 70m. Operculina, starting at 20m, sandy substrates, rare individuals on rubble)

Hoibian, T. (1984)- La microfaune benthique traceur de l'evolution d'un systeme deltaique sous climat equatorial: le delta de la Mahakam (Kalimantan). Doct. Thesis Universite Bordeaux, p. 1-169. *(Unpublished)*
('Benthic microfauna tracing evolution of a delta system under equatorial climate: the Mahakam Delta (Kalimantan)')

Ho Kiam Fui (1971)- Distribution of recent benthonic foraminifera in the "inner" Brunei Bay. Brunei Museum J. 2, 3, p. 124-137.
(Three foraminifera assemblage in nearly landlocked Brunei Bay: (1) Trochammina cf. lobata and other small arenacous species (tidal inlets); (2) Ammobaculites (large part of inner bay); (3) Asterorotalia trispinosa with Ammonia, Elphidium, Florilus (seaward part of inner bay))

Holbourn, A., A.S. Henderson & N. Macleod (2013)- Atlas of benthic foraminifera. Wiley-Blackwell, p. 1-654.
(Atlas describing and illustrating 300 common Jurassic- Recent deep-sea benthic foraminifera species)

Hope, G.S. (1973)- The vegetation history of Mt Wilhelm. Ph.D. Thesis Australian National University, Canberra, p. 1-461.
(online at: https://digitalcollections.anu.edu.au/handle/1885/11103)
(Palynology study of sites on Mt Wilhelm PNG. Before 10,000 BP cooler and drier conditions prevailed in mountains and most mountains over 3800m in New Guinea were glaciated. Glaciers started retreating after 14,000 BP. By 8300 BP subalpine forests colonised Mt Whilhelm up to 4000m. Forests extensively cleared after 800 BP when planting of gardens took place below 2500m)

Hope, G.S. (1976)- The vegetational history of Mt Wilhelm, Papua New Guinea. J. of Ecology 64, p. 627-663.
(Pollen diagrams and 14 C dating from sites between 4420- 2740m on Mt Wilhelm, PNG Highlands allow determination of position of vegetation zones. From >22 000 years ago until 10 200 yrs B.P. tree-line stood well below 2700m and glaciers were present on mountain)

Horton, B.P., S.J. Culver, M.I.J. Hardbattle, P. Larcombe, G.A. Milne et al. (2007)- Reconstructing Holocene sea-level change for the central Great Barrier reef (Australia) using subtidal foraminifera. J. Foraminiferal Research 37, 4, p. 47-63.
(Samples behind barrier reef in water depths from 4.2- 48m. Two foraminiferal zones: inner shelf with Elphidium hispidulum, Pararotalia venusta, Planispirinella exigua, Quinqueloculina venusta and Triloculina oblonga; and middle shelf dominated by Amphistegina lessonii, Dendritina striata and Operculina complanata)

Horton, B.P., P. Larcombe, S.A. Woodroffe, J.E. Whittaker, M.R. Wright & C. Wynn (2003)- Contemporary foraminiferal distributions of a mangrove environment, Great Barrier Reef coastline, Australia: implications for sea-level reconstructions. Marine Geology 198, p. 225-243.
(Modern foraminifera and associated environmental information from Cocoa Creek, a mesotidal fringing mangrove environment on Great Barrier Reef. Three elevational zones. Zones I (highest) and II dominated by agglutinated species Trochammina inflata and Miliammina fusca, respectively; and Zone III (lowest) dominated by calcareous species, notably Ammonia tepida and Elphidium discoidale multiloculum. These assemblage zones similar to those found in both tropical and temperate intertidal environments)

Horton, B.P., J.E. Whittaker, K.H. Thomson, M.I.J. Hardbattle, S.A. Woodroffe & M.R. Wright (2005)- The development of a modern foraminiferal data set for sea-level recontructions, Wakatobi Marine National Park, Southeast Sulawesi, Indonesia. J. Foraminiferal Research 35, 1, p. 1-14.
(Intertidal foraminifera from Tukang-Besi islands off SE Sulawesi. Agglutinated species like Arenoparella mexicana, Miliammina fusca and Trochammina inflata most common at landward margin, and small calcareous species Ammonia tepida, Elphidium advanum and Quinqueloculina dominant at seaward margin of mangrove belt)

Horton, B.P., Y. Zong, C. Hillier & S. Engelhart (2007)- Diatoms from Indonesian mangroves and their suitability as sea-level indicators for tropical environments. Marine Micropaleontology 63, p. 155-168.
(Modern diatoms from mangrove swamps of Kaledupa (Tukang Besi island). 95 species, dominated by mesohalobous species (Amphora coffeaeformis, Amphora turgida, Achnanthes delicatula, Nitzschia sigma, Tryblionella balatonis) and oligohalobous (Amphora veneta, Diploneis ovalis, Progonoia didiomatia) taxa)

Howe, H.V. & K.G. McKenzie (1989)- Recent marine ostracoda (Crustacea) from Darwin and North Western Australia. Northern Territory Museum Arts Sciences, Monograph Series 3, p. 1-50.

Hughes, G.W. (1977)- Recent foraminifera from the Honiara Bay area, Solomon Islands. J. Foraminiferal Research 7, 1, p. 45-57.

Hughes, G.W. (1984)- Recent foraminifera and selected biometrics of *Heterostegina* from Ontong Java Atoll, Solomon Islands, Southwest Pacific. J. Foraminiferal Research 15, p. 13-17.
(online at: http://jfr.geoscienceworld.org/content/15/1/13.full.pdf)
(56 species of foraminifera in Halimeda gravels from Ontong Java Atoll lagoon between 31-38 m water depth. Amphistegina lessoni and Heterostegina depressa two most common species in all samples)

Hughes, G.W. (1988)- Modern bathyal agglutinating foraminifera from the Vella Gulf and Blanche Channel, New Georgia, Solomon Islands, Southwest Pacific. J. Foraminiferal Research 18, p. 304-310.
(online at: http://jfr.geoscienceworld.org/content/18/4/304.full.pdf)
(Diverse agglutinated foraminiferal assemblages of 39 species between 510- 1225m water depth off Solomon Islands. Below ~600m Bathysiphon, Ammodiscus, Martinottiella, Karreriella appear. Increase in abundance and diversity between 865-1070m associated with regional oxygen-maximum zone between 450m and 750m and a salinity-minimum zone below 800m)

Hughes, G.W. (1995)- Recent foraminifera from inter-reef channels, nearshore North Rarotonga, Cook Islands, South Pacific. J. Micropalaeontology 14, 1, p. 29-36.
(online at: https://www.j-micropalaeontol.net/14/29/1995/jm-14-29-1995.pdf)
(Samples from 8-65 water depth in Avatiu and Avarua channels of Rarotonga Island, S Pacific. Foraminifera mainly Cymbaloporetta bradyi, Borelis schlumbergeri, Heterostegina depressa, Peneroplis pertusus, Planorbulinella larvata, Siphogenerina raphanus, Sorites marginalis, Reussella simplex, Spirillina vivipara, Rosalina globularis, Amphistegina radiata, Planispirinella exigua and small miliolids)

Hughes, G.W. (2008)- Recent brackish Foraminifera and Thecamoebae from Sedili River, West Malaysia. Malaysia. In: M.A. Kaminski & R. Coccioni (eds.) Proc. 7th Int. Workshop on agglutinated foraminifera, Grzybowski Foundation Special Publ. 13, p. 41-45.

(Sedili River of S Malay Peninsula enters S China Sea through wide estuary. Lower estuarine, mangrove-fringed regime with normal marine salinity (34 ppt) has mixed calcareous-agglutinated autochthonous foraminiferal assemblage with Trochammina, Tiphotrocha comprimata, Haplophragmoides and Ammobaculites and surge-transported inner neritic calcareous benthic forams (Asterorotalia, Cellanthus, Triloculina, Ammonia, Elphidium, etc.) and rare planktonics. Estuarine, mangrove-fringed regime upstream of lower estuarine contains diverse agglutinated foraminiferal assemblages dominated by Trochammina spp., Tiphotrocha, Cribrostomoides and Ammobaculites. Upper estuarine, freshwater-slightly brackish Pandanu-grass-fringed tidal regime with low diversity agglutinated foram assemblages dominated by Miliammina fusca and Spirolocammina sp. Upstream freshwater areas of Nipa palm and grass-fringed river banks barren of foraminifera. Thecamoebae include Difflugia oblonga, Cucurbitella tricuspis and Nebela colaris)

Hunt, C.O., D.D. Gilbertson & G. Rushworth (2012)- A 50,000-year record of Late Pleistocene tropical vegetation and human impact in lowland Borneo. Quaternary Science Reviews 37, p. 61-80.
(Palynology from exposures in Great Cave of Niah, Sarawak, spanning ~52,000-5000 BP records climate-driven vegetation changes. Interstadials (interglacial periods) are marked by lowland forest, stadials by taxa from open environments or, as at the Last Glacial Maximum by disturbed open forest. Glacial stadials also characterised by taxa now restricted to 1000-1600 m above sea level, suggesting temperature declines of ~ 7–9 °C relative to todayt. Biomass burning appears associated with earliest human activity in cave)

Hunt, C.O. & R. Premathilake (2012)- Early Holocene vegetation, human activity and climate from Sarawak, Malaysian Borneo. Quaternary International 249, p. 105-119.
(40 m core from Loagan Bunut yielded high-resolution E Holocene (11.3- 6.75 ka) sequence of marginal-marine deposits)

Hussain, S.M. (2017)- An overview of ostracoda studies from the freshwater, marginal marine and marine ecosystems of Andaman and Nicobar Islands and the coasts of India. In: P K Kathal et al. (eds.) Micropaleontology and its applications, Scientific Publishers, India, Chapter 10, p. 135-162.

Hussain, S.M., P. Ganesan, G. Ravi, S.P. Mohan & S.G.D. Sridhar (2007)- Distribution of ostracoda in marine and marginal marine habitats off Tamil Nadu and adjoining areas, SE coast of India and Andaman Islands: environmental implications. Indian J. Marine Sciences 36, 4, p. 369-377.
(online at: http://nopr.niscair.res.in/bitstream/123456789/68/1/IJMS%2036(4)%20(2007)%20369-377.pdf)

Hussain, S.M., R. Krishnamurthy, M.S. Gandhi, K. Ilayaraja, P. Ganesan & S.P. Mohan (2006)- Micropaleontological investigations of tsunamigenic sediments of Andaman Islands. Current Science 91, p. 1655-1667.
(online at: https://www.currentscience.ac.in/Volumes/91/12/1655.pdf)
(Diverse marine foraminifera and ostracods from likely tsunami deposits on Andaman Islands. Common Amphistegina., Operculina ammonoides, Calcarina, Textularia, Ammonia, etc. Also deeper marine elements, sediments derived from Shallow littoral to neritic depths)

Hussain, S.M., S.P. Mohan & M.P. Jonathan (2010)- Ostracoda as an aid in identifying 2004 tsunami sediments: a report from SE coast of India. Natural Hazards 55, p. 513-522.
(Presence of marine ostracods in 2004 coastal tsunami deposits)

Hustedt, F. (1938)- Systematische und okologische Untersuchungen uber die Diatomeen-Flora von Java, Bali und Sumatra nach dem Material der Deutschen Limnologischen Sunda-Expedition. 1: Systematischer Archiv. Hydrobiologie, Suppl. Band 15, p. 1-790. *(Reprinted 1980 by Otto Koeltz Science Publishers, Konigstein)*
('Systematic and ecological investigations of the diatom floras of Java, Bali and Sumatra…'. Recent freshwater diatoms)

Hustedt, F. (1942)- Susswasser-Diatomeen des indomalayischen Archipels und der Hawaii-Inseln. Nach dem Material der Wallacea-Expedition. Internationale Revue der gesamten Hydrobiologie und Hydrographie 42, 1-3, p. 1-252. *(Reprinted 1979, Otto Koeltz Science Publishers, Germany)*

('Freshwater diatoms of the Indo-Malayan Archipelago and the Hawaii islands-From the material collected by the Wallacea Expedition'. Diatoms from samples collected during the German Limnological Sunda Expedition, 1937-1939))'

Isnaniawardhani, V. (2006)- Biostratigrafi dan paleoekologi berdasarkan nannoplankton dan foraminifera daerah Perairan Madura sejak Pliosen hingga Resen. Ph.D. Thesis Institut Teknologi Bandung (ITB), p. 1- . *(Unpublished)*
('Biostratigraphy and paleoecology based on nannoplankton and foraminifera in the Madura waters from Pliocene to Recent'. 10 biozones in waters S of Madura. Climate trends: warm conditions characterized by nannoplankton Discoaster quinqueramus and foram Globorotalia tumida)

Isnaniawardhani, V. (2009)- Environmental control of nannoplankton and foraminifera assemblages in Madura waters. Bulletin of the Marine Geology 24, 1, p. 1-12.
(online at: http://isjd.pdii.lipi.go.id/admin/jurnal/24109112.pdf)
(Distribution of nannoplankton and foraminifera in 26 shallow marine surface sediment samples from Madura Strait and 24 samples from open marine water N of Madura)

Isnaniawardhani, V. (2012)- Karakteristik sedimen dan mikroorganisma permukaan dasar laut perairan Madura bagian utara. Bull. Scientific Contribution (UNPAD) 10, 1, p. 18-30.
(online at: http://jurnal.unpad.ac.id/bsc/article/view/8275/3822)
('Characteristics and microorganisms of seafloor sediments north of Madura'. Clay and silt seafloor samples in water depths 5-77m with 20 nannoplankton species (mainly Emiliania huxleyi, Gephyrocapsa oceanica), 30 planktonic forams species (mainly Globigerinoides ruber) and 16 benthic foraminifera species (common Ammonia, Quinqueloculina, Eponides, Triloculina, Asterorotalia, Cibicides, Cancris, Elphidium, Textularia). Pseudorotalia, Cibicides and Anomalinella more abundant in N, away from coast. Abundance and diversity increase with depth. Gephyrocapsa mainly in samples closest to shoreline)

Isnaniawardhani, V. (2016)- Environmental control of nannoplankton and foraminifera assemblages in Madura waters. Bulletin of the Marine Geology 24, 1, p. 1-13.
(online at: http://ejournal.mgi.esdm.go.id/index.php/bomg/article/viewFile/10/10)
(Distributions of calcareous nannofossils (mainly Gephyrocapsa oceanica) and foraminifera in Recent sediments around Madura Island, divided into four zones: (I) inner shelf (< 30 m) in Madura Strait, (II) inner shelf in open marine N of Madura, (III) outer shelf (water depth 30- 80 m) in Madura Strait, and (IV) outer shelf in open marine N of Madura)

Isnaniawardhani, V. & R. Kapid (2016)- Kumpulan nannoplankton dan foraminifera di Selat Madura, sebarannya dalam batimetri serta letak terhadap sungai dan pantai. Bul. Ilmiah Mineral dan Energi (MINDAGI) 10, 2, p. 47-60.
('Collection of nannoplankton and foraminifera in Madura Strait, bathymetric distribution and location of rivers and beaches'. Recent foraminifera and nannoplankton in 38 seafloor samples from 4-72m water depth)

Isnaniawardhani, V. & F. Muhammadsyah (2015)- Kelimpahan, keanekaragaman dan spesies khas dari kumpulan foraminifera bentik pada sedimen permukaan dasar laut di perairan Tambelan. Bull. Scientific Contribution (UNPAD) 13, 3, p. 259-269.
(online at: http://jurnal.unpad.ac.id/bsc/article/view/8413/3920)
('Abundance, diversity and species of benthic foraminifera in sea floor sediments in waters of Tambelan'. Benthic foraminifera from shallow waters around Tambelan Islands, Natuna Sea)

Isnaniawardhani, V., E. Suparka, R. Kapid & H. Latief (2002)- Calcareous nannoplankton and foraminifera in the surface sediments of Madura waters. Proc. 31st Annual Conv. Indonesian Association Geologists (IAGI), Surabaya, 1, p. 380-386.
(online at: https://www.iagi.or.id/web/digital/17/CALCAREOUS-NANNOPLANKTON-AND-FORAMINIFERA-IN-THE-SURFACE-SEDIMENTS-OF-MADURA-WATERS.pdf)

(Recent calcareous nannoplankton assemblages N of Madura island dominated by Gephyrocapsa oceanica, Emiliana huxleyi and Florisphaera. Not much detail (see also Isnaniawardhani 2009, 2012, 2016))

Isnaniawardhani, V., E. Suparka, R. Kapid & H. Latief (2003)- Nannoplankton and foraminifera assemblages and their relations to bathymetry in Madura waters. Proc. 8th International Congress Pacific Neogene Stratigraphy, Chiang Mai 2003, p.

Jell, J.S., W.H.G. Maxwell & R.G. McKellar (1965)- The significance of the larger foraminifera in the Heron Island reef sediments. Journal of Paleontology 39, 2, p. 273-279.
(Distribution of foraminiferal detritus in sediments of Heron Island Reef, Great Barrier Reef Province. Dominant genus Calcarina spp., Baculogypsina sphaerulata and Marginopora vertebralis mainly in outer parts of reef flats. Amphistegina, Elphidium, Operculina, Peneroplis and Alveolinella quoii relatively rare)

Jian, Z. & L. Wang (1997)- Late Quaternary benthic foraminifera and deepwater paleoceanography in the South China Sea. Marine Micropaleontology 32, p. 127-154.
(In deepwater S China Sea different foram assemblages associated with Intermediate Water Mass (Globocassidulina subglobosa), Deep Water Mass (Astrononion novozealandicum and Bulimina aculeata) and Deep water below CCD (Eggerella bradyi))

Jian, Z., L. Wang, M. Kienast, M. Sarnthein, W. Kuhnt, H. Lin & P. Wang (1999)- Benthic foraminiferal paleoceanography of the South China Sea over the last 40,000 years. Marine Geology 156, 1, p. 159-186.
(During periods of high organic carbon flux during last glacial maximum (~10 ka; possibly due to increased surface productivity, induced by increased input of nutrients from nearby river runoff) detritus feeders like Bulimina aculeata and Uvigerina peregrina dominated benthic foraminiferal assemblages. Suspension feeders like Cibicidoides wuellerstorfi and 'opportunistic' species like Oridorsalis umbonatus, Melonis barleeanum and Chilostomella ovoidea gradually became more abundant as soon as organic carbon flux decreased)

Jones, R.W. (1994)- The Challenger Foraminifera. Oxford University Press, p. 1-149.
(Updated taxonomy and reproductions of foraminifera from classic H.B. Brady (1884) Challenger report)

Jones, R.W. (2014)- Supplemental notes on Challenger Foraminifera. In: A.J. Bowden et al. (eds.) Landmarks in foraminiferal micropalaeontology: history and development, The Micropalaeontological Society, Special Publ. 6, Geological Society, London, p. 31-45.
(Updates to Jones (1994) updates to names of foraminifera in Brady Challenger Report)

Jouse, A., P. & G.H. Kazarina (1974)- Pleistocene diatoms from site 262 leg 27, DSDP. Initial Reports Deep Sea Drilling Project (DSDP) 27, U.S. Government Printing Office, Washington, p. 925-946.
(online at: www.deepseadrilling.org/27/volume/dsdp27_42.pdf)
(Site 262 near axis of the Timor Trough, 75 km S of W tip of Timor in 2315m water depth. Mainly nannofossil oozes with some terrigenous material. Diatoms only in upper 250m (M-U Pleistocene; 5 zones), not numerous, well preserved, 97 species, dominantly oceanic (38). All samples with of Thalassionema and Thalassiotrix)

Joyce, E.M., K.R. Thiele, J.W.F. Slik & D.M. Crayn (2020)- Plants will cross the lines: climate and available land mass are the major determinants of phytogeographical patterns in the Sunda-Sahul Convergence Zone. Biological J. Linnean Society, London, 132, 2, p. 374-387.
(online at: https://academic.oup.com/biolinnean/article/132/2/374/6040689)
(Miocene convergence of Sunda shelf, Sahul shelf and Wallacea facilitated exchange of previously isolated floras across the Sunda-Sahul Convergence Zone (SSCZ). Lydekker's and Wallace's Lines not reflected in flora compositions: 46% of genera distributed across these biogeographical lines. Modern phytogeographical patterns influenced more by environmental variables (rainfall, seasonality) and land areas available for colonization than by proposed barriers to movement such as Wallace's and Lydekker's Lines)

Jumnongthai, J. (1983)- Recent smaller foraminifera from the Gulf of Thailand. J. Geological Society Thailand 6, p. 39-53.

(online at: http://library.dmr.go.th/Document/J-Index/1983/88.pdf)
(Foram distribution in 18 samples from water depth 29-74m in N Gulf of Thailand >99% benthics (83% calcareous). Common Asterorotalia pulchella, Cellanthus craticulatus, Elphidium, Pseudorotalia spp., Quinqueloculina, Textularia, etc.))

Jumnongthai, J. (2001)- Brackish foraminifera from southern provinces along the Gulf of Thailand. Dept. Mineral Resources (DMR), Bangkok, Technical Report No. GSD 254/2001, p. 1-40.
(online at: http://library.dmr.go.th/Document/DMR_Technical_Reports/2001/1746.pdf)
(Brackish water foraminifera from five provinces along Gulf of Thailand 53 benthic species. Low salinity facies with Ammobaculites, Ammotium cassis, Arenoparella, Miliammina fusca and Trochammina. Higher salinity assemblages with more calcareous forms Elphidium, Pararotalia nipponica))

Jumnongthai, J. (2002)- Recent brackish foraminifera from southern peninsular Thailand. J. Geological Society Thailand 1, p. 35-46.
(online at: http://library.dmr.go.th/Document/J-Index/2002/136.pdf)
(92 species of brackish foraminifera in estuaries and coastal zones along Andaman Sea. Arenacous taxa Arenoparella mexicana, Haplophragmoids, Miliammini fusca and Trochammina dominant in mangrove forests; calcareous taxa Ammonia beccarii, Elphidium and Pararotalia nipponica dominant in coastal areas)

Jurnaliah, L. & Winantris (2015)- Distribusi submikrofosil (polen dan foraminifera) pada delta front di delta Mahakam, Kalimantan Timur. Bull. Scientific Contribution (UNPAD) 13, 3, p. 169-181.
(online at: http://jurnal.unpad.ac.id/bsc/article/view/8404/3911)
('Distribution of microfossils (pollen and foraminifera) in the Mahakam delta, East Kalimantan'. 29 samples from delta front. Pollen 24 species of palmae, 21 mangrove and 117 species of non-mangrove and 7 biofacies. Foraminifera 82 species of small benthics, also 7 biofacies, from inner shelf to mangrove swamp)

Jurnaliah, L. & Winantris (2021)- Biozona Delta Mahakam modern berdasarkan kumpulan foraminifera bentik kecil dan polen, Kalimantan Timur. J. Geologi Kelautan 19, 2, p. 96-102.
('Biozones in the modernMahakam Delta, based on small benthic foraminifera and pollen, E. Kalimantan')
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/742/539)
(83 sediment samples from 3 parts of Mahakam Delta show 4 biozones: I (Palmae pollen Oncosperma, Nypa) located in delta plain; II (Foraminifera "A"; Ammonia spp, Asterorotalia); III (Foraminifera "B";Elphidium, Hateroleoa, Ammonia) in delta front; IV (non-mangrove pollen like Graminae, Cyperaceae) mainly in delta plain and distributary channel. N part of Mahakam Delta more diverse than S part)

Jurnaliah, L., I. Syafri, A. Sudradjat & R. Kapid (2019)- Biofasies dan ekologi perairan Jawa Tengah bagian utara berdasarkan kumpulan foraminifera bentik kecil. J. Geologi Kelautan 17, 2, p. 89-98.
(online at: https://garuda.kemdikbud.go.id/documents/detail/1643798)
('Biofacies and ecology of the waters of Java Sea N of Central Java, based on benthic foram assemblages'.32 bottom samples from shallow part of Java Sea suggest 4 Recent benthic foraminifera biofacies)

Kalebos, R.C., J.M. Mamuaja, M.T. Lasut, M. Ompi, R.M. Rampengan & K. Kemer (2024)- Komposisi foraminifera genus *Calcarina* di perairan desa Wineru, Kecamatan Likupang Timur, Sulawesi Utara. Jurnal Pesisir dan Laut Tropis 12, 2, p. 72-81.
(online at: https://ejournal.unsrat.ac.id/v3/index.php/jplt/article/view/57804/47496)
('The composition of foraminifera of the genus Calcarina in the waters of Wineru village, East Likupang District, North Sulawesi'. Samples of Recent Calcarina from 1-2m deep reef habitats in N-most N Sulawesi. Dominant species in all locations is C. spengleri, followed by C. sp., C hispida, C. gaudichaudii and C. defrancei)

Kamaludin b. Hassan (1989)- Significance of palynology in Late Quaternary sediments in Peninsular Malaysia. Bull. Geological Society Malaysia 24, p. 57-66.
(online at: https://gsmpubl.files.wordpress.com/2014/09/bgsm1989b05.pdf)

Kapustin, D.A., A.M. Glushchenko, J.P. Kociolek & M.S. Kulikovskiy (2021)- *Encyonopsis indonesica* sp. nov. (Bacillariophyceae, Cymbellales), a new diatom from the ancient lake Matano (Sulawesi, Indonesia). PhytoKeys 175, 61044, p. 1-11.
(online at: https://www.researchgate.net/publication/350123595_Encyonopsis_indonesica_sp_nov_Bacillariophyceae_Cymbellales_a_new_diatom_from_the_ancient_lake_Matano_Sulawesi_Indonesia)
(New diatom species, Encyonopsis indonesica described from ancient lake Matano, C Sulawesi)

Kapustin, D.A., M. Kulikovskiy & J.P. Kociolek (2018)- *Celebesia* gen. nov., a new cymbelloid diatom genus from the ancient Lake Matano (Sulawesi Island, Indonesia). Nova Hedwigia, Beiheft 146, p. 147-155.
(Cymbella distinguenda Hustedt 1942 is rare endemic freshwater diatom species from ancient Lake Matano, Sulawesi. Here placed in A new genus, Celebesia

Karmini A.S., Mimin (1996)- Foraminifera sedimen permukaan perairan Teluk Semangko & Lepas pantai sebelah Barat Sumatra Selatan. Proc. 25th Annual Conv. Indonesian Association Geologists (IAGI), Bandung, p. 71-87.
('Foraminifera in seafloor sediments of Semangko Bay and Lepas beach, S Sumatra')

Kathal, P.K. (2002)- Taxonomy, distribution patterns and ecology of Recent littoral foraminifera of the East coast of India. Neues Jahrbuch Geologie Palaontologie, Abhandlungen 224, 1, p. 115-160.
(Foraminifera distributions in 96 samples of coastal sediments down to 3m depth along E coast India. Faunas dominated by Miliolidae, followed by Rotaliidae, Elphididae, Nonionidae, Cassidulinidae, etc.)

Kawagata, S., B.W. Hayward & A.K. Gupta (2006)- Benthic foraminiferal extinctions linked to late Pliocene-Pleistocene deep-sea circulation changes in the northern Indian Ocean (ODP sites 722 and 758). Marine Micropaleontology 58, p. 219-242.
(Late Pliocene- M Pleistocene decline and extinction of 63 species of elongate, bathyal-upper abyssal benthic foraminifera (Stilostomellidae, Pleurostomellidae, some Nodosariidae. Two Indian Ocean ODP sites show pulsed declines in Extinction Group abundance and richness, especially in glacial periods, with partial recoveries in interglacials. Glacial declines result of increased production of colder, well-ventilated Antarctic Bottom Water and Glacial North Atlantic Intermediate Water)

Kawamura, H. (2002)- Marine palynological records in the southern South China Sea over the last 44 kyr. Doctoral Dissertation Christian-Albrechts-University, Kiel, p. 1-145.
(online at: at http://deposit.ddb.de/cgi-bin/...)
(Mainly on Recent and Quaternary dinoflagellates in area of Molengraaff paleo-river at N margin of Sunda Shelf)

Kawamura, H. (2004)- Dinoflagellate distribution along a shelf to slope transect of an oligotrophic tropical sea (Sunda Shelf, South China sea). Phycological Research 52, 4, p. 355-375.
(51 surface samples along Sunda shelf to S China Sea slope with36 taxa of dinoflagellate cysts. Oligotrophic tropical shelf assemblages dominated by gonyaulacoids (Spiniferites, Operculodinium spp.). Slope assemblages dominated by protoperidinioids, possibly reflecting higher nutrient availability)

Keij, A.J. (1953)- Preliminary note on the Recent Ostracoda of the Snellius Expedition. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, B 56, 2, p. 155-168.
(13 species of podocopid and one platycopid ostracode species from 28 stations of 1929-1930 Snellius Expedition in E Indonesian Seas. Ostracoda still found in samples from 4000-5000m)

Keij, A.J. (1964)- The relative abundance of recent planktonic foraminifera in seabed samples collected offshore Brunei and Sabah. Annual Report Geological Survey Borneo Region Malaysia 1963, p. 146-153.
(Percentages of planktonic foraminifera increase with depth in 561 seafloor samples from narrow Brunei and Sabah shelf between 4-113m: rare between 0-20m, <5% between 20-40m, 5-40% between 40-100m, up to 80% between 100-200m. Distinct increases in relative abundance of Orbulina, Pulleniatina, Globorotalia menardii

below 40-50m. Globorotalia truncatulinoides, Gr. crassaformis and Sphaeroidinella dehiscens only below 100m)

Keij, A.J. (1975)- Some Recent Ostracoda of Manila (Philippines). Proc. Koninklijke Nederlandse Akademie van Wetenschappen B 78, p. 351-363.

Keijzer, C.J. (1935)- On variability in East Indian foraminifera. Doct. Thesis University of Leiden, Brill, p. 1-79. *(Unpublished)*
(Biometric study of selected modern foraminifera species from coasts of Java, Bali, Madura)

Khare, N., S.K. Chaturvedi & A. Mazumder (2007)- An overview of foraminiferal studies in nearshore regions off eastern coast of India, and Andaman and Nicobar Islands. Indian J. Marine Sciences 36, 4, p. 288-300.
(online at: http://nopr.niscair.res.in/bitstream/123456789/52/1/IJMS%2036(4)%20(2007)%20288-300.pdf)
(Review and bibliography of over 100 papers dealing with shallow marine and coastal Recent foraminifera)

Khustina, Y.C., D.D. Solihin & N.T.M. Pratiwi (2014)- Morphological identification and diversity analysis of fossil diatoms from diatomite Sangiran Central Java Indonesia. Jurnal Biologi Indonesia 10, 2, p. 247-258.
(online at: https://e-journal.biologi.lipi.go.id/index.php/jurnal_biologi_indonesia/article/view/2105)
(Pleistocene lower, middle and upper diatomites in Sangiran with 50 species identified. Highest diversity in lower diatomite (34 species, mixed marine and freshwater; dominated by Coscinodiscus kutzingi). Middle diatomite dominated by Fragilaria and Coscinodiscus lineatus. Upper diatomite mainly Fragilaria, Cyclotella and Coscinodiscus spp. Gradual changes reflect change from marine to freshwater environment)

Kleijne, A. (1990)- Distribution and malformation of extant calcareous nannoplankton in the Indonesian Seas. Marine Micropaleontology 16, p. 293-316.
(Calcareous nannoplankton distribution in 202 samples from Snellius-II Expedition in Banda Sea and adjacent seas. 36 living species recorded; most common Gephyrocapsa oceanica, Umbellosphaera irregularis, Emiliania huxleyi and U. sibogae. Coccolithophorids present, but devoid of coccoliths during NW monsoon, suggesting low salinity and nutrient depletion of surface waters restrict coccolith formation, since normal coccoliths do develop during SE monsoon when upwelling causes nutrient enrichment and normal salinity)

Kob, M.R.C. (1993)- Late Quaternary nannofossils from offshore Sabah, northwest Borneo. In: T. Thanasuthipitak (ed.) Proc. International Symposium Biostratigraphy of mainland Southeast Asia: facies & paleontology (BIOSEA), Chiang Mai 1993, Chiang Mai University, 2, p. 261-281.
(online at: http://library.dmr.go.th/Document/Proceedings-Yearbooks/M_1/1993/2298_2 ...)
(late Pleistocene- Recent calcareous nannofossils from 13m thick core KL139 in ~2700m water depth of Sabah Trough. Emiliana huxleyi, Gephyrocapsa oceanica and 'small Gephyrocapsa' >50% of total assemblages. Four zones: deeper zone C (Late Pleistocene Last Glacial Maximum?) with peak 'small Gephyrocapsa' represents rel. cool period, youngest zone A with peak G. oceanica rel. warm)

Koba, M. (1978)- Distribution and environment of Recent *Cycloclypeus*. Science Reports Tohuku University, ser. 7, 28, p. 283-311.
(online at: http://ir.library.tohoku.ac.jp/re/bitstream/10097/45065/1/AA0045945078466.pdf)
(Tropical larger foram Cycloclypeus widely distributed in Recent of Indo-Pacific oceanic region. Common on outer reef slopes, and confined by 200m isobath. Not present in pelagic environments, reef banks, lagoons or bays. In Recent samples found between 32-1419m depth, but clear abundance peak around 90m)

Langer, M.R. (1992)- New Recent foraminiferal genera and species from the lagoon at Madang, Papua New Guinea. J. Micropalaeontology 11, 1, p. 85-93.
(online at: https://www.j-micropalaeontol.net/11/85/1992/jm-11-85-1992.pdf)
(Samples from shallow water (0-55m) fore- and back-reef environments at Madang lagoon with larger foraminifera (Assilina spp., Heterostegina depressa, Alveolinella quoyi, Sorites spp., Amphisorus hemprichii, Marginopora vertebralis). Two new genera (Pseudolachlanella) and eight new species of benthic foraminifera)

Langer, M.R. (1995)- Oxygen and carbon isotopic composition of Recent larger and smaller foraminifera from the Madang Lagoon (Papua New Guinea). Marine Micropaleontology 26, p. 215-221.
(General trend in Recent large benthic foraminifera (Operculina, Heterostegina, Sorites, Marginopora, Alveolinella) from Madang Lagoon of depletion in heavier C and O isotopes with depth appears to be depth- and light-dependant)

Langer, M.R. & L. Hottinger (2000)- Biogeography of selected 'larger' foraminifera. Micropaleontology 46, Suppl. 1, Advances in the biology of foraminifera, p. 105-126.
(Global distributions of Recent larger foraminifera species. Main provinces: (1) Central Pacific with Baculogypsinoides spinosus, Schlumbergerella floresiana, Operculina heterosteginoides, Pseudorotalia indopacifica; (2) Indo-Pacific with Marginopora vertebralis, Alveolinella quoyii, Amphistegina radiata, Calcarina spp., Nummulites venosus and Cycloclypeus carpenteri; (3) W Indian Ocean and (4) Caribbean)

Langer, M.R. & J.H. Lipps (2003)- Foraminiferal distribution and diversity, Madang reef and lagoon, Papua New Guinea. Coral Reefs 22, p. 143-154.
(Benthic foram distribution shows four clusters in Madang lagoon at NE coast of PNG)

Lei, Y. & T. Li (2016)- Atlas of benthic foraminifera from China Seas, the Bohai Sea and the Yellow Sea. IUP Science Press, Beijing, Springer, p. 1-399.
(Descriptions of 183 species of mainly shallow marine foraminifera from northern South China Sea)

Lelono, E.B. (2006)- Pleistocene palynology of East Java. Lemigas Scientific Contributions 29, 3, p. 3-14.
(online at: https://journal.lemigas.esdm.go.id/index.php/SCOG/article/view/1027)
(Study of Pleistocene palynology in 3 unnamed wells in NE Java basin. Base Pleistocene characterised by index pollen of Dacrycarpidites australiensis and abundant grass pollen of Monoporites annulatus in the absence of spore of Stenochlaenidites papuanus. Interpretation supported by occurrence of nannoplankton Gephyrocapsa eocenica (NN19). Benthic zone lower NB1 (equivalent to E Pleistocene planktonic foram zone N22, based on appearance of Asterorotalia gaimardii and Pseudorotalia schroeteriana in absence of Asterorotalia gaimardii inermis, Pseudorotalia schroeteriana angusta, P. alveiformis, Asanoina globosa and Ammonia ikebei)

Lelono, E.B. (2018)- Dry climate expansion in the Pleistocene of Indonesia as recorded in its pollen assemblage. Scientific Contributions Oil and Gas (SCOG), Lemigas, Jakarta, 41, 1, p. 17-27.
(online at: http://journal.lemigas.esdm.go.id/ojs/index.php/SCOG/article/download/69/pdf)
(Pleistocene characterised by glacial and interglacial periods. In Indonesia reflected in dry/cool (abundant Gramineae grass pollen) and wet/warm climates (increases of coastal and mangrove palynomorphs). Quantitative palynology shows Pleistocene in E Java wells characterised by abundant Monoporites annulatus grass pollen, corresponding to glacial-period expansion of savanna vegetation. Abundant charred Gramineae cuticles reflect burning grass. In N Papua wells repetition of dry/ wet conditions or low/ high sea levels)

LeRoy, L.W. (1938)- A preliminary study of the microfaunal facies along a traverse across Peper Bay, West coast of Java. De Ingenieur in Nederlandsch-Indie (IV) 5, 8, p. IV.130- IV.133.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1938-08.pdf)
(Recent foraminifera off West coast of Java, SW of Labuan (deepest sample 32m). Three assemblages (1) Haplophragmoides- Haplophragmium facies (semi-brackish), (2) Operculina- Ozawaia facies (18-32m, marine) and (3) Dendritina-Aveolinella facies (7-26m; also Rotalia calcar, Calcarina spengleri, Marginopora, etc.))

Lessard, R.H. (1964)- Intertidal and shallow water foraminifera of the tropical Pacific Ocean. M.Sc. Thesis, University of Southern California, p. 1-112.
(online at: http://digitallibrary.usc.edu/cdm/ref/collection/p15799coll30/id/107855)
(Mainly on distribution of Baculogypsina and Tinoporus in western tropical Pacific)

Lessard, R.H. (1980)- Distribution patterns of intertidal and shallow water foraminifera of the tropical Pacific Ocean. Cushman Foundation Foraminiferal Research, Special Publ. 19, p. 40-58.

(online at: https://business.cushmanfoundation.org/ap/CloudFile/Download/rJDJ8V5P
(Recent foraminifera from intertidal and shallow-water sediment samples along two major sections across islands of tropical Pacific Ocean, dominated by Tinoporus, Baculogypsina, Atnphistegina, Marginopora and Heterostegina. With discussions on possible dispersal mechanisms. No distrubution maps)

Li, Z., Y. Saito, L. Mao, T. Tamura, Z. Li, B. Song, Y. Zhang, A. Lu, S. Sieng & J. Li (2012)- Mid-Holocene mangrove succession and its response to sea-level change in the upper Mekong River delta, Cambodia. Quaternary Research 78, 2, p. 386-399.
(Cores from upper Mekong River delta in Cambodia record transgressive sequence from floodplain freshwater marsh to tidal flat (~9.4- 6.3 ka), overlain by mangrove. At decelerated sea-level rise at ~8.3 ka pioneer (high-salinity tolerant) mangrove species Sonneratia alba and Sonneratia caseolaris appeared, then was replaced by regressive mangrove succession of increasing Rhizophora apiculata and Bruguiera spp.)

Ling, H.Y. & W.A. Anikouchine (1967)- Some spumellarian radiolaria from the Java, Philippines and Mariana trenches. Journal of Paleontology 41, 6, p. 1481-1491.
(Eight species of five genera of patagium-bearing and morphologically closely related spumellarian Radiolaria in three sediment cores from Java, Philippine, and Mariana Trenches. Java Trench samples from pelagic ooze in core from 3380m water depth off S Sumatra, with Euchitonia spp., Hymeniastrum, Dictyocorine and Rhopalodictyum. Also locally common diatiom Ethmodiscus rex)

Lipps, J.A. & K.P. Severin (1985)- *Alveolinella quoyi*, a living fusiform foraminifera, at Motupore Island, Papua New Guinea. Science in New Guinea (University of PNG, Port Moresby) 11, p. 126-137.
(Living species of Alveolinella quoyi in water depths of 3-12m, mainly on algae-covered coral rubble and around bases of living coral heads in rel. sheltered areas. Dead tests scattered over wider bathymetric range. Virtually absent on back-reef flats)

Lobegeier, M.K. (2002)- Benthic foraminifera of the family Calcarinidae from Green Island Reef, Great Barrier Reef Province. J. Foraminiferal Research 32, 3, p. 201-216.
(Three epiphytic calcarinid species on Great Barrier Reef and limited to W Indo-Pacific (absent from Indian Ocean and E of 170°W). Calcarina spengleri (=hispida) dominant and common in shallow water on reef flat. Calcarina mayori smaller and dominates in deeper water off reef flat. Baculogypsina sphaerulata is shallow water high-energy species. Best preserved Calcarina at Green Island Reef in windward shoals)

Li, B.H., X.Y. Wang, Z.M. Jian, P.X. Wang (2009)- Sea surface environment inferred from planktonic foraminifera in the southern South China Sea since the last glacial period. Palaeoworld 18, p. 23-33.
(Planktonic foraminifera from Site 17964 in southern S China Sea (SCS) show higher % warm-water species in Holocene, while temperate-water species increase during last glacial period. Pulleniatina obliquiloculata more common during glacial period. Orbulina universa test size larger than those in Indian and Atlantic Oceans, indicating warmer and less saline surface water in Equatorial- tropical W Pacific. Diameter and shell porosity of O. universa increased from last glacial to Holocene)

Lirdwitayaprasit, T. (1997)- Distribution of dinoflagellate cysts in the surface sediment of the South China Sea, Area I: Gulf of Thailand and East Coast of Peninsular Malaysia. Proc. 1st Techn. Seminar Marine fishery resource survey in the South China Sea, SE Asia Fisheries Development Centre (SEAFDEC), Training Dept., Thailand, S4/SB7, p. 294-309.
(online at: http://map.seafdec.org/downloads/pdf/collaborative%20research/AreaI_GOT/SCS_FRS1_18.pdf)
(Dinoflagellate cysts in Gulf of Thailand and off E coast of Peninsular Malaysia 20 goniolacoid, tuberculodinioid and peridinioid species. Spiniferites spp. (=Gonyaulax spp.) dominant cyst species)

Lirdwitayaprasit, T. (1997)- Distribution of dinoflagellate cysts in the surface sediment of the South China Sea, Area II: Sabah, Sarawak and Brunei Darussalam waters. Proc. 1st Techn. Seminar Marine fishery resource survey in the South China Sea, SE Asia Fisheries Development Centre (SEAFDEC), Training Dept., Thailand, S2/FB4, p. 310-322.

(online at: http://map.seafdec.org/downloads/pdf/collaborative%20research/AreaII_West%20Borneo/SCS_FRS2_16.pdf)
(Dinoflagellate cysts off N Borneo with Spiniferites bulloideus as dominant species)

Lloyd, A.R. (1973)- Foraminifera of the Great Barrier Reef bores. In: O.A. Jones & R. Endean (eds.) Biology and geology of coral reefs, 1, Academic Press, New York, p. 347-366.

Loeblich, A.R. & H. Tappan (1994)- Foraminifera of the Sahul Shelf and Timor Sea. Cushman Foundation Foraminiferal Research, Special Publ. 31, p. 1-661.
(online at: https://business.cushmanfoundation.org/ap/CloudFile/Download/PezgBY2L)
(Comprehensive study of Recent foraminifera from Timor Trough and Sahul Shelf, collected in 1961. Mainly species descriptions (946 species of 428 genera; 101 new species)

Luan, B.T. & J.P. Debenay (2005)- Foraminifera, environmental indicators in the highly impacted environments of the Mekong Delta. Hydrobiologia 548, p. 75-83.

Lucero, E.S. & L.P. de Silva (2006)- The occurrence and distribution of Recent benthic foraminifera in Subic Bay, Zambales, Philippines. J. Geological Society of the Philippines 61, p.

Lunt, P. (2014)- Stacked digital imaging of foraminifera. Berita Sedimentologi 29, p. 123-132.
(online at: https://journal.iagi.or.id/index.php/FOSI/article/view/148)
(Description of technique of imaging foraminifera by combining series of digital photographs in automated software to produce single optical image with high resolution and superior depth of focus. With examples of Indonesian Tertiary- Recent foram material)

Makled, W.A. & M.R. Langer (2011)- Benthic foraminifera from the Chuuk Lagoon Atoll System (Caroline Islands, Pacific Ocean). Neues Jahrbuch Geologie Palaontologie, Abhandlungen 259, 2, p. 231-249.
(online at: https://www.researchgate.net/publication/233654597_Benthic_Foraminifera_from_the_Chuuk_Lagoon_Atoll_System_Caroline_Islands_Pacific_Ocean)
(Excellent documentation of Recenet shallow marine benthic foraminifera from a western Equatorial Pacific atoll system formerly called Truk, in hotspot of marine tropical diversity in modern oceans. 104 species: 66% porcelaneous, 22% hyaline-perforate and 16% agglutinated species. Amphistegina lessonii and Heterostegina depressa most abundant. Other significant species include Alveolinella quoyi and Baculogypsina sphaerulata)

Maloney, B.K. (1992)- Late Holocene climatic change in Southeast Asia: the palynological evidence and its implications for archaeology. World Archaeology 24, 1, p. 25-34.
(Review of palynological data from SE Asia Holocene from deep sea cores, archeological sites and land cores)

Maloney, B.K. (1998)- That elm again! Ulmus at Pea Bullok, North Sumatra, and regional comparisons. Blumea 43, p. 121-127.
(online at: http://repository.naturalis.nl/document/565696)
(Quaternay Ulmus pollen present in Pua Bulok swamp in N Sumatra. Ulmus not necessarily indicator for seasonal dryness)

Maloney, B.K. & F.G. McCormac (1996)- Thirty thousand years of radiocarbon dated vegetation and climatic change in highland Sumatra. Proc. 15th Int. 14C Conference, Radiocarbon 37, 2, p. 181-190.
(online at: www.cambridge.org/core/services/aop-cambridge-core/content/view/AD19D1 Etc.)
(Pollen analysis and 14C sequences from two Sumatra highland sites, Pea Bullok (Toba Plateau) and Danau di-Atas, spanning last glacial period. Strong indications of extensive forest clearance after ~2 ka)

Maloney, B.K. & F.G. McCormac (1996)- Palaeoenvironments of North Sumatra: a 30,000 year old pollen record from Pea Bullok. Bull. Indo-Pacific Prehistory Association 14, p. 73-82.
(online at: https://journals.lib.washington.edu/index.php/BIPPA/article/view/11590)
(Palynology of two 3.5 and 8m deep cores from Pea Bullock, 10km S of Lake Toba)

Mamo, B.L. (2016)- Benthic foraminifera from the Capricorn Group, Great Barrier Reef, Australia. Zootaxa 4215, 1, p. 1-123.
(Taxonomy of 133 benthic foram species from Heron Island, One Tree Island, Wistari and Sykes Reefs)

Mao, L., D.J. Batten, T. Fujiki, Z. Li, L. Dai & C. Weng (2012)- Key to mangrove pollen and spores of southern China: an aid to palynological interpretation of Quaternary deposits in the South China Sea. Review Palaeobotany Palynology 176-177, p. 41-67.
(online at: https://www.semanticscholar.org/paper/Key-to-mangrove-pollen-and-spores-of-southern-an-to-Mao-Batten/b1103575729f2b3827e06999d3cd0d2502e4b66e)
(Pollen morphological "atlas" and identification keys for mangrove flora of coastal areas of S China)

Marquez, E.J. (1999)- Deep sea foraminiferal distribution of eastern South China Sea. J. Geological Society of the Philippines 54, 1-2, p.

Marquez, E.J. (2000)- The 1991 Mount Pinatubo eruption and Eastern South China Sea foraminifera: occurrence, composition and recovery. Island Arc 9, 4, p. 527-541.
(Pyroclastic materials from 1991 eruption of Mt Pinatubo resulted in decimation of most benthic foraminifera in E South China Sea. Samples above eruption layers in deep water cores much lower abundances, lower diversity and relative common of Quinqueloculina spp., probably form part of recolonization fauna)

Marquez, E.J., P.J. Militante-Matias, G.P. Yumul, M.M. De Leon, D.V. Faustino et al. (1999)- Deep-sea foraminiferal distribution of the central and eastern portions of the South China Sea. In: G.H. Teh (ed.) Proc. 9th Regional Congress Geology Mineral Energy Resources of SE Asia (GEOSEA '98), Kuala Lumpur 1998, Bull. Geological Society Malaysia 43, p. 529-536.
(online at: www.gsm.org.my/products/702001-100787-PDF.pdf)
(1991 eruption of Mount Pinatubo covered S China Sea with W-trending ash fan. Eruption affected foraminifera only in cores taken near Luzon. Not much detail (not one species name!))

Martin, S.Q. (2016)- Distribution and taxonomy of modern benthic foraminifera of the western Sunda Shelf (South China Sea) off Peninsular Malaysia. Masters Thesis, East Carolina University, p. 1-262. *(Unpublished)*
(also as Cushman Foundation Special Publ. 47, 2018, with S.J. Culver et al.; see below)
(Distribution and taxonomy of 125 modern benthic foraminifera species from 54 seafloor samples in southern S China Sea, between 8-60m. Main assemblages across Sunda Shelf: (1) nearshore areas (<40m) dominated by symbiont-bearing Amphistegina lessonii, A. radiata; (2) inner shelf (40-100m), sandy mud substrates and abundant Heterolepa. dutemplei; (3) outer shelf (100-200m), muddy substrates and Uvigerina schwageri)

Martin, S.Q., S. J. Culver; E. Leorri; D.J. Mallinson; M.A. Buzas; L.A.C. Hayek & N.A.M. Shazili (2018)- Distribution and taxonomy of modern benthic foraminifera of the Western Sunda Shelf (South China Sea) off Peninsular Malaysia. Cushman Foundation Foraminiferal Research, Special. Publ. 47, p. 1-108.
(60 seafloor sediment samplesfrom southern South China Sea off NE Peninsular Malaysia from 8- 60m water depths yielded 125 species. Shallow samples with sandy substrates dominated by Amphistegina radiata and Amphistegina lessonii. Mud substrates at (44–59 m) characterized by Heterolepa dutemplei and Asterorotalia milletti. At 60m rel. common Heterolepa and Assilina ammonoides)

Maryunani, Khoiril Anwar (2003)- Calibration of tropical Pacific marine sediment indices (d18O) to sea surface temperature. Buletin Geologi (ITB) 35, 1, p.

Matsumaru, K. & Y. Matsuo (1976)- Short note on the Recent benthic foraminiferids from the beach sediments of the subtropical and tropical islands in the Western Pacific region. J. Saitama University Fac. Education (Math. Nat. Science) 25, p. 15-26.

Matsuoka, K., Y. Fukoyo, D.P. Praseno, Q. Adnan & M. Kodama (1999)- Dinoflagellate cysts in surface sediments of Jakarta Bay, off Ujung Pandang and Larantuka of Flores island, Indonesia with special reference of *Pyrodinium bahamense*. Bull. Fac. Fisheries, Nagasaki University, 80, p. 49-54.
(online at: http://naosite.lb.nagasaki-u.ac.jp/dspace/bitstream/10069/9236/1/80-049.PDF)

McCloskey, B.J. (2009)- Foraminiferal responses to arsenic in shallow-water hydrothermal system in Papua New Guinea and in the laboratory. Ph.D. Thesis University of South Florida, p. 1-425. *(Unpublished)*
(On shallow marine, reef flat foram assemblages around hydrothermal vents in ~8m of water at Ambitle Island, Tutum Bay, E of New Ireland, PNG)

McKenzie, K.G. & A.J. Keij (1977)- *Pterobairdia* (Ostracoda, Bairdiidae), a new genus from the Indo-West Pacific. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, B, 80, 5, p. 367-374.
(Recent ostracodes from East Indonesia Snellius Expedition, incl. Pterobairdia maddocksae n. sp.)

Melis, R. & D. Violanti (2006)- Foraminiferal biodiversity and Holocene evolution of the Phetchaburi coastal area (Thailand Gulf). Marine Micropaleontology 61, p. 94-115.
(Study of foraminifera from surface sediments from Phetchaburi coastal area, Thailand Gulf, from margin of vegetated zone, to depth of ~20m, and core in Mae Khlong delta plain. Area characterized by inflow of Mae Khlong river. Four assemblages of low diversity brackish-water faunas, with salinity controlling factor)

Mertens, K.N., Y. Takano, M.J. Head and K. Matsuoka (2014)- Living fossils in the Indo-Pacific warm pool: a refuge for thermophilic dinoflagellates during glaciations. Geology (GSA) 42, 6, p. 531-534.
(Dinoflagellate 'refugium species' Dapsilidinium pastielsii thought to range from Eocene- E Pleistocene, but still living in SE Asia (Okinawa; Koror, Palau, Ambon, E Vietnam Sea and Philippines)

Miao, Q. & R.C. Thunell (1993)- Recent deep-sea benthic foraminiferal distributions in the South China and Sulu Seas. Marine Micropaleontology 22, p. 1-32.
(Benthic foraminifera in surface sediment samples from S China and Sulu Seas. Low abondances but high diversity. Four faunal assemblages: (1) Globocassidulina subglobosa/ Uvigerina <1500m; within oxygen minimum zone); (2) Bulimina aculeata between 1700-2000m in SE S China Sea, also associated with high organic carbon content; (3) Astrononion pusillum in S China Sea between 1500-3200 m; (4) below lysocline (~3200m) in S China Sea agglutinated Rhabdammina abyssorum assemblage, in water mass that is highly undersaturated with respect to calcite. In Sulu Sea Pyrgo murrhina assemblage 1400-2200m; below 2200m assemblages dominated by Oridorsalis umbonatus)

Miao, Q. & R.C. Thunell (1996)- Late Pleistocene-Holocene distribution of deep-sea benthic foraminifera in the South China Sea and Sulu Sea- palaeoceanographic implications. J. Foraminiferal Research 26, p. 9-23.

Millett, F.W. (1898-1904)- Report on the Recent foraminifera of the Malay Archipelago collected by Mr. A. Durrand, parts I-XVII. J. Royal Microscopical Society, 1898-1904, p. 1-248.
(Reprinted by Antiquariaat Junk, Lochem, 1970. 17 papers on foraminifera from samples collected by A. Durrand at 30 stations along transect from N Australia to Malay Peninsula)

Minhat, F.I., S.M. Ghandhi, N.S.M. Ahzan, N.A. Haq, O.A.R.A. Manaf, S.M. Sabohi et al. (2021)- The occurrence and distribution of benthic foraminifera in tropical waters along the Strait of Malacca. Frontiers in Marine Science 8, 647531, p. 1-10.
(online at: https://www.frontiersin.org/articles/10.3389/fmars.2021.647531/full)
(24 seafloor samples from Malaysian waters in northern Strait of Malacca, in water depths 9-86m. With 49 species of benthic foraminifera, dominated by calcareous hyaline taxa Asterorotalia pulchella (= A. trispinosa; most dominant), Pseudorotalia schroeteriana, Discorbinella bertheloti, Ammonia tepida, and Heterolepa praecincta. Three groups)

Minhat, F.I., P.R. Parham, M.L. Husain & B. Satyanarayana (2014)- Review of foraminiferal studies in nearshore areas, Peninsular Malaysia. Geological Society Malaysia, Proc. National Geoscience Conference (NGC), Terengganu 2014, P063, p. 122-124. *(Extended Abstract)*
(online at: https://gsm.org.my/articles/702001-100318/)
(Brief review of current status of intertidal- shallow marine foraminiferal distribution work in Malaysia)

Minhat, F.I., B. Satyanarayana, M.L. Husain & V.V.V. Rajan (2016)- Modern benthic foraminifera in subtidal waters of Johor: implications for Holocene sea-level change on the East coast of Peninsular Malaysia. J. Foraminiferal Research 46, 4, p. 347-357.
(Modern subtidal benthic foraminifera on E coast of Johor 279 species, dominated by Asterorotalia pulchella (= A. tripinosa), Discorbinella bertheloti, Pseudorotalia indopacifica, Ammonia and Cavarotalia annectens. Agglutinated species Textularia pseudosolita, T. agglutinans, Bigenerina nodosaria and T. foliacea in middle-shelf (>20m), calcareous genera Elphidium, Pararotalia and Ammonia in inner-shelf (<20m))

Minhat, F.I., K. Yahya, A. Talib & O. Ahmad (2013)- A survey of benthic foraminiferal assemblages in tropical coastal waters of Pulau Pinang, Malaysia. Tropical Life Sciences Research 24, 1, p. 35-43.
(Samples from subtidal zone to 1200m offshore Pinang Island, Malacca Straits, in water depths 1.5-10m with 9 nine genera of foraminifera, dominated Ammonia, Bigenerina, Ammobaculites and Elphidium)

Minhat, F.I., K. Yahya, A. Talib & O. Ahmad (2014)- Benthic foraminiferal distributions as bioindicators in coastal waters of Penang National Park, Malaysia. J. Foraminiferal Research 44, 2, p. 143-150.
(Distribution of benthic foraminifera off NW corner of Penang Island. Water depths 1.5-10 m, with predominantly muddy substrate. All assemblages dominated by Ammonia; also Elphidium, Ammobculites, etc.)

Mohamed, Mahani, S. Jirin, S.S. Hasan & N. Mohsin (2010)- Biofacies characterisation in the marginal marine environments of the Malay Basin using agglutinated foraminifera. Proc. Petroleum Geology Conference Exhibition (PGCE), Kuala Lumpur 2010, p. *(Extended Abstract)*
(Malay Basin dominated by paralic facies, where agglutinated foraminifera are useful in characterising biofacies. Three modern localities studied for biofacies analogs: Sedili Besar Estuary, Klang-Langat Delta and Pahang River Delta. Occurrences of species such as Ammobaculites exiguus, Textularia sp and Arenoparrella mexicana used to differentiate nearshore, shallow marine and brackish intertidal depositional settings)

Mohamed, Mahani, S.S. Hasan & S. Jirin (2014)- Recent agglutinated foraminiferal trends and assemblages of the Sedili Besar River and its adjacent offshore area, southeastern Peninsular Malaysia. Berita Sedimentologi 29, p. 73-79.
(online at: https://journal.iagi.or.id/index.php/FOSI/article/view/144)
(Distribution of Recent foraminifera from Sedili river estuary to shallow marine offshore S China Sea along SE coast of Malay Peninsula. Calcareous forms increase with water depth, agglutinated foraminifera show reverse trend. Five biofacies zones differentiated, coinciinge with the upper brackish intertidal, lower brackish intertidal, estuary mouth, beach/nearshore and inner shelf depositional settings)

Mohamed, Mahani, S.S. Hasan, A.M. Yakzan & S. Jirin (2011)- Agglutinated foraminiferal trends and assemblages of the Sedili Besar River and its offshore area, southeastern Peninsular Malaysia. In: M.A. Kaminski & S. Filipescu (eds.) Proc. 8th Int. Workshop on agglutinated foraminifera, Romania, Grzybowski Foundation Special Publ. 16, p. 131-136.
(Similar to paper above)

Mohamed, Mahani, S. Jirin & S.S. Hasan & N. Mohsin (2011)- Salinity stratification and its effects on the Malay Basin biofacies assemblages. In: Petroleum Geology Conference and Exhibition 2011, Kuala Lumpur, Poster 7, p. 141-143. *(Extended Abstract)*
(Foraminifera in Sedili Besar River Estuary dominated by Ammonia cf. takanabensis (also identified as Ammonia beccarii) in stratified water column of marine base and freshwater top. In Klang-Langat and Pahang Deltas, where minimal salinity stratification, Ammonia assemblages are quite scattered. Agglutinated forms (mainly Arenoparrella group) dominate less stratified water column)

Mohan, P.M., P. Dhivya & K. Narayanamurthy (2013)- Distribution of live planktonic and benthic foraminifera in the shelf off Port Blair and Hut Bay, Andaman Group of Islands, India. In: K. Venkataraman et al. (eds.) Ecology and conservation of tropical marine faunal communities, Springer-Verlag, Berlin, p. 19-42.
(189 shelfal marine foram species off Andaman Islands (no water depths of sample locations given))

Montaggioni, L.F. & M.T. Venec-Peyre (1993)- Shallow-water foraminiferal taphocoenoses at Site 821: implications for the Pleistocene evolution of the central Great Barrier Reef shelf, Northeastern Australia. In: J.A. McKenzie et al. (eds.) Proc. Ocean Drilling Program (ODP), Scientific Results, College Station, 133, p. 365-378.
(online at: www-odp.tamu.edu/publications/133_SR/VOLUME/CHAPTERS/sr133_26.pdf)
(Useful overview of foram distribution on and around Great Barrier reef)

Morley, R.J., H.P. Morley, A.A.H. Wonders, Sukarno & S. van der Kaars (2004)- Biostratigraphy of modern (Holocene and Late Pleistocene) sediment cores from Makassar Straits. In: R.A. Noble et al. (eds.) Proc. Deepwater and frontier exploration in Asia and Australasia, Indonesian Petroleum Association, Jakarta 2004, p. 361-371.
(Palynology and foraminifera from two shallow Late Pleistocene- Holocene cores from Makassar Straits and offshore SW Sulawesi)

Mostafawi, N. (1992)- Rezente Ostracoden aus dem mittleren Sunda-Schelf, zwischen der Malaiischen Halbinsel und Borneo. Senckenbergiana Lethaea 72, p. 129-168.
('Recent ostracods from the central part of the Sunda Shelf, between the Malay Peninsula and Borneo'. 116 species of Recent ostracodes in 44 seafloor samplesalong W-E transect, all <100m depth. Four new genera.Major control on abundance / diversity appears to be substrate. Many species range across all water depths, some more diagnostic for water depth: (1) <~50m with Atjehella kingmai, Keijia spp., Hemicytheridea cf. oculosa, etc.; (2) >50m with Abracythereis malaysiana, Bythocytheridea carinatum, etc.)

Mostafawi, N., J.P. Colin & J.F. Babinot (2005)- An account on the taxonomy of ostracodes from recent reefal flat deposits in Bali, Indonesia. Revue de Micropaleontologie 48, p. 123-140.
(Ostracods from recent reefal flat sample off Sanur, SE Bali, at depth of ~1.5m. Assemblage of 34 species, dominated by Loxoconcha peterseni, Auradilus convolutus, A. australiensis, Paranesidea conulifera, etc. Fauna belongs to East Indian biogeographical province of Titterton and Whatley (1988) in tropical littoral zone of Indo-W Pacific. Associated with foraminifera Schlumbergella floresiana, Calcarina hispida, etc.)

Mousson, A. (1849)- Die Land- und Susswasser–Mollusken von Java. Nach den Sendungen des Herrn Seminardirektors Zollinger. Friedrich Schulthess, Zurich, p. 1-126.
(online at: https://www.e-rara.ch/zuz/content/zoom/11468827)
('The land- and freshwater molluscs of Java, from shipments of Mr. Seminary Director Zollinger'. One of first descriptions of freshwater molluscs and land snails of Java by Swiss naturalist of University of Zurich, Prof. Albert Mousson, collected by Swiss naturalist H. Zollinger (see also freshwater mollusc papers by E. von Martens 1897, K. Martin, W. van Benthem Jutting, Neuberth et al. 2023, etc.))

Mousson, A. (1849)- Uber die Land- und Susswassermollusken von Java. Mittheilungen Naturforschenden Gesellschaft Zurich, 1 (3, 30), p. 264-273.
('The land- and freshwater molluscs of Java'. Abbreviated version of Mousson (1849, above)

Muller, G.W. (1906)- Die Ostracoden der Siboga-Expedition. In: M. Weber (ed.) Siboga Expeditie, Uitkomsten op zoologisch, botanisch, oceanographisch en geologisch gebied, Monograph 30, Brill, Leiden, 40, p. 1-40.
(online at: http://ia800807.us.archive.org/26/items/dieostracodender00mlle/dieostracodender00mlle.pdf)
('The ostracodes of the Siboga Expedition'. 56 Recent species from shallow to deep water seafloor samples of 1899-1900 Siboga Expedition. Material unusually rich in Cypridinids (25 species, 15 new). Halocypriden mainly at depths>750m. Cypridiniden generally <70m)

Muller, J. (1965)- Palynological study of Holocene peat in Sarawak. In: Symposium on ecological research in humid tropics vegetations, Kuching, Sarawak. UNESCO, p. 147-156.

Murphy, S. (1996)- Distribution and ecology of Holocene foraminifera within the Lassa distributary of the Rajang River delta, Sarawak, East Malaysia. M.S. Thesis, Southern Illinois University, Carbondale, p. 1-121. *(Unpublished)*

Murray, J.W. (1994)- Larger foraminifera from the Chagos Archipelago: their significance for Indian Ocean biogeography. Marine Micropaleontology 24, p. 43-55.
(Chagos Archipelago is thin limestone cap on Eocene volcanic basement in central Indian Ocean. Principal larger foraminifera in surface samples from 0-43m. Amphisorus hemprichii and Sorites orbiculus widespread in shallow lagoon Heterostegina depressa patchy distribution, most common between 18-25m. Operculina ammonoides generally in deeper lagoon, below 12m)

Murray, J.W. (2006)- Ecology and applications of benthic foraminifera. Cambridge University Press, New York, p. 1-426.

Murray, J.W. & C.W. Smart (1994)- Distribution of smaller benthic foraminifera in the Chagos Archipelago, Indian Ocean. J. Micropalaeontology 13, 1, p. 47-53.
(online at: https://www.j-micropalaeontol.net/13/47/1994/jm-13-47-1994.pdf)
(Chagos Archipelago in C Indian Ocean close to the equator. Relatively high energy conditions in shallow waters around reefs. On oceanic side of atoll reefs Amphistegina lessonii dominant, with minor miliolidsand up to 20% planktonics. Lagoon assemblages dominated by Calcarina calcar, with minor miliolids)

Muruganantham, M. & P.M. Mohan (2015)- First report of three benthic foraminifera from the waters of Andaman Islands, India. Biodiversity Journal 6, 4, p. 789-794.
(online at: www.biodiversityjournal.com/pdf/6(4)_789-794.pdf)
(Living benthic foraminifera Nevillina coronata, Sigmoihauerina involuta and Loxostomina limbata reported from inner shelf regions of Andaman Islands. Nevillina coronata very common in NE, flourishing in low T of rainy season, while two other species abundant in non rainy months)

Muruganantham, M. & P.M. Mohan (2015)- The assemblages of benthic foraminifera In the muddy and sandy sediments of Andaman Islands. J. Andaman Science Association 20, 2, p. 199-208.
(online at: http://asapb.org/15%20-%20The%20Assemblages%20of%20Benthic.pdf)
(28 species of tropical marginal- shallow marine benthic foraminifera around South Andaman Island)

Muruganantham, M., P. Ragaven & P.M. Mohan (2017)- Diversity and distribution of living larger foraminifera from coral reef environments, South Andaman Islands, India. J. Foraminiferal Research 47, 3, p. 252-257.
(Larger foraminifera at six reef sites (4-30 m) around South Andaman Islands 16 species, incl. Amphistegina lessonii, A. radiata and Calcarina spengleri)

Myers, E.H. (1945)- Recent studies of sediments in the Java Sea and their significance in relation to stratigraphic and petroleum geology. In: P. Honig & F. Verdoorn (eds.) Science and scientists in the Netherlands Indies, New York, p. 265-269.
(online at: https://archive.org/details/in.ernet.dli.2015.183759)

Newman, K., B. Hakim, A.A. Oktaviana, B. Burhan, D. McGahan & A. Brumm (2022)- The missing deposits of South Sulawesi: New sources of evidence for the Pleistocene/Holocene archaeological transition. Archaeological Research in Asia 32, 100408, p. 1-16.
(S Sulawesi archeologically one of most intensively investigated parts of Indonesia. Pleistocene assemblages scarce, but S Sulawesi contains sites with deposits >50,000 years old, along with rock art of similar age. Gap in archeological record between ~20,000–10,000 years ago (dating of archeological breccias at Leang Bulu Bettue, in Maros Regency. Archeological breccias in caves important sources of information)

Newsome, J.C. (1985)- Late Quaternary vegetational history of the West Sumatran Highlands. Ph.D. Thesis, University of Hull, p. 1-583.
(online at: https://hydra.hull.ac.uk/assets/hull:12775a/content)
(Palynological investigation of sediments from montane West Sumatra, from shallow cores at Danau di Atas and Telago Climates formerly cooler, particularly from ~17.9- 12.9 kyrs, when vegetation of area included important gymnosperm element. During this cool period mean annual T 1.6-5.2°C cooler than present, but no evidence that climate was drier)

Newsome, J.C. (1988)- Late Quaternary vegetational history of the Central Highlands of Sumatra. I. Present vegetation and modern pollen rain. J. of Biogeography 15, 2, p. 363-386.

Newsome, J. & J.R. Flenley (1988)- Late Quaternary vegetational history of the Central Highlands of Sumatra. II. Palaeopalynology and vegetational history. J. of Biogeography 15, 4, p. 555-578.
(Late Quaternary pollen from two swamps at ~1500 m asl in Sumatran Highlands (Danau di Atas, Telago). In Late Pleistocene, montane forest vegetation zones depressed by ~800m and mean annual T was 1.6- 5.2 °C cooler than now)

Ngisomuddin, Akmaluddin & H. Amijaya (2007)- Benthic foraminifera as indicator of Recent tsunami deposit sources at Pangandaran coast, Ciamis and Parangendog coast, Yogyakarta. Proc. Joint Convention 36th IAGI, 32nd HAGI, and 29th IATMI, Bali 2007, p. 1110a-1110d. *(online at: https://www.iagi.or.id/web/digital/28/2007_IAGI-Bali_Benthic-Foraminifera-as-Indicator-of-Recent.pdf)*
(Recent tsunami deposit from Pangandaran with Ammonia, Elphidium, Amphistegina, Cibicides, Bigenerina. Bolivina, Bathysiphon, Nodosaria and Quinqueloculina, suggesting source from shallow to deep marine environents. Recent tsunami sediments at Parangendog Beach with Ammonia beccarii and Elphidium advenum, suggesting sediments came from lagoonal to shallow marine environment)

Noraswana, N.F., K.R. Mohamed & R. Omar (2017)- Species diversity and distribution of Ostracoda (Crustacea) in Sungai Kubang Badak, Langkawi, Kedah, Malaysia. Malayan Nature Journal 69, 2, p. 83-92.
(25 samples of estuarine- nearshore marine ostracod faunas of Sungai Kubang Badak river and estuary, N Langkawi. 16 genera and 27 species, most abundant species Hemicytheridea wangi and Neomonoceratina delicata. Three ostracod biofacies, (1) lower estuary and nearshore: dominated by H. wangi, N. delicata and Stigmatocythere indica; (2) upper estuary: lowest diversity, dominated by H. wangi and Xestoleberis malaysiana; (3) lower estuary: dominated by N. delicata and Hemicytheridea ornata)

Noraswana, Nor Faiz & R. Omar (2009)- Ostrakod baharu di dalam sedimen luar pantai di Sekitar Pulau Tioman, Pahang. Sains Malaysiana 38, 1, p. 9-20.
('Recent Ostracoda in offshore sediment around Pulau Tioman, Pahang'. 34 species of ostracodes in shallow waters around Tioman Island. Most common species Loxoconcha paiki, Pistocythereis bradyi, Venericythere papuensis)

Noraswana, N.F., Omar Ramlan & S. Norashikin (2014)- Distribution of Recent Ostracoda in offshore sediment of selected stations in the Sulu Sea. Sabah. Malaysian Applied Biology 43, 2, p. 49-57.
(online at: http://journalarticle.ukm.my/8679/1/43_2_07.pdf)
(32 species of Recent ostracods from 12 stations off eastern Sabah in 24-205m water depth. Dominant species Phlyctenophora orientalis)

Nugroho, S.H. (2018)- State of knowledge on marine palynology in Indonesia. Proc. Global Colloquium on GeoSciences and Engineering, Bandung 2017, IOP Conference Series: Earth and Environmental Science 118, 012012, p. 1-7.
(online at: http://iopscience.iop.org/article/10.1088/1755-1315/118/1/012012/pdf)
(Brief review of Quaternary (transported) spores-pollen studies in marine environments in Indonesia)

Nugroho, S.H., M.D. Sisca, P.S. Putra, S.W.A. Suedy & M. Izzati (2019)- Distribusi polen pada sedimen permukaan Bawah Laut di Perairan Sumba, Nusa Tenggara Timur. J. RISET Geologi dan Pertambangan (LIPI) 29, 1, p. 105-114.
(online at: https://jrisetgeotam.lipi.go.id/index.php/jrisgeotam/article/view/975/pdf)
(Distribution of pollen in 13 seafloor samples around Sumba Island helps interpret sedimentation effect of fluvial inputs, marine and terrestrial vegetation changes. Most dominant pollen from Family Poaceae)

Nugroho, S.H., Y. Zaim, E. Yulianto, Y. Rizal, A. Kurniasih, P. Sulastya Putra, S. Prasetyo A. Wibowo & A. Amar (2021)- A preliminary study in vertical distribution of planktonic foraminifera and marine ecological conditions of Simeulue sub-basin, Aceh, Indonesia. Bull. Geological Society Malaysia 72, p. 137-149.
(online at: https://gsm.org.my/wp-content/uploads/gsm_file_2/702001-101916-PDF.pdf)
(Quaternary planktonic foraminifera in 2m sediment core from 870m water depth in Simeulue sub-basin, off Aceh, W Sumatra forearc ('usual suspects': Globigerinoides ruber, Neogloboquadrina dutertrei, Pulleniatina obliquiloculata, Globigerina calida, Globigerinoides elongates, Gs cyclostomus, Gs sacculiferus, etc.))

Nurani, N., L. Jurnaliah & Winantris (2014)- Penentuan spesies foraminifera bentonik kecil dominan pada perairan Semarang, Provinsi Jawa Tengah. Bull. Scientific Contribution (UNPAD) 12, 1, p. 1-7.
(online at: http://jurnal.unpad.ac.id/bsc/article/view/8364/3885)
('Determination of small benthic foraminifera species dominant in the waters of Semarang, C Java Province'. Foraminifera in 20 samples from 38-54m in Java Sea N of Semarang dominated by Heterolepa (36%), Anomalina, Ammonia spp. (12%), Pseudorotalia, Quinqueloculina (9%) and Asterorotalia trispinosa (6%))

Nurdin, N., L. Gustiantini, S. Marina & M. Yosi (2014)- Analyses of foraminifers microfauna as environmental bioindicators in Kotok Besar, Kotok Kecil and Karang Bongkok Islands, Kepulauan Seribu, DKI Jakarta Province. Bulletin of the Marine Geology 29, 1, p. 21-28.
(online at: http://ejournal.mgi.esdm.go.id/index.php/bomg/article/view/62/63)
(Kepulauan Seribu (Thousand Islands) samples from coral reefs with ~20 genera of benthic foraminifera, dominated by Amphistegina and Calcarina, also Operculina, Quinqueloculina, Peneroplis, and Discorbis)

Nurdin, N. & I.R. Silalahi (2014)- Distribusi foraminifera bentik di perairan Aceh. J. Geologi Kelautan 12, 1, p. 25-31.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/243/233)
('Distribution of benthic foraminifera in Acaeh waters'. Shallow marine foraminifera distribution off NW tip of Sumatra, in water depths of 7-170m. 45 species of benthic foraminifera, dominated by Amphistegina papillosa. Ammonia tepida dominant in N part of Weh island)

Okada, H. (1983)- Modern nannofossil assemblages in sediments of coastal and marginal sea along the Western Pacific Ocean. Utrecht Micropaleontological Bulletin 30, p. 171-187.
(online at: https://dspace.library.uu.nl/bitstream/1874/205889/1/Utrecht+Micropaleontological+Bulletins-30-Meulenkamp.pdf)
(On distribution of modern calcareous nannofossils off Japan, Taiwan, Gulf of Thailand (mainly Emiliania huxleyi, Gephyrocapsa oceanica, Florisphaera profunda; G. oceanica most abundant in coastal stations) and Arafura Sea- Gulf of Carpenteria (mainly G. oceanica and E. huxleyi; F. profunda rel. rare). F. profunda dominates associations in deep basins, G. oceanica dominates floras in shallow marginal basins)

Omar, R. & N.N. Faiz (2014)- Distribution of Recent ostracoda in offshore sediment of the South China Sea. Pertanika J. Science Technology 22, 2, p.:433-444.
(Two faunal assemblages of ostracoda around S China Sea: (1) shallow water (Hemikrithe orientalis, Neomonoceratina iniqua, Stigmatocythere indica, Cytherelloidea leroyi, Neocytheretta snellii) and deep water (Paracypris sp., Alataconcha pterogona, Bythocytheropteron alatum, Keijella paucipunctata, Actinocythereis scutigera). Dominant species Foveoleberis cypraeoides)

O'Neill, M.C. (1986)- Distribution and ecology of Recent foraminifera from Pulau Pari (Pulau Seribu), West Java Sea. M.Sc. Thesis, p. 1-132.

Ongkosongo, O.S.R., S. Soeka & Susmiati (1979)- Foraminifera Resen dari daerah kehidupan hutan bakau di Teluk Ambon. Proc. Seminar Ekosistem Hutan Mangrove, Jakarta 1978, p. 129-138.
('Recent foraminifera from mangrove forests in Ambon Bay' (see also Suhartati Natsir 2010))

Oostingh, C.H. (1923)- Recent shells from Java, Part I. Gastropoda. Mededelingen Landbouwhoogeschool Wageningen 26, 3, Verhandelingen, p. 1-174.
(online at: https://edepot.wur.nl/294606)
(Description of Recent marine and freshwater gastropods from across Java, mainly collected by J. van Baren and J.M. Geerts)

Oostingh, C.H. (1925)- Report on a collection of Recent shells from Obi and Halmahera (Moluccas). Mededeelingen Landbouw-Hoogeschool Wageningen 29, 1, p. 1-362.
(online at: https://edepot.wur.nl/294577)
(Systematic descriptions of collection of rent shells from Obi Major and Halmahera. No localities or illustrations?))

Oostingh, C.H. (1931)- Beitrage zur Kenntnis der Molluskenfauna von Sud Sumatra. Archiv fur Molluskenkunde 63, 4-6, p. 166-255.
(online at: https://www.zobodat.at/pdf/Archiv-fuer-Molluskenkunde_63_0166-0255.pdf)
('Contributions to the knowledge of the mollusc fauna of S Sumatra'. Listings of recent molluscs from southernmost Sumatra (S tip of Bengkulu and along Sunda straits)

Oppenoorth, W.F.F. (1920)- het gebruik der microfauna voor de kennis der stratigrafie van het jong-tertiair ten behoeve van olie-opsporingen. Mededelingen Algemeen Ingenieurs Congres, Batavia 1920, 5, Mijnbouw en Geologie, Prae-advies 5, p. 3-4.
('The use of microfauna for the knowledge of stratigraphy of the Late Tertiary, as needed for oil exploration'. Brief pamphlet promoting use of foraminifera for oilfield stratigraphic subdivision))

Orpin, A.R., D.W. Haig & K.J. Woolfe (1999)- Sedimentary and foraminiferal facies in Exmouth Gulf, in arid tropical northwestern Australia. Australian J. Earth Sciences 46, 4, p. 607-621.
(Recent foraminifera distribution in shallow water grab samples, Exmouth Gulf (generally <20m). Benthic foraminifera dominant; planktonics only 1-2%. Six cluster groups, mainly controlled by bottom sediment type)

Palmieri, V. (1976)- Modern and relict foraminifera from the central Queensland continental shelf. Queensland Government Mining J. 77, p. 406-423.

Pam, B.& P. Mohan. (2022)- Distribution of Larger Benthic Foraminifera (LBF) in selected islands of Marine National Park, Port Blair, South Andaman, India. Open J. Marine Science, 12, 4, p. 141-160.
(online at: https://www.scirp.org/pdf/ojms_2022101714282600.pdf)
(22 taxa of modern LBF in southern Andaman islands. Most common families Amphisteginidae, Calcarinidae, Nummulitidae, Peneroplidae and Soritidae. With listings of preferred water depth, substrate, symbonts, etc.)

Panchang, R. & R. Nigam (2012)- High resolution climatic records of the past ~489 years from Central Asia as derived from benthic foraminiferal species, *Asterorotalia trispinosa*. Marine Geology 307-310, p. 88-104.
(Recent benthic foraminiferal distribution on Ayeyarwaddy Delta Shelf, off Myanmar shows Asterorotalia trispinosa has preference for low salinity. Variations in abundance used as proxy for delineation of past wet and dry periods: dry climate prior to 1650 AD and warm and wet climate since 1650 to present)

Parker, J.H. (2009)- Taxonomy of foraminifera from Ningaloo Reef, Western Australia. Memoir Assoc. Australasian Palaeontologists (AAP) 36, p. 1-810.
(Comprehensive inventory of modern foraminifera from 334 samples in 0-34m water depth of subtropical Ningaloo Reef in E Indian Ocean, off NW Australia. Descriptions and illustrations of 404 species)

Parker, J.H. & E. Gischler (2011)- Modern foraminiferal distribution and diversity in two atolls from the Maldives, Indian Ocean. Marine Micropaleontology 78, p. 30-49.
(Two coral-reef lagoons comprise eight foraminifera assemblages with 270 species. Three assemblages reefal and dominated by Amphistegina and Calcarina. One lagoon assemblage with abundant Ammonia and smaller miliolids. Species diversity in Maldives higher than W Indian Ocean, but not as high as central Indo-Pacific)

Pflaumann, U. & Z. Jian (1999)- Modern distribution patterns of planktic foraminifera in the South China Sea and Western Pacific: a new transfer technique to estimate regional sea-surface temperatures. Marine Geology 156, 1, p. 41-83.

Pickett, E.J., S.P. Harrison, G. Hope, K. Harle, J.R. Dodson, A.P. Kershaw, I.C. Prentice, J. Backhouse et al. (2004)- Pollen based reconstructions of biome distributions for Australia, Southeast Asia and the Pacific (SEAPAC region) at 0,6000 and 18,000 14C yr BP. J. of Biogeography 31, p. 1381-1444.
(Reconstruction of vegetation patterns in Australia and SE Asia in mid-Holocene and Last Glacial Maximum from pollen data. At 18ka SEAPAC region drier than today and, in tropics, colder)

Pirrung, M., G. Buechel, U. Schulte-Vieting, B.W. Scharf & H.C. Treutler (2005)- Dating of lacustrine sediments from the tropical maar Ranu Klindungan, East Java (Indonesia). Zeitschrift Deutschen Gesellschaft Geowissenschaften 156, 4, p. 557-571.
(On Late Quaternary lake sediments from 126 m deep volcanic maar lake near Pasuruan, E Java. Bottom sediments down to 1.06 m sediment depth consist of calcareous diatom gyttja with frequent turbidites. Seasonal variations of terrigenic layers deposited during rainy season (November–April) and diatom gyttja layers with thin distal turbidites, deposited during dry season)

Polhaupessy, A.A. (2009)- Pollen dispersal into the Liang Bua cave, Ruteng, Manggarai region, West Flores. Jurnal Sumber Daya Geologi (JSDG) 14, 3 (147), p. 113-117.
(Quaternary pollen grains in Liang Bua cave derived mainly from local upland vegation around cave)

Polhaupessy, A.A. (2009)- Palynology of Togi Ndrawa cave, coastal area of Nias Island, North Sumatera. Bulletin of the Marine Geology 24, 2, p. 99-115.
(online at: http://ejournal.mgi.esdm.go.id/index.php/bomg/article/view/19/19)
(Spores-pollen in 10 samples from Pleistocene- Holocene cave deposits on Nias)

Poliakova, A. (2015)- The Late Holocene history of vegetation, climate, fire dynamics and human impacts in Java and Southern Kalimantan. Doct. Thesis Georg-August Universitat, Gottingen, p. 1-186.
(online at: https://d-nb.info/1113875569/34)
(Collection of papers/ manuscripts of palynological studies in four shallow cores in Holocene deposits in Java Sea off NE Java (Solo River) and S Kalimantan (off Jelai and Pembuang rivers))

Poliakova, A. & H. Behling (2015)- Pollen and fern spores recorded in recent and late Holocene marine sediments from the Indian Ocean and Java Sea in Indonesia. Quaternary International 392, p. 251-314.
(Documentation of Quaternary and Recent pollen (138) and spores (41) in cores from coastal regions of Java Sea off Pembuang River in SW Kalimantan and NE Java and in marine sediment trap off W Java)

Poliakova, A., K.A.F. Zonneveld, L.S. Herbeck, T.C. Jennerjahn, H. Permana, C. Kwiatkowski & H. Behling (2017)- High resolution multi-proxy reconstruction of environmental changes in coastal waters of the Java Sea, Indonesia, during the late Holocene. Palynology 41, 3, p. 297-310.
(A 134-cm-long sediment core from ~50 km off Pembuang River mouth, S Kalimantan. Mixed terrestrial and marine organic matter, with low pollen-spore concentrations. Dinoflagellate cysts mainly Operculodinium and Spiniferites with minor Impagidinium (mainly I. strialatum). After ca. 2480 cal yr BP, bottom waters became increasingly ventilated. After 1530 cal yr BP, more pronounced influence of Pembuang River indicated by nutrient-sensitive Lingulodinium machaerophorum and Nematosphaeropsis labyrinthus)

Post, A.L., L. Sbaffi, V. Passlow & D.C. Collins (2009)- Benthic foraminifera as environmental indicators in Torres Strait- Gulf of Papua. In: B.J. Todd & H.G. Greene (eds.) Mapping the seafloor for habitat characterization, Geolpgical Association of Canada (GAC), Special Paper47, p. 329-348.
(online at: https://www.researchgate.net/publication/286972215_Benthic_foraminifera_as_environmental_indicators_in_Torres_Strait-Gulf_of_Papua)
(Study of benthic forams along transect from Fly River Delta to shelf edge (~140m depth), near N end of Great Barrier Reef. Three areas different benthic foram assemblages. High relict content in surface samples)

Prayudi, S.D., Suhartati M. Natsir & M.A. Kaminski (2025)- Extant benthic foraminifera from the Indonesian seas: An update of what we know so far and implications for future studies. Marine Micropaleontology 197-198, 102471, p.
(Benthic foraminifera from Indonesian waters have been studied since the Challenger Expedition in 1876. To date, 1164 species of Recent benthic foraminifera documented here, belonging to 448 genera. Numerous instances of taxonomic misnaming)

Pudjoarinto, A. (1999)- Palynological evidence for environmental change in Dieng Highland. Indonesian J. Geography 31, 77-78, p. 11-24.
(Palynology of 18m core from small lake in Dieng Plateau, C Java. Montane forest assemblages record climate changes)

Putra, P.S. & S.H. Nugroho (2020)- Holocene climate dynamics in Sumba Strait, Indonesia: a preliminary evidence from high resolution geochemical records and planktonic foraminifera. Studia Quaternaria 37, 2, p. 91-99.
(online at: https://journals.pan.pl/Content/116874/PDF/4_Putra.pdf)
(Planktonic foraminifera in 243cm drop core ST14 in 1283m water depth in Sumba Strait. Base Holocene determined from first occurrence of Globigerina calida calida at 156 cm. Six Holocene climate zones (warm-humid versus cold-dry episodes) identified. Cool water Globigerina bulloides more abundant in Pleistocene than in Holocene sediment. Globorotalia menardii, Gr. tumida, Pulleniatina obliquiloculata peaks in warm humid zones. Etc.)

Rai, A.K. & V.B. Singh (2001)- Late Neogene deep-sea benthic foraminifera at ODP Site 762B, eastern Indian Ocean: diversity trends and palaeoceanography. Palaeogeogr. Palaeoclim. Palaeoecology 173, p. 1-8.
(Pliocene-Pleistocene deep sea benthic foraminifera from ODP Site 762B off Exmouth Plateau. Species diversity inverse relationship with abundance of Uvigerina proboscidea and precentage infaunal taxa))

Rai, A.K. & V.B. Singh (2012)- Response of eastern Indian Ocean (ODP Site 762B) benthic foraminiferal assemblages to the closure of the Indonesian seaway. Oceanologia 54, 3, p. 449-472.
(Pliocene-Pleistocene deep sea benthic foraminifera from ODP Site 762B off Exmouth Plateau in E Indian Ocean. Diverse fauna in E Pliocene (>3.5 Ma) relatively oligotrophic and warm bottom water conditions. At beginning of Late Pliocene (i.e. ~ 3 ± 0.5 Ma) increase in Uvigerina proboscidea, infaunal taxa and high productivity taxa and decline in faunal diversity suggest development of pronounced upwelling. Reduced inflow of warm and oligotrophic water masses from SW Pacific to E Indian Ocean due to effective closure of Indonesian seaway increased surface water productivity. Closing of Indonesian seaway in Late Pliocene also responsible for cessation of warm, S-flowing Leeuwin Current)

Rajshekhar, C. (2013)- The Late Holocene foraminifera from Andaman Islands, Andaman Sea, Bay of Bengal. In: K. Venkataraman et al. (eds.) Ecology and conservation of tropical marine faunal communities, Springer Verlag, p. 3-18.
(S Andaman Island three distinct environments: rocky shore with Elpidium, Amphistegina; (2) sandy shore with common Calcarina and (3) intertidal muddy region with Trochammina inflata is common in intertidal clays)

Ramlan, Omar & Noraswana Nor Faiz (2014)- Distribution of recent ostracoda in offshore sediment of the South China Sea. Pertanika J. Science and Technology 22, 2, p. 433-444.
(online at: http://www.pertanika.upm.edu.my/pjst/browse/archives?article=JST-0379-2012)

(30 sediment samples in S China Sea off eastern Malay Penisula contain 79 species of ostracods. Dominant species Foveoleberis cypraeoides. Two faunal assemblages: shallow water (<50m) (Hemikrithe orientalis, Neomonoceratina iniqua, Stigmatocythere indica, Cytherelloidea leroyi and Neocytheretta snellii) and deep water (>50m) (Paracypris sp., Alataconcha pterogona, Bythocytheropteron alatum, Keijella paucipunctata and Actinocythereis scutigera).

Rao, N.R., M. Jayaprakash & P.M. Velmurugan (2013)- The ecology of *Asterorotalia trispinosa* (Thalmann, 1933)- new insights from Muthupet Lagoon, Southeast Coast of India. J. Foraminiferal Research 43, p. 14-20.
(Asterorotalia trispinosa dominates sparse foraminiferal assemblage of shallow, mud-dominated Muthupet Lagoon on SE coast of India, where salinities are slightly lower than normal marine)

Rasheed, D.A. (1968)- Distribution of foraminifera in the Coral Sea, South of Papua, New Guinea. J. of the Madras University B37-38, p. 73-80.

Rasheed, D.A. (1968)- Some foraminifera belonging to Miliolidae and Opthalmididae from the Coral Sea, south of Papua, New Guinea. J. of the Madras University B37-38, p. 19-68.

Rasheed, D.A. (1970)- Some Recent arenaceous foraminifera from the Coral Sea. south of Papua, New Guinea. J. of the Madras University B39-40, p. 41-58.

Rasheed, D.A. (1970)- Some recent calcareous foraminifera belonging to the families Peneroplidae. Alveolinellidae, Lagenidae and Polymorphinidae from the Coral Ssea, South of Papua (New Guinea). J. of the Madras University B39-40, p. 77-110.

Rasheed, D.A. (1970)- Some calcareous foraminifera belonging to the families Rotaiiidae, Cymboloporidae, Anomalinidae, Calcarinidae, Globigerinidae and Bulinidae from the Coral Sea, South of Papua (New Guinea). J. of the Madras University B 39-40, p. 150-201.

Rasheed, D.A. (1971)- Some foraminifera belonging to Miliolidae and Ophthalmidiidae from the Coral Sea, South of Papua (New Guinea), Part II. J. of the Madras University B37-38, p. 150-201.

Rathburn, A.E. & B.H. Corliss (1994)- The ecology of living (stained) deep-sea benthic foraminifera from the Sulu Sea. Paleoceanography 9, p. 87-150.
(Deep-sea benthic foraminifera distibution in 8 box cores from 510-4515 m in thermospheric (>10°C) Sulu Sea. Site 510m dominated by Cibicidoides, Uvigerina, and Bolivina. 1005m core: mainly Siphonina Cibicidoides, Uvigerina. 2000m cores: Cibicidoides, Gyroidinoides, and Oridorsalis. 3000 and 4000m cores: Cibicidoides bradyi and Oridorsalis umbonatus dominant. Infaunal Valvulineria mexicana in sediments of 4515m core. Low bottom water oxygen values do not necessarily yield 'low-oxygen taxa' like Bolivina, Uvigerina, Chilostomella, Bulimina and Globobulimina)

Rathburn, A.E., B.H. Corliss, K.D. Tappa & K.C. Lohmann (1996)- Comparisons of the ecology and stable isotopic compositions of living (stained) benthic foraminifera from the Sulu and South China Seas. Deep Sea Research 1, 43, 10, p. 1617-1646.
(Significant differences between living deep-sea benthic foraminifera in thermospheric (>10°C) environments of Sulu Sea and psychrospheric (<10°C) conditions in S China Sea. Gavelinopsis, Bolivinopsis, Astrononion, Osangularia and Ceratobulimina common taxa in S China Sea, but rare in Sulu Sea. Siphonina and Valvulineria dominant genera at certain depths in Sulu Sea, but rare in S China Sea. Differences result from large differences of bottom-water temperatures)

Rathburn, A.E. & Q. Miao (1995)- The taphonomy of deep-sea benthic foraminifera: comparisons of living and dead assemblages from box and gravity cores taken in the Sulu Sea. Marine Micropaleontology 25, p. 127-149.
(Benthic foraminifera from 500- 4000m water in Sulu Sea. Bolivina, Bulimina, Globobulimina, Chilostomella and Uvigerina most abundant <1500m, but rel. rare in deeper water. Dominant taxa below 2000m Cibicidoides bradyi and Oridorsalis umbonatus)

Razak, N.S.A., H. Shaari, F.I. Minhat, E.H. Ariffin, F. Shariful, S. Yang, R. Suriadi & H.M. Sharif (2023)- Benthic foraminifera as environmental indicators in the lagoon and mangrove environments of Langkawi, Malaysia. Marine Geology 462, 107095, p. 1-11.
(Foraminiferal distribution in the surface sediments of Sungai Kilim in Kilim Geoforest Park, Pulau Langkawi, NW Malaysia. Twenty-eight genera of benthic foraminifera from surface samples. Elphidium (13-83%) most abundant in mangrove and lagoon environments.Other common genera Trochaminna, Ammotium, Ammobaculites and Haplophragmoides)

Reeves, J.M. (2004)- The use of ostracoda in the palaeoenvironmental reconstruction of the Gulf of Carpenteria, Australia, from the last interglacial to present. Ph.D. Thesis University of Wollongong, p. 1-447.
(online at: http://ro.uow.edu.au/cgi/viewcontent.cgi?filename=0&article=1211&context=theses&type=additional)

Reeves, J.M., A.R. Chivas, A. Garcia & P. De Deckker (2007)- Palaeoenvironmental change in the Gulf of Carpentaria (Australia) since the last interglacial based on Ostracoda. Palaeogeogr. Palaeoclim. Palaeoecology 246, p. 163-187.
(Throughout last glacial cycle, region between Australia and New Guinea (now Gulf of Carpentaria) oscillated from open shallow marine conditions to freshwater lake behind Arafura sill. Six ostracod biofacies in last 130 ka: (1) open shallow marine with bairdiids, pectocytherinids, cytherettids; (2) shallow marine dominated by Cytherella and Hemikrithe; (3) marginal marine with Xestoleberis and Praemunita; (4) tidal channel dominated by Loxoconcha; (5) estuarine with Venericythere and Leptocythere; (6) non-marine facies: brackish lagoon/lake dominated by Cyprideis and Leptocythere and freshwater with Ilyocypris, Cyprinotus and Cypretta. Also morphological variations within species tied to paleoenvironments)

Renault-Miskovsky, J. & A.M. Semah (1998)- Palynology of the Quaternary in temperate and tropical areas: chronostratigraphy, palaeoclimatology and vegetal environment of fossil man. In: N.M. Dutta et al. (eds.) Current concepts in pollen-spore and biopollution research (S. Chandra volume), Chapter 20, Research Periodicals and Book Publishing House, Houston, p. 297-317.
(online at: https://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_7/b_fdi_53-54/010021194.pdf)
(General review of spores-pollen studies and applications in chronostratigraphy, paleoclimate and paleoenvironments. With examples from SE Asia, including work on Java Pleistocene hominid sites and Pleistocene climate changes on New Guinea, Sumatra, etc.)

Renema, W. (2003)- Larger foraminifera on reefs around Bali. Zoologische Verhandelingen, Leiden, 345, p. 337-366.
(online at: www.repository.naturalis.nl/document/46850)
(Recent larger foraminifera from Bali total 19 species. Species richness similar to SW Sulawesi and Cebu, but different composition. Schlumbergerella locally abundant and geographically restricted to Lesser Sunda Islands and Java. Very low abundance of imperforate species probably due to climatic or oceanographic parameters, most likely periodic upwelling, which causes seasonal seawater temperature drops)

Renema, W. (2005)- Depth estimation using diameter-thickness ratios in larger benthic foraminifera. Lethaia 38, p. 137-141.
(Diameter-thickness ratio (D/T) of Amphistegina and Operculina varies with depth. Increased turbulence thickens test, whilst decreased light intensity causes flatter tests)

Renema, W. (2006)- Large benthic foraminifera from the deep photic zone of a mixed siliciclastic-carbonate shelf off East Kalimantan, Indonesia. Marine Micropaleontology 58, p. 73-82.
(Modern large benthic forams on Berau shelf two-three depth-controlled assemblages (1) shallow (20-50m; dominated by Operculina ammonoides) and (2) deeper (50-85m; dominated by Operculina complanata and Planostegina operculinoides). Deepest living LBF at 115m. Cycloclypeus carpenteri between 55-95m)

Renema, W. (2006)- Habitat variables determining the occurrence of large benthic foraminifera in the Berau area (East Kalimantan, Indonesia). Coral Reefs 25, 3, p. 351-359.
(Composition of larger foram assemblages (35 species) on Berau carbonate shelf with barrier reef system and some reefs outside barrier. Four clusters corresponding to substrate type)

Renema, W. (2008)- Habitat selective factors influencing the distribution of larger benthic foraminiferal assemblages over the Kepulauan Seribu. Marine Micropaleontology 68, p. 286-298.
(On distribution of symbiont-bearing larger foraminifera on 'Thousand Islands' off Jakarta. Diversity and habitat fractionation increases as terrestrial and nutrient influence decline. Assemblages in nearshore reefs dominated by generalist species, while, additionally, more specialistic species occur at more offshore reefs)

Renema, W. (2010)- Is increased calcarinid (foraminifera) abundance indicating a larger role for macro-algae in Indonesian Plio-Pleistocene coral reefs? Coral Reefs 29, p. 165-173.
(Comparisons of large benthic foraminifera between areas with high (Kepulauan Seribu) and lower (Spermonde Archipelago) anthropogenic influence. Reefal habitats dominated by algae are inhabited by Calcarinidae larger foraminifera)

Renema, W. (2018)- Terrestrial influence as a key driver of spatial variability in large benthic foraminiferal assemblage composition in the Central Indo-Pacific. Earth-Science Reviews 177, p. 514-544.
(online at: https://www.sciencedirect.com/science/article/pii/S0012825217303410)
(Thorough review of Recent Indo-Pacific larger foraminifera and their distributions. Nutrient availability, hydrodynamic energy, light intensity, substrate type and temperature important environmental parameters controlling occurrence of larger foraminifera species. Terrestrial run-off also important environmental parameter. Calcarinids more abundant with increased algal cover)

Renema, W., R.J. Beaman & J.M. Webster (2013)- Mixing of relict and modern tests of larger benthic foraminifera on the Great Barrier Reef shelf margin. Marine Micropaleontology 101, p. 68-75.
(On distribution of live, well preserved, and relict larger foraminifera off shelf margin near Hydrographers Passage in Great Barrier Reef, Australia. Eight living species, mostly of families Amphisteginidae and Nummulitidae, separated into a shallow (platform) and deep (terrace) assemblages. Relict fauna more diverse, with taxa from much shallower settings (downslope transport from nearby Pleistocene coral reef structures))

Renema, W., D.R. Bellwood, J.C. Braga, K. Bromfield, R. Hall, K.G. Johnson, P. Lunt et al. (2008)- Hopping hotspots: global shifts in marine biodiversity. Science 321, p. 654-657. *(online at: www.researchgate.net/publication/23140202_Hopping_Hotspots_Global_Shifts_in_Marine_Biodiversity)*
(Fossil and molecular evidence reveals at least three hotspots of high marine biodiversity in past 50 million years. They moved across globe, with timing and locations coinciding with major tectonic events. Birth and death of successive hotspots highlights link between environmental change and biodiversity patterns. Antiquity of taxa in modern Indo-Australian Archipelago hotspot emphasizes role of pre-Pleistocene events)

Renema, W., B.W. Hoeksema & J.E. van Hinte (2001)- Larger benthic foraminifera and their distribution patterns on the Spermonde shelf, South Sulawesi. Zoologische Verhandelingen, Leiden, 334, p. 115-149.
(online at: www.repository.naturalis.nl/document/46294)
(Distribution patterns of 20 species of larger benthic foraminifera in Spermonde Archipelago, off SW Sulawesi. 13 transects sampled, down to 33m water depth. Substrate type, hydrodynamic energy, light intensity, nutrient availability and environmental stabiltity determine distribution)

Renema, W. & J. Hohenegger (2005)- On the identity of *Calcarina spengleri* (Gmelin 1791). J. Foraminiferal Research 35, 1, p. 15-21.
(online at: https://www.researchgate.net/publication/236849648_On_the_identity_of_Calcarina_Spengleri_GMELIN_1791)
(On identity of Recent reef dwelling larger foram Calcarina spengleri (Gmelin 1791). Commonly confused with Calcarina mayori Cushman 1924, C. gaudichaudii d'Orbigny 1840 and C. hispida Brady 1876)

Renema, W. & S.R. Troelstra (2001)- Larger foraminifera distribution on a mesotrophic carbonate shelf in SW Sulawesi (Indonesia). Palaeogeogr. Palaeoclim. Palaeoecology 175, p. 125-146.
(online at: https://www.academia.edu/28094197/Larger_foraminifera_distribution_on_a_mesotrophic_carbonate_shelf_in_SW_Sulawesi_Indonesia_)
(Modern larger foram distribution on Spermonde Shelf)

Robba, E., L.D. Geronimo, N. Chaimanee, M. Negri & R. Sanfilippo (2007)- Holocene and Recent shallow soft-bottom molluscs from the Western Gulf of Thailand: Pak Phanang Bay, and additions to Phetchchaburi fauna. Bolletino Malacologico 42, Supplement 5; p. 1-98.
(online at: https://www.researchgate.net/publication/235087893_Holocene_and_Recent_shallow_soft-bottom_mollusks_from_the_Western_Gulf_of_Thailand_Pak_Phanang_Bay_and_additions_to_Phetchaburi_fauna)
(>150 species of gastropods and bivalves from Holocene Bangkok Bay from modern shallow bottoms in Phetchaburi coastal area)

Robles, E. (2007)- Palynological investigation of a laminated sediment core from Lake Guyang Warak, Java, Indonesia. Masters Thesis Museum National Histoire Naturelle, Paris, p. 1-53. *(Unpublished)*
(online at: http://hopsea.mnhn.fr/pc/thesis/Emil_Robes_2007.pdf)
(Palynology of Quaternary lake deposits on karst surface of Gunung Sewu (Southern Mountains), C Java)

Romero, O.E., M. Mohtadi, P. Helmke & D. Hebbeln (2012)- High interglacial diatom paleoproductivity in the westernmost Indo-Pacific Warm Pool during the past 130,000 years. Paleoceanography 27, PA3209, p. 1-14.
(online at: http://onlinelibrary.wiley.com/doi/10.1029/2012PA002299/epdf)
(Late Pleistocene diatoms in core GeoB10038-4 off S Sumatra show highest paleoproductivity during interglacials, due to nutrient input after rise in sea level. In Marine Isotope Stage 5 response of diatom productivity and upwelling intensity to boreal summer insolation. Resting spores of Chaetoceros, typical of nutrient-rich waters, dominant during periods of highest diatom paleoproductivity)

Rottger, R., R. Kruger & S. de Rijk (1990)- Larger Foraminifera; variation in outer morphology and prolocular size in *Calcarina gaudichaudii.* J. Foraminiferal Research 20, 2, p. 170-174.
(online at: http://jfr.geoscienceworld.org/content/20/2/170.full.pdf)
(Study of larger foram Calcarina gaudichaudii, abundant in high-energy shallow reefal facies of W Pacific. Based on material from Adorius island, Micronesia and Komodo island, Indonesia))

Rositasari, R. (1990)- *Calcarina* sebagai genus penciri lingkungan terumbu karang Resen. Proc. 17th Annual Conv. Indonesian Association Geologists (IAGI), Jakarta 1988, p. 79-85.
(online at: https://www.iagi.or.id/web/digital/46/17th-volume-(-12-15-Des-1988-)-82-89.pdf)
(On Calcarina larger foram in Recent reefal limestone deposits. Calcarina spp. dominant on Recent reefs of Pulau Seribu (Calcarina calcar, C. spengleri, C. venusta))

Rositasari, R. (1993)- Foraminifera sebagai bioindikator lingkungan tercemar. Proc. 22nd Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 2, p. 1091-1199.
(online at: https://www.iagi.or.id/web/digital/53/22nd-Volume-2-(6-9-Des-1993)-489-497.pdf)
('Foraminifera as bioindicators for contaminated deposits'. Foraminifera from Jakarta Bay and Sunda Straits)

Rositasari, R. (2010)- Recent foraminifera communities in Makassar Strait. J. Coastal Development 14, p. 26-34.
(online at: https://scispace.com/pdf/recent-foraminiferal-communities-in-makassar-strait-26gpolhhtb.pdf)
(Globigerina ooze forms major fraction of bottom sediment in Makassar Strait. Diversity and richness of benthic foraminifera decrease with water depth. High abundance of Uvigerina asperula may be proxy of oxygen minimum zone)

Rositasari, R. (2010)- Karakteristik komunitas foraminifera di perairan Teluk Jakarta. J. Ilmu Teknologi Kelautan Tropis (JITKT; Bogor) 3, 2, p. 100-111.

(online at: http://repository.ipb.ac.id/bitstream/handle/123456789/53434/08%20Karakteristik%20Komunitas.pdf)
(The characteristics of the foraminiferal community in Jakarta Bay'. Benthic foraminifera in Jakarta Bay: (1) coastal water and estuary dominated by Ammonia beccarii; (2) Calcarina and other larger foraminifera common in coral reef area; (3) Elphidium and Nonion depressulum common in open waters area. Higher diversity than coastal water of Semarang and Cirebon. Reef area has highest diversity)

Rositasari, R. & L. Effendi (1994)- Foraminifera aglutinin dan kemungkinan pengaplikasiannya sebagai indikator lingkungan yang mengalami tekanan. Proc. 23rd Annual Conv. Indonesian Association Geologists (IAGI), Jakarta, 1, p. 155-161.
(online at: https://www.iagi.or.id/web/digital/62/14.pdf)
('Agglutinated foraminifera and possible application as indicators of environments under stress'. Discussion of agglutinated foraminifera and presence in shallow, brackish waters of coastal creeks around Jakarta Bay. Main taxa Eggerella, Ammobaculites agglutinans, Trochammina, Haplophragmoides, Textularia conica, Cyclammina pusilla, etc.)

Rositasari, R., Suhartati M.N., T. Susana & Helfinalis (1994)- Tipe estuari sebagai faktor pembatas pada komunitas foraminifera; hasil penelitian di Muara Sungai Ciawi dan Muara Sungai Bekasi. Proc. 23rd Annual Conv. Indonesian Association Geologists (IAGI), Jakarta, 1, p. 162-172.
(online at: https://www.iagi.or.id/web/digital/62/15.pdf)
('Type of estuary as factor on foraminifera communities; results of study at mouths of Ciawi and Bekasi Rivers'. Ciawi river mouth (Ujung Kolon) with abundant Operculina ammonoides, Amphistegina lessonii, Calcarina calcar, Elphidium, Quinqueloculina, etc. Bekasi river mouth dominated by Ammonia beccarii, Trochammina hadai and Ammobaculites agglutinans)

Rottman, M.L. (1979)- Dissolution of planktonic foraminifera and pteropods in South China Sea sediments. J. Foraminiferal Research 9, p. 41-49.
(Pteropods common in Recent sediments of S China Sea (but aragonitic tests highly susceptible to dissolution, so rarely fond as fossils))

Rottman, M.L. (1980)- Net tow and surface sediment distributions of pteropods in the South China Sea region: comparison and oceanographic implications. Marine Micropaleontology 5, p. 71-110.
(Distributions of Recent aragonitic pelagic pteropod species on Sunda Shelf and in Java Sea very similar, but some exceptions)

Rowe, C., I.J. McNiven, B. David, T. Richards & M. Leavesley (2013)- Holocene pollen records from Caution Bay, southern mainland Papua New Guinea. The Holocene 23, 8, p. 1130-1142.
(Palynological data from tide-dominated shoreline of Caution Bay, W of Port Moresby, PNG, suggest late-Holocene mangrove to mudflat transition)

Rugmai, W., P.J. Grote, C. Chonglakmani, R. Zetter & D.K. Ferguson (2008)- A Late Pleistocene palynoflora from the coastal area of Songkhla Lake, southern Thailand. Science Asia 34, p. 137-145.
(online at: http://scienceasia.org/2008.34.n2/scias34_137.pdf)

Rymer-Jones, F.W.O. (1874)- On some Recent forms of Lagenae from deep-sea soundings in the Java Seas. Transactions Linnean Society, London, 30, 1, p. 45-69.
(online at: https://www.biodiversitylibrary.org/page/27557854#page/66/mode/1up)
(On many 'varieties' of Lagena vulgaris (incl. Oolina) from seafloor samples at 1080 fathoms, 10 miles S of 'Sandalwood Island' (= Sumba?, but lat-longs closer to South Bali; unlikely from Java Sea). Many now viewed as variations of single species. Associated with common Rotalia, Uvigerina, Bulimina, Globigerina, diatoms, ostracods, sponge needles, etc.)

Sahidin, A., G. Muhammad, Z. Hasan, M.C.W. Arief, R.M. Marwoto & A. Komaru (2021)- Indonesian freshwater bivalves: a meta-analysis of endemicity, ecoregion distributions, and conservation status. AACL Bioflux 14, 6, p. 3750-3775.
(online at: http://bioflux.com.ro/docs/2021.3750-3775.pdf)
(Geographic distribution of freshwater bivalve molluscs in Indonesia. Three main ecoregions: 1) Sundaland (Sumatra, Java, and Kalimantan), dominated by Unionidae; 2) Wallacea (Sulawesi), dominated by Cyrenidae; and 3) the Sahul region (Papua): hotspot of the Hyriidae)

Sakai, K. & M. Nishimura (1980)- Population study of the benthic foraminifer *Baculogypsina sphaerulata* on the Okinawan reef flat and preliminary estimation of its annual production. Proc. Fourth International Coral Reef Symposium, Manila, 2, p. 736-766.

Saputro, E., L. Fauzielly, I.R. Silalahi & Winatris (2019)- Distribusi spasial foraminifera di Perairan Teluk Cenderawasih, Papua Barat. J. Geologi Kelautan, 17, 2, p. 77-88.
(online at: https://core.ac.uk/download/pdf/287301008.pdf)
('Distribution of foraminifera in the waters of Cenderawasih Bay, West Papua'.Foraminifera from 20 bottom samples between 87 and 1266m water depth, with 13 species of planktonic foraminifera and 87 species of benthic foraminifera)

Sarasin, P. & F. Sarasin (1897)- Ueber die Molluskenfauna der grossen Susswasser Seen von Central-Celebes. Zoologischer Anzeiger 20, 536, p. 241-245.
('On the mollusc fauna of the large freshwater lakes of Central Sulawesi')

Saraswat, R., M. Manasa, T. Suokhrie, M.S. Saalim & R. Nigam (2017)- Abundance and ecology of endemic *Asterorotalia trispinosa* from the western Bay of Bengal: implications for its application as a paleomonsoon proxy. Acta Geologica Sinica (English Ed.) 91, 6, p. 2268-2282.
(manuscript online at: https://drs.nio.res.in/drs/bitstream/handle/2264/7753/Acta_Geolo_Sinica_91_2268.pdf)
(In samples from continental shelf and slope of W Bay of Bengal Asterorotalia trispinosa abundance ranges from 0- 31%, with highest abundance near outfall region of Ganges-Brahmaputra Rivers and decreases away from river mouths. Abundance of A. trispinosa indicates warmer and marginally hyposaline environment)

Sathyanarayana, B., M.L. Husain, R. Ibrahim, S. Ibrahim & F.D. Guebas (2014)- Foraminiferal distribution and association patterns in the mangrove sediments of Kapar and Matang, West Peninsular Malaysia. J. Sustainability Science Management 9, p. 32-48.
(online at: www.ulb.ac.be/sciences/biocomplexity/pub/Satyanarayanaetal_2014_JSustainSciManage.pdf)
(28 foram species in mangrove surface sediments on W coast of Malay Peninsula. Calcareous forms mainly Ammonia beccarii and Buccella frigida. Agglutinated species mainly Arenoparrella and Haplophragmoides)

Sawai, Y., K. Jankaew, M.E. Martin, A. Prendergast, M. Choowong &, T. Charoentitirat (2009)- Diatom assemblages in tsunami deposits associated with the 2004 Indian Ocean tsunami at Phra Thong Island, Thailand. Marine Micropaleontology 73, 1, p. 70-79.
(Diatom assemblages in fining-upward m-f sandy deposits of 2004 tsunami at Phra Thong Island, Thailand: (1) lowermost sand mainly unbroken beach and subtidal species that live attached to sand grains; (2) shift to marine planktonic species in middle of the tsunami deposit and (3) mix of freshwater, brackish, and marine species near top. Trends are consistent with expected changes in current velocities of tsunami through time)

Shen, L., M. Chen, B. Lan, H. Qi, A. Zhang, D. Lan & Qi Fang (2017)- Diatom distribution as an environmental indicator in surface sediments of the West Philippine Basin. Chinese J. Oceanology Limnology 35, 2, p. 431-443.
(Distribution of oceanic diatoms in W Philippine Basin. Ethmodiscus rex dominant species. 68 species in 4 assemblages, related with North Equatorial Current and Kuroshio Current patterns)

Schepman, M.M. (1908)- The Prosobranchia of the Siboga Expedition, Part I. Rhipidoglossa and Docoglossa. In: M. Weber (ed.) Siboga Expeditie, Uitkomsten op zoologisch, botanisch, oceanographisch en geologisch gebied verzameld in Nederlandsch Oost-Indie 1899-1900, Monographie 49a, Brill, Leiden, p. 1-107.
(online at: https://ia600405.us.archive.org/32/items/prosobranchiaofs13sche/prosobranchiaofs13sche.pdf)
('The Prosobranchs of the Siboga Expedition'. First of series of six monographs published in 1908-1913 on Recent marine gastropods from Indonesia, collected during Siboga Expediton 1899-1900)

Schepman, M.M. (1909)- The Prosobranchia of the Siboga Expedition, Part II. Taenioglossa and Ptenoglossa. In: M. Weber (ed.) Siboga Expeditie, Uitkomsten op zoologisch, botanisch, oceanographisch en geologisch gebied verzameld in Nederlandsch Oost-Indie 1899-1900, Monographie 49b, Brill, Leiden, p. 108-231.
('The Prosobranchs of the Siboga Expedition, Part II, Taenioglossa and Ptenoglossa'. Part 2 of series of six monographs on Recent marine gastropods from Indonesia, collected during Siboga Expediton 1899-1900)

Schepman, M.M. (1909)- The Prosobranchia of the Siboga Expedition, Part III. Gymnoglossa. In: M. Weber (ed.) Siboga Expeditie, Uitkomsten op zoologisch, botanisch, oceanographisch en geologisch gebied verzameld in Nederlandsch Oost-Indie 1899-1900, Monographie 49c, Brill, Leiden, p. 233-246.
('The Prosobranchs of the Siboga Expedition, Part III, Gymnoglossa'. 32 species. Part 3 of series of six monographs on Recent marine gastropods from Indonesia, collected during Siboga Expediton 1899-1900)

Schepman, M.M. (1911)- The Prosobranchia of the Siboga Expedition, Part IV. Rachiglossa. In: M. Weber (ed.) Siboga Expeditie, Uitkomsten op zoologisch, botanisch, oceanographisch en geologisch gebied verzameld in Nederlandsch Oost-Indie 1899-1900, Monographie 49d, Brill, Leiden, p. 247-364.
(online at: http://ia600409.us.archive.org/26/items/sibogaexpeditie58sibo/sibogaexpeditie58sibo.pdf)
('The Prosobranchs of the Siboga Expedition, Part IV, Rachiglossa'. Part 4 of series of six monographs on Recent marine gastropods from Indonesia, collected during Siboga Expediton 1899-1900)

Schepman, M.M. (1911)- The Prosobranchia of the Siboga Expedition, Part V. Toxoglossa. In: M. Weber (ed.) Siboga Expeditie, Uitkomsten op zoologisch, botanisch, oceanographisch en geologisch gebied verzameld in Nederlandsch Oost-Indie 1899-1900, Monographie 49e, Brill, Leiden, p. 365-452.
('The Prosobranchs of the Siboga Expedition, Part V, Toxoglossa'. Part 5 of series of six monographs on Recent marine gastropods from Indonesia, collected during Siboga Expediton 1899-1900)

Schepman, M.M. (1913)- The Prosobranchia of the Siboga Expedition, Part VI. Pulmonata and Opisthobranchia Tectibranchiata, tribe Bullomorpha. In: M. Weber (ed.) Siboga Expeditie, Uitkomsten op zoologisch, botanisch, oceanographisch en geologisch gebied verzameld in Nederlandsch Oost-Indie 1899-1900, Monographie 49f, Brill, Leiden, p. 453-494.
('The Prosobranchs of the Siboga Expedition, Part VI, Pulmonata and Opisthobranchia, Tectibranchiata, tribe Bullomorpha'. Part 6 of series of six monographs on Recent gastropods from Indonesia, collected during Siboga Expediton. 19 species of land- and freshwater molluscs (2 new), 41 species of Opisthobranchia Tectibranchiata Bullomorpha with 15 new sp.)

Schmidt, C., P. Heinz, M. Kucera & S. Uthicke (2011)- Temperature-induced stress leads to bleaching in larger benthic foraminifera hosting endosymbiotic diatoms. Limnology and Oceanography 56, 5, p. 1587-1602.
(online at: www.aslo.org/lo/toc/vol_56/issue_5/1587.pdf)
(Aquarium experiments on living larger forams Amphistegina radiata and Heterostegina depressa, collected from Great Barrier Reef. Function normally at temperatures of 23-28°C, but show bleaching and lack of growth at T of 31°C and higher)

Schonfield, J. (1994)- Biostratigraphy and assemblage composition of benthic foraminifera from the Manihiki Plateau, southwestern tropical Pacific. J. Micropalaeontology 14, 1, p. 165-175.
(online at: https://www.j-micropalaeontol.net/14/165/1995/jm-14-165-1995.pdf)
(Deep water late Pliocene- Pleistocene benthic foraminifera from Sonne cruise SO67 core on Manihiki Plateau in SW Tropical Pacific (2612m water depth). Dominated by Nodogenerina, Cibicidoides wuellerstorfi,

Oridorsalis umbonatus, Pleurostomella, Dentalina, etc. Remarkable absence of 'high-productivity taxa' Bolivina, Bulimina, Chilostomella and Uvigerina, suggesting low flux of organic matter to sea floor)

Schubert, R.J. (1900)- Uber die recente Foraminiferenfauna von Singapore. Zoologischer Anzeiger 23, p. 500-502.
('On the Recent foraminifera fauna of Singapore'. Foram faunas in shallow marine calcareous sand off Singapore dominated by miliolids)

Schudack, M.E. & J. Reitner (1996)- Holocene Ostracoda from the Satonda crater lake (Indonesia). Gottinger Arbeiten Geologie Palaeontologie, Sonderband SB2, p. 119-123.
(Holocene Ostracoda from 80cm deep digs on beach of Satonda Crater Lake off Sumbawa: (1) lowermost samples monospecific associations of fresh-brackish Aglaiocypris; (2) middle part higher diversity including marine Tenedocythere, etc.; (3) in upper parts of digs and in alkaline waters of today's crater lake, same cypridid species as in lowermost horizons, reflecting re-establishment of more stressful environment)

Severin, K.P. (1983)- The size-frequency distribution of the foraminifer *Marginopora vertebralis* on seagrass through time. Science in New Guinea 10, p. 187-195.

Shafik, S. (1978)- The near-surface sediments of the Scott Plateau and Java Trench: nannofossil assessment and implications. BMR J. Australian Geology Geophysics 3, p. 341-345.
(online at: www.ga.gov.au/corporate_data/80975/Jou1978_v3_n4_p341.pdf)
(Quaternary sediments in N Scott Plateau and Java Trench. Quaternary calcareous nannofossils associated with reworked U Cretaceous and Tertiary forms, possibly caused by bottom currents eroding parts of C Scott Plateau. Late Pleistocene-Holocene calcareous nannofossils in upper ~1 m thick of N Scott Plateau, but absent from Java Trench suggesting that present Nanno Solution Depth is between 3290-4950m water depth)

Shuto, T. (1970)- Taxonomical notes on the Turrids of the Siboga-Collection originally described by M.M. Schepman, 1913 (Part 1). Venus (Japanese J. Malacology) 28, 4, p. 161-178.

Shuto, T. (1970)- Taxonomical notes on the Turrids of the Siboga-Collection originally described by M.M. Schepman, 1913 (Part 2). Venus (Japanese J. Malacology) 29, 2, p. 37-54.

Shuto, T. (1971)- Taxonomical notes on the Turrids of the Siboga-Collection originally described by M.M. Schepman, 1913 (Part 3). Venus (Japanese J. Malacology) 30, 1, p. 5-22.

Sidiq, A., S. Hadisusanto & K.T. Dewi (2016)- Foraminifera bentonik kaitannya dengan kualitas perairan de wilayah barat daya Pulau Morotai, Maluku Utara. J. Geologi Kelautan 14, 1, p. 13-22.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/336/264)
('Benthic foraminifera and relation to water quality off SW Morotai Island, North Moluccas'. Seafloor samples from 16-36m depth off Morotai, N Moluccas, with 28 species of benthic foraminifera, dominated by abundant Amphistegina spp. and Operculina spp. Also Alveolinella, Calcarina, Heterolepa, Baculogypsinoides, Elphidium, Peneroplis, Schlumbergerella and Sorites)

Sijinkumar, A.V., B.N. Nath, G. Possnert & A. Aldahan (2011)- *Pulleniatina* minimum events in the Andaman Sea (NE Indian Ocean): implications for winter monsoon and thermocline changes. Marine Micropaleontology 81, p. 88-94.
(Late Quaternary record of Pulleniatina obliquiloculata in cores from Andaman Sea. As in Pacific Ocean, Pulleniatina obliquiloculata Minimum Event exists in Indian Ocean between 4.5- 3.0 ka. Two additional minimum events in Younger Dryas and late Last Glacial Maximum (20-18 ka). PMEs of Andaman Sea characterized by fewer thermocline species, indicating increased depth of thermocline during minimum events)

Silalahi, I.R., M.K. Adisaputra, R. Kapid & M. Hendrizan (2012)- Album mikrofosil foraminifera dan nanoplankton perairan Indonesia. Puslitbang Geologi Kelautan RI (Research Center of Marine Geology), Bandung, p. 1-143.

('Album of foraminifera and nannoplankton microfossils from Indonesian waters')

Siregar, M.S. (1978)- Pengaruh musim terhadap penyebaran foraminifera di Teluk Jakarta. J. RISET Geologi dan Pertambangan (LIPI) 1, 3, p. 33-78.
('The influence of the seasons on the distribution of foraminifera in Jakarta Bay'. Seasonal influence shown by modern foraminifera Rotalia and Quinqueloculina)

Smith, B.J. & M. Djajasasmita (1988)- The land molluscs of the Krakatau Islands, Indonesia. Philosophical Transactions Royal Society London, Ser. B, 322, 1211, p. 379-400.
(Land molluscs fauna from 1984-85 Zoological Expeditions to Krakatau Islands 19 species: 16 from Rakata, 12 from Sertung, 7 from Panjang and 1 from Anak Krakatau (first record of land snail from that island))

Soemodihardjo, S. & A. Matsukuma (1989)- Ecology of sandy beach bivalves of Pari Island off the coast of Jakarta Bay. Indonesia. Bull. National Science Museum, Tokyo, Ser. A, 15, p. 197-212.
(online at: http://ci.nii.ac.jp/lognavi?name=nels&lang=en&type=pdf&id=ART0006479304)

Somboon, J.R.P. (1990)- Palynological study of mangrove and marine sediments of the Gulf of Thailand. J. Southeast Asian Earth Sciences 4, 2, p. 85-97.

Southward, E.C., A. Schulze & V. Tunnicliffe (2002)- Vestimentiferans (Pogonophora) in the Pacific and Indian Oceans: a new genus from Lihir Island (Papua New Guinea) and the Java Trench, with the first report of *Arcovestia ivanovi* from the North Fiji Basin. J. Natural History 36, p. 1179-1197.
(Example of occurrences of tube worms at cold gas seeps and hot hydrothermal vent sites in SW Pacific and S Java deepwater seafloor settings (also known from Sumatra forearc, etc.))

Staub, W. (1915)- Uber die Verbreitung einiger lebender und versteinter Lamellibranchier und Gastropodenarten am Ausgange der Sangkulirangbai (Ost Borneo), einem Aestuarium der tropischen Zone. Vierteljahrschrift Naturforschenden Gesellschaft Zurich 61, p. 120-135.
('On the distribution of some living and fossilized bivalve and gastropod species at the mouth of Sangkulirang Bay (East Borneo), an estuary of the tropical sea'. Early paper on distribution of land, brackish and marine molluscs in estuary of NE Kalimantan)

Stelbrink, B. (2014)- A biogeographic view on Southeast Asia's history. Doct. Thesis Humboldt-Universitat, Berlin, p. 1-270.
(online at: http://edoc.hu-berlin.de/dissertationen/stelbrink-bjoern-2014-12-19/PDF/stelbrink.pdf)
(On significance of modern fauna and floral distribution for evolution of SE Asia, with focus on Sulawesi)

Sterrenburg, F.A.S, P.L.A. Erftemeijer & P.H. Nienhuis (1995)- Diatoms as epiphytes on seagrasses in South Sulawesi (Indonesia); comparison with growth on inert substrata. Botanica Marina 38, p. 1-8.

Stidolph, S.R., F.A.S. Sterrenburg, K.E.L. Smith & A. Kraberg (2012)- Stuart R. Stidolph diatom atlas. U.S. Geological Survey (USGS) Open File Report 2012-1163.
(online at: http://pubs.usgs.gov/of/2012/1163/)
(Spectacular photographs of modern coastal marine diatoms, including Indonesian material on Plates 30-33 (Semarang, Sumatra))

Stuijts, I. (1984)- Palynological study of Situ Bayongbong, West Java. Modern Quaternary Research in Southeast Asia 8, Balkema, Rotterdam, p. 2-17.
(Palynology of 8m sediment core from Bayombong swamp at ~1200m above s.l. in SW Java, showing Late Pleistocene- Holocene (~17 ky- today) vegetational shift from upper montane forest to lower montane forest around ~10,000 yrs B.P.)

Stuijts, I. (1993)- Late Pleistocene and Holocene vegetation of West Java, Indonesia. Modern Quaternary Research in Southeast Asia 12, Balkema, Rotterdam, p. 1-173. *(also Thesis Rijksuniversiteit Groningen)*

(Late Pleistocene- Holocene palynology of shallow core holes from montane localities above 1000m in West Java (Gede-Panggrango, Gunung Patuha, Danau Ciharus))

Stuijts, J.C., J.C. Newsome & J.R. Flendley (1988)- Evidence for Late Quaternary vegetational change in the Sumatran and Javan highlands. Review Palaeobotany Palynology 55, p. 207-216.
(Late Pleistocene pollen records from above 2000m on tropical mountains indicate cooler climates and more arid climates below 1200m. Sumatran and Javan sites at intermediate altitudes show higher altitude vegetation from ~18,200 yr B.P. to ca. 12,400 yr B.P., suggesting much lower forest altitudinal boundaries than today's)

Sugawara, D., K. Minoura, N. Nemoto, S. Tsukawaki, K. Goto & F. Imamur (2009)- Foraminiferal evidence of submarine sediment transport and deposition by backwash during the 2004 Indian Ocean tsunami. Island Arc 18, 3, p. 513-525.
(online at: http://onlinelibrary.wiley.com/doi/10.1111/j.1440-1738.2009.00677.x/epdf)
(Nearshore to offshore sediments from SW coast of Thailand clarify submarine sediment transport during 2004 Indian Ocean tsunami. Benthic foraminifera showed seaward migration after tsunami event (brackish agglutinated foraminifera in post-tsunami foreshore to offshore, transported offshore with tsunami backwash). Offshore planktonic and benthic species slight evidence of landward migration by tsunami)

Suhartati, M.N. (1992)- Preliminary study on the benthic foraminifera and its association with ostracoda in Porong Delta, East Java. Toyama University, p. 1-10.
(15 sediment samples from 0.6- 21.5m along Porong delta front, Madura Straits. Most abundant species Ammonia beccarii, Calcarina calcar and Elphidium advenum. Also common Elphidium crispum, Asterorotalia trispinosa, Pseudorotalia schroeteriana and Quinqueloculina)

Suhartati, M.N. (1994)- Foraminifera bentonik dan kaitannya dengan kandungan zat hara di perairan Padang Lamun, Goba Besar, Pulau Pari, Kepulauan Seribu. Proc. 23rd Annual Conv. Indonesian Association Geologists (IAGI), Jakarta, 1, p. 582-590.
(online at: https://www.iagi.or.id/web/digital/62/49.pdf)
('Foraminifera bentonic and their relation to nutrient content in the waters of Padang Lamun, Goba Besar, Pulau Pari, Thousand Islands'. Diverse benthic foram assemblage on seagrass beds of Pari Island, dominated by Quinqueloculina spp., Spiroloculina spp., Ammonia, Peneroplis, Rosalina, Elphidium, Calcarina, etc.)

Suhartati, M.N. (1994)- Benthic foraminifera in the seagrass beds of Pari island- Seribu islands, Jakarta. In: S. Sudara et al. (eds.) Proc. Third ASEAN-Australia Symposium Living Coastal Resources, Bangkok, 2, p. 323-329.
(Dominant species in reef-flat seagrass areas are miliolids Quinqueloculina spp., and Spiroloculina spp. Also common Ammonia beccarii, Calcarina calcar, Elphidium spp.)

Suhartati, M. Natsir (1998)- First record of brackish water agglutinated foraminifera from Java. Reopical Biodiversity 5, 1, p. 57-63.
(Ammobaculites agglutinans and Textularia pseudogramen common in Recent sediments near Solo and Poreng River mouths, E Java)

Suhartati, M. Natsir (2005)- Distribusi foraminifera bentik (*Textularia*) di Delta Porong, Jawa Timur. Agritek 4, 2, p. 1-7.
('Distribution of benthic foraminifera (Textularia) in the Porong Delta, East Java')

Suhartati, M. Natsir (2009)- Distribusi dan kelimpahan foraminifera bentik Resen di Pulau Opak Besar, Kepulauan Seribu. Lingkungan Tropis 3, 2, p. 95-103.
(Online at: www.lingkungan-tropis.org/distribusi-dan-kelimpahan-foraminifera-suhartati-m-natsir)
('Distribution and abundance of Recent benthic foraminifera in Opak Besar, Seribu islands'. Off NW Java. Samples from water depths 27-36m around Opak Island dominated by Calcarina calcar, followed by miliolids)

Suhartati, M.N. (2010)- First record of agglutinated foraminifera from Lombok. J. Coastal Development 13, 1, p. 48-55.
(online at: www.omicsonline.com/open-access/first-record-of-agglutinated-foraminifera-from-lombok-1410-5217-13-276.pdf)
(Benthic foram assemblages around Gili islands, NW Lombok, have more agglutinated individuals in stations close to bay, mainly Ammobaculites agglutitans and Haplophragmoides canariensis)

Suhartati M. Natsir (2010)- The distribution of benthic foraminifera in Damar and Jukung Island, Seribu Islands. Marine Research in Indonesia (LIPI) 35, 2, p. 9-14.
(online at: https://www.researchgate.net/publication/327826272_THE_DISTRIBUTION_OF_BENTHIC_FORAMINIFERA_IN_DAMAR_ISLAND_AND_JUKUNG_ISLAND_SERIBU_ISLANDS)
(Benthic foraminifers on Jakarta Bay islands. Jukung Island higher diversity than Damar Besar Island. Larger foraminifera of both islands Amphistegina, Calcarina, Heterostegina, Marginophora, and Operculina)

Suhartati M. Natsir (2010)- Foraminifera bentik sebagai indikator kondisi lingkungan terumbu karang perairan Pulau Kotok Besar dan Pulau Nirwana, Kepulauan Seribu. Oseanologi dan Limnologi di Indonesia 36, 2, p. 181-192.
('Benthic foraminifera as indicator of environmental conditions of coral reefs in Kotok Besar and Nirwana islands of Seribu islands'. Kotok Besar Island healthy reef growth due to FORAM Index of ~7.6. Dominant symbiont bearing foraminifera are Amphistegina, Calcarina and Tinoporus. Nirwana Island was dominated by opportunistic foraminifera Ammonia, Elphidium, Quinqueloculina and Spiroloculina, showing stressed conditions unsuitable for reef growth as shown by FORAM Index of 1.6-1.9)

Suhartati M. Natsir (2010)- Kelimpahan foraminifera Resen pada sedimen permukaan di Teluk Ambon. J. Ilmu Teknologi Kelautan Tropis (JITKT; Bogor) 2, 1, p. 9-18.
(online at: http://repository.ipb.ac.id/jspui/bitstream/123456789/53406/1/2_foraminifera.pdf)
('The abundance of recent foraminifera in surface sediment of Ambon Bay'. Bottom samples from Ambon Bay with 61 species of benthic and 25 species of planktonic foraminifera. Dominant benthics Amphistegina lessonii, Ammonia beccarii, Elphidium craticulatum, Operculina ammonoides and Quinqueloculina. Forams generally abundant on sand substrate sand, but no foraminifera on mud substrate)

Suhartati M. Natsir (2012)- The benthic foraminiferal assemblages on Handeuleum Islands, Ujung Kulon National Park of Banten, Indonesia. J. Shipping and Ocean Engineering 2, p. 86-91.
(online at: www.davidpublishing.com/...)
(Nine sediment samples from around Handeuleum Islands off Ujung Kulon Peninsula contain 14 genera of benthic foraminifera and some Ostracoda and Bryozoa. Most specimens from sand sediments of coral reefs community. Most common foraminifera are opportunistic taxa such as Ammonia beccari and Elphidium craticulatum and E. crispum. Also present are symbiont bearing foraminifera Amphistegina, Calcarina, Sorites also Cymbaloporetta, Oolina, Quinqueloculina and Spiroloculina)

Suhartati M. Natsir (2014)- The distribution of benthic foraminifera in Indonesian shallow waters. Berita Sedimentologi 29, p. 66-72.
(online at: https://journal.iagi.or.id/index.php/FOSI/article/view/143)
(Summary of studies on Recent benthic foraminifera in Indonesia shallow waters by Research Center for Oceanography- LIPI)

Suhartati M. Natsir & K.T. Dewi (2015)- Foraminifera bentik terkait dengan kondisi lingkungan perairan sekitar Pulau Damar, Kepulauan Seribu. J. Geologi Kelautan 13, 3, p. 165-171.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/271/261)
('Benthic foraminifera related to marine environments around Damar Island, Pulau Seribu'. 64 species of benthic foraminifera from 11-37m depth around Damar Island reef, S part of Thousand Islands. Common forms associated with coral reef incl. Amphistegina lessonii, A. radiata, Sorites marginalis, Heterostegina and Calcarina calcar)

Suhartati M. Natsir, K.T. Dewi & S. Ardhyastuti (2017)- Keterkaitan foraminifera dan kedalaman perairan sebelah tenggara Pulau Seram, Maluku. J. Geologi Kelautan 15, 2, p. 73-80.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/389/424)
('The relation between foraminifera and water depth of waters SE of Seram Island, Moluccas'. Nine samples between 512-1177m water depth, with 95-100% planktonic foraminifera. Rare benthic foraminifera incl. Bulimina, Buliminella, Bolivinella)

Suhartati M. Natsir, A. Firman, I. Riyantini & I. Nurruhwati (2015)- Struktur komunitas foraminifera pada sedimen permukaan dan korelasinya terhadap kondisi lingkungan lepas pantai Balikpapan, Selat Makassar. J. Ilmu Teknologi Kelautan Tropis (JITKT; Bogor) 7, 2, p. 671-680.
(online at: http://journal.ipb.ac.id/index.php/jurnalikt/article/view/11059/8774)
('Community structure of foraminifera in surface sediments and correlation with environmental conditions in offshore waters of Balikpapan, Makassar Strait'. Foraminifera from 6 seafloor samples off Balikpapan (no water depths for samples (54-73m?), no species identifications))

Suhartati M. Natsir & Z.A. Muchlisin (2012)- Benthic foraminiferal assemblages in Tambelan Archipelago, Indonesia. AACL Bioflux 5, 4, p. 259-264.
(online at: https://bioflux.com.ro/docs/2012.259-264.pdf)
(Recent foraminifera in shallow waters around Tambelan islands, S China Sea. Dominated by Amphistegina and Assilina ammonoides; also common Quinqueloculina, Pseudorotalia, Amphistegina, Elphidium)

Suhartati, M.N., Ricky R. & Helfinalis (1994)- Foraminifera bentonik dan spesifikasinya pada beberapa lingkungan perairan Dangkal di Indonesia. Proc. 23rd Annual Conv. Indonesian Association Geologists (IAGI), Jakarta, 1, p. 603-604.

Suhartati M. Natsir & Rubiman (2010)- Distribusi foraminifera Resen di Laut Arafura. J. Ilmu Teknologi Kelautan Tropis (JITKT; Bogor) 2, 2, p. 74-82.
(online at: www.itk.fpik.ipb.ac.id/ej_itkt22/jurnal/ML_185_final%20(74-82).pdf)
('The distribution of Recent benthic foraminifera in the Arafura Sea'. Arafura Sea shallow waters S of Papua. Shallow-water ecosytems such as mangrove, seagrass beds and coral reefs. Samples mainly between 30-90m, deepest station 13 at 341m. 37 species, most common Ammonia beccarii and Pseudorotalia schroeteriana, except station 13 which has abundant Bolivina spp. and Anomalina rostrata)

Suhartati & Subadri (1993)- Foraminifera bentonik di perairan menpawah dan Sungai Duri- Pontianak Kalimantan Barat. Proc. 22nd Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 2, p. 1181-1188.
(online at: https://www.iagi.or.id/web/digital/53/22nd-Volume-2-(6-9-Des-1993)-579-586.pdf)
('Benthic foraminifera in the waters offhore Sungai Duri, Pontianak, W Kalimantan'. Shallow marine (0.5-5m) seafloor samples off Mempawah and Sunai Duri with 24 species of benthic foraminifera. Dominated by Quinquelocuina spp., Asterorotalia trispinosa, Ammonia beccarii, Pseudorotalia schroeteriana, etc.))

Suhartati M. Natsir & M. Subkhan (2012)- The distribution of agglutinated foraminifera in Porong and Solo deltas, East Java. J. Environmental Science and Engineering A 1, p. 918-923.
(Recent foraminiferal assemblages around Porong and Solo River Deltas dominated by small agglutinated forams, mainly Textularia pseudogramen, Ammobaculites agglutinans, Haplophragmoides, Ammotium, etc.)

Suhartati M. Natsir & M. Subkhan (2012)- The distribution of benthic foraminifera in coral reefs community and seagrass bed of Belitung Islands based on foram index. J. Coastal Development 15, 1, p. 51-58.
(online at: http://ejournal.undip.ac.id/index.php/coastdev/article/view/1997/1775)
(Benthic foraminiferal from six sampling sites around Belitung Islands 29 species of 18 genera. Most abundant benthic foraminifera in Nasik Strait on coarse sand substrate with coral reef (Peneroplis, Calcarina, Operculina, etc.). Seagrass beds of Nasik Strait dominated by opportunistic foraminifera Heterostegina, Calcarina, Elphidium, Ammonia, Acervulina, Spirolina, Quinqueloculina and Lenticulina. Most abundant species of all sites is Peneroplis pertusus)

Suhartati M. Natsir & M. Subkhan (2012)- Foraminifera bentik sebagai indikator kualitas perairan ekosistem terumbu karang di Pulau Bidadari dan Ringit, Kepulauan Seribu. Lingkungan Tropis 5, 1, p. 1-10.
('Benthic foraminifera as indicators for water quality of coral reefs ecosystem in Bidadari and Ringit Islands, Thousand Islands', off NW Java.)

Suhartati M. Natsir, M. Subkhan, Rubiman &, S.P.A. Wibowo (2011)- Komunitas foraminifera bentik di perairan kepulauan Natuna. J. Ilmu Teknologi Kelautan Tropis (JITKT; Bogor) 3, 2, p. 21-31.
('Benthic foraminifera community in the Natuna islands group'. Over 50 species, dominated by Amphstegina lessonii, Ammonia beccarii, Operculina ammonoides, Quinqueloculina, etc. Water depths??)

Suhartati M. Natsir, M. Subkhan, M.S. Tarigan, S.P.A. Wibowo & K.T. Dewi (2012)- Benthic foraminifera in South Waigeo waters, Raja Ampat, West Papua. Bulletin of the Marine Geology 27, 1, p. 1-6.
(online at: http://ejournal.mgi.esdm.go.id/index.php/bomg/article/view/40/41)
(Foraminifera from 12 sites in S part off Waigeo Island, Raja Ampat group, W Papua. Faunas dominated by symbiotic bearing benthic foraminifera, mainly Amphistegina lessonii, also Baculogypsina, Calcarina, Tinoporus, Heterostegina, etc.)

Suhartati M. Natsir, M. Subkhan & V.I. Wardhani (2012)- The distribution of benthic foraminiferal assemblages in Tambelan Islands of Riau Islands. Proc. Soc. Indonesian Biodiversity International Conference, 1, p. 107-112.
(online at: https://biosains.mipa.uns.ac.id/P/P0101/P010119.pdf)
(Foraminifera from 5 shallow shelf stations around Tambelan Archipelago in South China Sea, off NW Kalimantan (34-50m water depth). Sediments mainly clay and mud. Assemblages 64 species, dominated by Operculina ammonoides ('Assilina depressa'), Amphistegina lessonii and miliolids)

Sukandarrumidi (1990)- The new species of *"Quinqueloculina"* and "*Triloculina*" from the bottom sea sediment of the Java Sea, Bali Strait and Karimata Strait, Indonesia. Media Teknik (UGM) 13, 2, p. 132-143.
(Sea floor samples from Java Sea, Bali Strait and Karimata Strait with 3 new species of miliolid benthonic foraminifera, Quinqueloculina aberensis, Triloculina malayensis and Triloculina siuriensis)

Sukapti, W.S. & K.T. Dewi (2023)- Perubahan lingkungan Holosen akhir berdasarkan rekaman polen dari Delta Berau-Kalimantan Timur. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 24, 1, p. 23-29.
(online at: https://jgsm.geologi.esdm.go.id/index.php/JGSM/article/view/713)
('Pollen study of Late Holocene environment changes in Berau Delta-East Kalimantan'. Pollen from 100 cm long core in Berau Delta, showing 80 types of pollen (dominated by lowland forest species) and 13 types of spores. Some fluctuations in % of mangrove- dry land- grassland and (sub-)montane pollen)

Suleiman, A, C.C.S. Wahyu & A. Bachtiar (2011)- Quaternary benthic foraminifera from bathyal zone seabed of Mamuju offshore, North Makassar Basin, West Sulawesi. Proc. Joint 36th HAGI and 40th IAGI Annual Conv., Makassar, JCM2011-424, p. 1-5.
(online at: https://www.iagi.or.id/web/digital/10/2011_IAGI_Makassar_Quarternary-Benthic-Foraminifera.pdf)
(Summary of benthic foraminifera from 1700-1800m depth, sampled during geohazard survey, offshore, Lariang Basin. Mainly calcareous benthics usually found in outer shelf- upper bathyal instead of arenaceous tests that are common in bathyal zone. This suggests deposition is at least partly allochthonous sediment debris from upslope (no data details))

Sumawinata, B. (1998)- Sediments of the lower Barito basin in South Kalimantan: fossil pollen composition. J. Southeast Asian Studies, Kyoto, 36, 3, p. 293-316.
(online at: https://kyoto-seas.org/pdf/36/3/360302.pdf)
(Palynology and successions of depositional environments inf Holocene sediments from 8 shallow (2-3m) cores in Lower Barito and Martapura River basins. Pollen diagrams dominated by mangrove forest taxa)

Sun, H.J., T.G. Li, R.T. Sun, X.K. Yu & F.M Chang & Z. Tang (2011)- Calcareous nannofossil bioevents and microtektite stratigraphy in the Western Philippine Sea during the Quaternary. Chinese Science Bull. 56, 25, p. 2732-2738.
(Seven calcareous nannofossil bioevents identified over past 2.36 Ma in two sediment cores from Benham Rise, W Philippine Sea, E of Luzon. Bioevents and Australasian microtektite impact event calibrated to oxygen isotope stratigraphy. Age of highest concentration of microtektites 792 ± 2 ka, near boundary of isotope stages MIS20/19)

Sun, X., Y. Luo, F. Huang, J. Tian & P. Wang (2003)- Deep-sea pollen from the South China Sea: Pleistocene indicators of East Asian monsoon. Marine Geology 201, p. 97-118.
(High-resolution pollen record for last 820 ka of ODP Site 1144, northern S China Sea. 29 pollen zones, mainly defined by alternations of Pinus-dominant (interglacial) vs. herb-dominant (glacial) zones correspond to Marine Oxygen Isotope Stages 1-29. Clear 100 ka Milankovich cyclicity)

Suriadi, R., H. Shaari, S.J. Culver, M.L. Husain, V.R. Vijayan, P.R. Parham, A. Sulaiman & N. Sapon (2019)- Inner shelf benthic foraminifera of the South China Sea, East Coast Peninsular Malaysia. J. Foraminiferal Research 49, 1, p. 11-28.
(online at: https://www.academia.edu/download/67521350/Suriadi_et_al._2019.pdf)
(Distribution of modern benthic foraminifera on inner shelf of southern S China Sea, off E coast of Malaysia, between Johor and Terengganu. Samples from <50 m water depth with 266 species. Two biofacies: (A) common Amphistegina papillosa in sandier sediments, (B) common Pseudorotalia schroeteriana and other small rotaliids in muddy sediments. Assemblages reflect strong fluvial/terrestrial influence on tropical shelf environment)

Suriadi, R., H. Shaari, S.J. Culver, A. Sulaiman, S. Yang, W.N.W. Saelan & M.M. Sharif (2023)- Holocene transgression and sea level change offshore Peninsular Malaysia (innermost Sunda Shelf) as indicated by benthic foraminiferal, lithologic and geochronologic data. Quaternary International 649, p. 46-57.
(open manuscript online at: https://www.sciencedirect.com/science/article/pii/S1040618223000034)
(Evidence of Holocene transgression from inner shelf of eastern Peninsular Malaysia, based on lithologic and foraminiferal analyses of 24 cores from 13.5- 46.8 m water depth)

Szarek, R. (2001)- Biodiversity and biogeography of Recent benthic foraminiferal assemblages in the south-western South China Sea (Sunda Shelf). Ph.D. Thesis, Christian Albrechts Universitat, Kiel, p. 1-273.
(online at: https://macau.uni-kiel.de/servlets/MCRFileNodeServlet/dissertation_derivate_00000537/d537.pdf)
(Benthic foraminifera distribution patterns on Vietnam Shelf and Sunda Shelf of SW S China Sea, based on 75 sites along two transects in 50-2000m water depth. Shallow water (<200m) assemblages from Vietnam and Sunda Shelves significantly different species composition and distinct distribution patterns. Bathyal faunas exhibit more uniform species composition)

Szarek, R., W. Kuhnt, H. Kawamura & H. Kitazato (2006)- Distribution of Recent benthic foraminifera on the Sunda Shelf (South China Sea). Marine Micropaleontology 61, p. 171-195.
(Recent benthic foraminifera distribution on Sunda Shelf around Natuna Island between 60-226m depth. Four biofacies: (A) inner shelf (Ammomassilina alveoliniformis- Asterorotalia pulchella), in fine grained sediments; (B) high-energy inner shelf (Heterolepa dutemplei- Textularia lythostrota, Asterorotalia gaimardii) in sand and silt dominated sediments NE of Natuna; (C) high-energy outer shelf biofacies (Cibicidoides pachyderma-Textularia bocki, Operculina ammonoides) in neritic relict sand; (4) outer shelf (Facetocochlea pulchra-Bulimina marginata, Bolivina) in area covered with modern silt and mud)

Szarek, R., W. Kuhnt, H. Kawamura & H. Nishi (2009)- Distribution of Recent benthic foraminifera along continental slope of the Sunda Shelf (South China Sea). Marine Micropaleontology 71, 1-2, p. 41-59.
(Benthic foraminiferal distribution from the winter upwelling region off Borneo on continental slope of Sunda Shelf and from continental slope of S Vietnam Shelf. Faunas highly diverse. Four biofacies: (1) Upper bathyal (Siphotextularia foliosa- Cibicidoides robertsonianus); (2) Middle bathyal (Uvigerina auberiana- Nuttallides

rugosus; within oxygen minimum zone); (3) uppermost Lower bathyal (Lagenammina difflugiformis- Uvigerina peregrina) and (4) Lower bathyal (Paratrochammina challengeri- Parrelloides bradyi))

Tan Koh Siang (2021)- Deep-sea Protobranchia (Bivalvia) from the SJADES biodiversity cruise, with a description of a new species of *Propeleda* (Nuculanidae). Raffles Bull. of Zoology, Supplement. 36, p. 346-385.
(online at: https://www.science.nus.edu.sg/wp-content/uploads/sites/11/2024/07/s36rbz346-385.pdf)
(Study of Recent deep marine (100-2000m) bivalve molluscs collected during South Java Deep-Sea (SJADES) Biodiversity Expedition 2018. 37 stations off SE Sumatra and SW Java in Sunda Strait and Indian Ocean. 37 species in 17 genera belonging to families Nuculidae (incl. Nucula), Solemyidae (incl. Acharax), Nucinellidae, Nuculanidae, Malletiidae and Neilonellidae). Several evidences of methane venting on seafloor)

Tanaka, G., T. Komatsu & N.D. Phong (2009)- Recent ostracod assemblages from the northeastern coast of Vietnam and biogeographical significance of euryhaline species. Micropaleontology 55, p. 365-382.
(75 species, 3 biofacies controlled by salinity)

Taylor, A.M. (1988)- The taxonomy, ecology and zoogeographical significance of Recent reef Ostracoda from Singapore. M.Sc. Thesis, University of Wales, Aberystwyth, p. 1-203. *(Unpublished)*

Taylor, P. D. & S.H.A.Tan (2015)- Cheilostome bryozoa from Penang and Langkawi, Malaysia. European Journal of Taxonomy 149, p. 1-34.
(online at: https://europeanjournaloftaxonomy.eu/index.php/ejt/article/view/271/522)
(Twenty-three species of Recent cheilostome bryozoans from beaches of Penang and Langkawi islands)

Thanikaimoni, G. (1983)- Palynological investigation on the Borobudur monument. Bull. Ecole française d'Extreme-Orient 72, p. 237-250.
(online at: www.persee.fr/docAsPDF/befeo_0336-1519_1983_num_72_1_1458.pdf)
(Palynomorphs from soil material used for construction of base of Borobudur monument. Collected from alluvial deposits in open area, not covered by dense vegetation. Absence of marsh and aquatic elements like Typha and Nymphaea in samples suggests not derived from lake or marsh)

Thomas, E.A., T. Bond, J.L. Kolbusz, Y. Niyazi, D.J.B. Swanborn & A.J. Jamieson (2024)- Deep-sea ecosystems of the Indian Ocean >1000m. Science of the Total Environment 957, 176794, p. 1-35)
(https://www.sciencedirect.com/science/article/pii/S0048969724069511
(Comprehensive review of present-day deep-sea Indian Ocean physical features and associated ecosystems)

Thomas, M.L. (2015)- Holocene palynology of the Gulf of Papua, Papua New Guinea: using modern palynomorph distribution to better constrain paleoenvironmental changes. Ph.D. Thesis, Louisiana State University, Baton Rouge, p. 1-207.
(online at: http://digitalcommons.lsu.edu/cgi/viewcontent.cgi?article=1786&context=gradschool_dissertations)

Thomas, M.L., D.T. Pocknall, S. Warny, S.J. Bentley, A.W. Droxler & C.A Nittrouer (2015)- Assessing palaeobathymetry and sedimentation rates using palynomaceral analysis: a study of modern sediments from the Gulf of Papua, offshore Papua New Guinea. Palynology 39, 3, p. 410-433.
(Palynofacies analyses of organic matter of 64 seafloor samples in Gulf of Papua, PNG, from river mouth to shelf to slope. Five categories of palynomacerals distinguished: (1, 2) brown wood, (3) leaf cuticle, (4) black debris, (5) structureless organic matter. Palynomacerals 1-3 more common in nearshore facies, 4 more common in deeper offshore sites)

Thomas, M.L., S. Warny, D.M. Jarzen, S.J. Bentley, A.W. Droxler, B.B. Harper, C.A. Nittrouer & X. Xu (2018)- Palynomorph evidence for tropical climate stability in the Gulf of Papua, Papua New Guinea, over the latest marine transgression and highstand (14,500 years BP to today). Quaternary International 467, B22, p. 277-291.
(online at: https://sites01.lsu.edu/faculty/swarny/wp-content/uploads/sites/30/2018/02/Thomas-et-al.-2018-QI.pdf)

(Palynological data indicate climatic conditions at sea level around Gulf of Papua remained warm, wet and stable for past 14.5 kyr, with sea surface T >14 °C)

Titterton, R. & R.C. Whatley (1988)- The provincial distribution of shallow water Indo-Pacific and marine ostracoda: origins, antiquity, dispersal routes and mechanisms. In: T. Hanai et al. (eds.) Evolutionary biology of ostracoda: its fundamentals and applications, Proc. 9th International Symposium on Ostracoda (ISO), Shizuoka 1985, Elsevier, Amsterdam, p. 759-786.
(Tertiary to Recent ostracods from Indo-Pacific and Southern Ocean fall into 13 zoogeographical provinces. East Indian and SW Pacific regions were locus from which ostracods migrated out since Miocene)

Titterton, R. & R.C. Whatley (1988)- Recent Bairdiinae (Crustacea, Ostracoda) from the Solomon Islands. J. Micropalaeontology 7, 2, p. 111-142.
(online at: https://www.j-micropalaeontol.net/7/111/1988/jm-7-111-1988.pdf)
(21 species of Bairdiinae ostracods (13 new) from littoral and inner shelf of Solomon Islands)

Titterton, R. & R.C. Whatley (2005)- Recent marine Ostracoda from the Solomon Islands. Part 2. Cytheracea (Xestoleberidae). Revista Espanola Micropaleontologia 37, 2, 291-313.
(online at: http://revistas.igme.es/index.php/revista_micro/article/viewFile/314/312)

Titterton, R. & R.C. Whatley (2006)- Recent marine Ostracoda from the Solomon Islands. Part 1: Cypridoidea, Platycopina and Cladocopina. J. Micropalaeontology 25, p. 73-94.
(online at: https://www.j-micropalaeontol.net/25/73/2006/jm-25-73-2006.pdf)
(16 species of Cypridoidea described, comprising 15% of the ostracode fauna of 160 species from Solomon Islands. Platycopids/ Cladocopids 13% of total fauna)

Titterton, R. & R.C. Whatley (2006)- Recent marine Ostracoda from the Solomon Islands. Part 3: Cytheroidea, Bythcytheroidea, Cytherideidae, Krithidae, Neoytherideidae, Cytheruridae. Revista Espanola Micropaleontologia 38, p. 169-189.
(online at: http://revistas.igme.es/index.php/revista_micro/article/view/325/322)

Titterton, R. & R.C. Whatley (2008)- Recent marine Ostracoda from the Solomon Islands. Part 4: Cytheroidea; Hemicytheridae, Thaerocytheridae. J. Micropalaeontology 27, p. 13-33.
(online at: https://www.j-micropalaeontol.net/27/13/2008/jm-27-13-2008.pdf)
(11 species of family Hemicytheridae and 7 of family Thaerocytheridae described, together comprising 14% of Recent ostracod fauna from Solomon Islands. Hemicytherids more endemic than thaerocytherids)

Titterton, R., R.C. Whatley & J.E. Whittaker (2001)- A review of some key species of mainly Indo-Pacific ostracoda from the collections of G.S. Brady. J. Micropalaeontology 20, 1, p. 31-44.
(online at: https://jm.copernicus.org/articles/20/31/2001/jm-20-31-2001.pdf)
(Review and re-naming of 15 Recent (mainly deep water?) ostracode species from G.S. Brady's Challenger collectionsin the Hancock Museum in Newcastle-upon-Tyne and the Natural History Museum, London)

Todd, R. (1960)- Some observations on the distribution of *Calcarina* and *Baculogypsina* in the Pacific. Science Reports Tohoku University, Sendai, ser. 2 (Geology), Special Volume 4, p. 100-107.
(online at: https://tohoku.repo.nii.ac.jp/record/10741/files/KJ00004163739.pdf)
(Calcarina spengleri and Baculogypsina sphaerulata common reef-dwelling foram species of tropical W Pacific between ~170°W and 120°E. Both first appeared late in Tertiary. Common from Australia Great Barrier Reef to Philippines and Ryukyus in W and from Marianas to Marshalls, Gilberts, Phoenix Islands, Samoa, and Niue in E (not known from C-E Pacific))

Todd, R. (1961)- Foraminifera from Onotoa Atoll Gilbert Islands. U.S. Geological Survey Professional Paper 354-H, p. 1-34.
(online at: https://pubs.usgs.gov/pp/0354h/report.pdf)

(Example of foraminifera distribution on a SW Pacific coral reef atoll. 168 species identified. Dominant species in reef sediments Calcarina spengleri, Baculogypsina sphaerulata, Amphistegina madagascariensis and Marginopora vertebralis. Main taxa in lagoon sediments Amphistegina madagascariensis, Heterostegina suborbicularis, Marginopora vertebralis and Elphidium striato-punctatum)

Todd, R. (1965)- The foraminifera of the tropical Pacific Collections of the "Albatross", 1894-1900, Part IV, Rotaliform families and planktonic families. Bull. U.S. National Museum, 161, p. 1-139.
(online at: https://repository.si.edu/handle/10088/10220)
(Descriptions of smaller rotaliform benthic and planktonic foraminifera from tropical Pacific Ocean (Part 4 of Cushman 1932-1942 monographs))

Toruan, L.N.L., D. Soedharma & K.T. Dewi (2013)- Komposisi dan distribusi foraminifera bentik di ekosistem terumbu karang pada Kepulauan Seribu. J. Ilmu Teknologi Kelautan Tropis (JITKT; Bogor) 5, 1, p. 1-16.
(online at: http://journal.ipb.ac.id/index.php/jurnalikt/article/view/7741)
('The composition and distribution of benthic foraminifera at coral reef ecosystem in the Thousand Islands'. Benthic foraminifera from 11 stations at Karang Bongkok, Pramuka, and Onrust Island. Highest composition of symbiont-bearing foraminiferal assemblages associated with reef ecosystem in East Pramuka (78%) and lowest in South Onrust (22%). Opportunistic types highest in S Onrust, indicating high nutrients)

Troelstra, S.R., H.M. Jonkers & S. de Rijk (1996)- Larger Foraminifera from the Spermonde Archipelago (Sulawesi, Indonesia). Scripta Geologica 113, p. 93-120.
(online at: https://repository.naturalis.nl/pub/317518)
Modern larger foram distribution in Spermonde reefal province off SW tip of Sulawesi. Near-reef facies dominated by few species (Calcarina, Elphidium). Mid-shelf reefs mainly with Heterostegina depressa and Amphistegina radiata. Outer platform reefs more open ocean influence with Amphisorus hemprichii and Amphistegina lessonii)

Troelstra, S.R., G.J. Klaver, A. Kleijne, D. Kroon, L.J. van Marle, W. Meyboom, P.J. van de Paverd, M. Situmorang & I.M. van Waveren (1989)- Actuomicropalaeontology and sediment distribution of three transects across the Banda Arc, Indonesia (Snellius-II expedition, cruise G5). Proc. Snellius II Symposium, Jakarta 1987, Netherlands J. of Sea Research 24, 4, p. 477-489.
(online at: https://www.researchgate.net/publication/222322748_Actuomicropalaeontology_and_sediment_distribution_of_three_transects_across_the_Banda_Arc_Indonesia_Snellius-II_Expedition_Cruise_G5)
(Results of 1984/1985 systematic sampling along three traverses across Banda Arc resulted in recognition of distinct litho- and biofacies. Depths of lysocline and CCD determined at 1700 and 3300 m respectively. With table of benthic foram depth ranges. Peak radiolaria in sediments between 2000-3000m. Etc.)

Troelstra, S.R. & D. Kroon (1989)- Note on extant planktonic foraminifera from the Banda Sea, Indonesia (Snellius-II Expedition, Cruise G-5). In: J.E. van Hinte et al. (eds.) Proc. Snellius II Symposium, Jakarta 1987, Netherlands J. of Sea Research 24, p. 459-463.
(Recent planktonic foraminifers collected from oligotrophic Banda Sea surface waters during NW monsoon (April/May 1985) show five associations which correspond to measured sea surface temperatures)

Tulung, G.F., J.M. Mamuaja, R.M. Rampengan, R. Djamaluddin & R.C. Kepel (2023)- Komposisi foraminifera bentik besar pada sedimen mangrove. Jurnal Pesisir dan Laut Tropis 11, 1, p. 38-49.
(online at: https://ejournal.unsrat.ac.id/v3/index.php/jplt/article/download/53332/45065/127891)
('Composition of large benthic foraminifera in mangrove sediments', In Manado Bay, N Sulawesi, dominant foraminifera genera (>5%) at two mangrove study sites were Ammonia, Amphistegina, Calcarina, Elphidium and Neorotalia)

Van Benthem Jutting, W.S.S. (1925)- On molluscs of the Krakatau-isles. Treubia (Buitenzorg (Bogor), 6, 2, p. 140-145.
(First of many papers by Dutch malacologist WWSVBJ on Recent mainly non-marine molluscs from Indonesia. N.B.: W.S.S. (Woutera) van Benthem Jutting is the same person as T. (Tera) van Benthem Jutting)

Van Benthem Jutting, T. (1932)- On Prehistoric shells from Sampoeng Cave (Central Java). Treubia 14, 1, p. 103-108.
(online at: https://archive.org/details/CFGEDI193201010141932000l/page/102/mode/2up)
(Mollusc shells excavated from Neolithic rock shelter at Sampoeng village on slopes of Wilis volcano by P.V. van Stein Callenfels are mixture of marine and non-marine species. Freshwater shells were presumably used for food, while marine shells may have been valued as ornamental objects at this far-from-the-coast location)

Van Benthem Jutting, T. (1933)- Non marine mollusca from Dutch North New Guinea including an annotated list of the species of Papuina. Nova Guinea (Zoology) 17, p. 71-150.

Van Benthem Jutting, W.S.S. (1934)- Non marine Mollusca from Nias Island. Miscellanea Zoologica Sumatrana, Medan, 84/85, p. 1-17.

Van Benthem Jutting, W.S.S. (1941)- Non-marine Mollusca from the satellite islands surrounding Java. Archives Neerlandaises de Zoologie, Leiden, 5, 3, p. 251-348.
(online at: https://www.delpher.nl/nl/boeken/view?identifier=MMKB31:047144000:00006)
(Non-marine molluscs from series of islands around Java, incl. Meeuwen Eiland, Prinsen Eiland, Popole, Krakatau, Sebesi, Dwars in den Weg, Pulau Seribu and Bay of Batavia islands, Karimon Djawa, Bawean, Madura, Kangean, Nusa Baron, Nusa Kembangan, Trowers, etc. Non-marine molluscs and land snails generally migrated by currents mainly from Java, possibly also Sumatra or Borneo. No plates?)

Van Benthem Jutting, T. (1941)- On a collection of non-marine mollusca from the Talaud Islands and from Morotai (Moluccas). Treubia, Bogor,18, 1, p. 1-27.
(online at: http://e-journal.biologi.lipi.go.id/index.php/treubia/article/view/2593/2231)
(Freshwater molluscs from northern Moluccas islands collected by H.J. Lam in 1926)

Van Benthem Jutting, W.S.S. (1948)- On the present state of the malacological research in the Malay Archipelago. Chronica Naturae 104, 5, p. 129-138.

Van Benthem Jutting, W.S.S. (1950)- Systematic studies on the non-marine mollusca of the Indo-Australian Archipelago. II. Critical revision of the Javanese Pulmonate land-shells of the Families Helicariontdae, Pleurodontidae, Fruticicolidae and Streptaxida. Treubia (Museum Zoologicum Bogoriense), Bogor, 20, 3, p. 381-505.
(online at: https://ejournal.brin.go.id/treubia/article/view/10016/7711)
(Extensive review of Recent pulmonate land snails on Java)

Van Benthem Jutting, W.S.S. (1950)- Systematic studies on the non-marine mollusca of the Indo-Australian Archipelago. IV. Critical revision of the freshwater bivalves of Java. Treubia (Museum Zoologicum Bogoriense), Bogor, 22, 1, p. 19-73.
(online at: https://e-journal.biologi.lipi.go.id/index.php/treubia/article/view/1562/1453)
(Review of 16 species of modern freshwater bivalves from lakes/ rivers of Java. Three families: Unionidae, Corbiculidae, Sphaeriidae (Pisidium spp.))

Van Benthem Jutting, W.S.S. (1956)- Systematic studies on the non-marine mollusca of the Indo-Australian Archipelago. V. Critical revision of the Javanese gastropods. Treubia (Museum Zoologicum Bogoriense), Bogor, 23, 2, p. 259-477.
(online at: http://e-journal.biologi.lipi.go.id/index.php/treubia/article/view/2713/2325)
(Review of modern freshwater gastropods (89 species) and land gastropods (171 species) of Java)

Van Benthem Jutting, W.S.S. (1958)- Landmollusken von Sumba. Wissenschaftliche Ergebnisse der Sumba-Expedition des Museums fur Volkerkunde und des Naturhistorischen Museums in Basel 1949, Verhandlungen der Naturforschenden Gesellschaft Basel 69, 1, p. 90-117.
(‘*Land molluscs from Sumba’*)

Van Benthem Jutting, W.S.S. (1958)- Non-marine Mollusca of the island of Misool. Nova Guinea, New Series, Leiden, 9, 2, p. 293-338.

Van Benthem Jutting, W.S.S. (1959)- Catalogue of the non-marine mollusca of Sumatra and of its satellite islands. Beaufortia, Zoological Museum Amsterdam, 7, 83, p. 41-191.
(online at: www.repository.naturalis.nl/document/548339)

Van Benthem Jutting, W.S.S. (1959)- Non-marine mollusca of the North Moluccan islands Halmahera, Ternate, Batjan and Obi. Treubia (Museum Zoologicum Bogoriense) 25, p. 25-87.
(online at: http://e-journal.biologi.lipi.go.id/index.php/treubia/article/view/2731/2341)

Van Benthem Jutting, W.S.S. (1963-1965)- Non-marine mollusca of West New Guinea. Nova Guinea (Zoology), vols. 20, 23, 26, 32.
(Part 1, Mollusca from fresh and brackish waters, vol. 20, p. 409-521. Part 2, Operculate land shells, vol. 23, p. 653-726. Part 3, Pulmonata, 1, vol. 26, p. 1-74. Part 4, Pulmonata, 2, vol. 32, p. 205-304)

Van Benthem Jutting, W.S.S. (1973)- Systematic studies on the non-marine mollusca of the Indo-Australian Archipelago. Linnaeus Press, Amsterdam, p. 1-477.
(Reprint of 5 papers on Recent fresh and brackish water molluscs from Indonesia, originally published in 'Treubia' (vol.19 (1948; p. 539-604; 20 (1950; p. 381-505; 21 (1952; p. 291-435; 22 (1953; p. 19-73) and 23 (1956; p. 259-477) (see also above)

Van den Bold, W.A. (1950)- *Hemikrithe*, a new genus of ostracoda from the Indopacific. Annals and Magazine Natural History 12, 3, p. 900-904.
(Short note listing 44 species of ostracods from two samples, one off Sarawak, one from Batu Island, W coast of Sumatra. Incl. new genus Hemikrithe orientalis)

Van de Paverd, P.J. & K.R. Bjorklund (1989)- Frequency distribution of polycystine radiolarians in surface sediments of the Banda Sea, Eastern Indonesia. In: J.E. van Hinte et al. (eds.) Proc. Snellius II Symposium, Jakarta 1987, Netherlands J. of Sea Research 24, 2, p. 511-521.
(Numbers of radiolarians in sea-floor sediments of Banda Sea vary widely: low from 0-950m, high between 950-4800m, and low again below 4800m water depth. Distribution reflect sediment influx and occurrence of highly productive areas in surface water)

Van de Paverd, P.J. & K.R. Bjorklund (1996)- Preservation and density of Late Quaternary radiolaria in piston cores from the Banda Sea, eastern Indonesia. Revista Espanola Micropaleontologia 28, 3, p. 139-152.

Van der Kaars, W.A. (1991)- Palynology of eastern Indonesian marine piston-cores: a Late Quaternary vegetational and climatic record for Australasia. Palaeogeogr. Palaeoclim. Palaeoecology 85, p. 239-302.
(Pollen analyses on Late Quaternary sediments from E Indonesia marine piston cores show vegetation and environmental record for E Indonesia and N Australia. On Halmahera and N Australia montane oak forest largely replaced tropical lowland vegetation during last glacial period, while climate was cooler and drier than today, with maximum grassland cover at ~18 ka. One piston-core (G6-4) extends to 300 ka. and also shows glacial periods characterised by expanding grassland vegetation (Graminae pollen peaks), and during interglacials increases in woodland and fern cover. Mangrove vegetation expansions suggest rises in sea-level at ~244, 220 and 130 ka)

Van der Kaars, S. (1998)- Marine and terrestrial pollen records of the last glacial cycle from the Indonesian region: Bandung Basin and Banda Sea. Palaeoclimates 3, p. 209-219.

Van der Kaars, S.F. Bassinot, P. de Deckker & F. Guichard (2010)- Changes in monsoon and ocean circulation and the vegetation cover of southwest Sumatra throughout the last 83,000 years: the record from marine core BAR94-42. Palaeogeogr. Palaeoclim. Palaeoecology 296, p. 52-78.

(Palynological record from deep-sea core off SW Sumatra used to reconstruct monsoon circulation and vegetation of SW Sumatra over the last 83 ky. During marine isotope stage (MIS) 5a, SW Sumatra was covered by rainforest. During MIS 4 conditions became drier, cooler and weaker monsoon. Vegetation most open during MIS 3, between ~52- 43 ky, driest of last glacial, also increase in montane pollen. After ~43 ky everwet climate gradually developed as monsoonal circulation intensified)

Van der Kaars, S. & R. Dam (1997)- Vegetation and climate change in West-Java, Indonesia during the last 135,000 years. Quaternary International 37, p. 67-71.
(Sediment cores from intramontane Bandung basin (W Java) provide paleoclimatic record for last 135,000 years. Anomalously dry conditions for penultimate glacial last glacial periods, very warm and humid conditions during last interglacial. For Last Glacial Maximum temperatures 4-7°C lower than present)

Van der Kaars, S., D. Penny, J. Tibby, J. Fluin, R.A.C. Dam & P. Suparan (2001)- Late Quaternary palaeoecology, palynology and palaeolimnology of a tropical lowland swamp: Rawa Danau, West-Java, Indonesia. Palaeogeogr. Palaeoclim. Palaeoecology 171, 3-4, p. 185-212.
(Open vegetation and drier climate suggested for Late Glacial time in Java, but no evidence for cooler conditions. Onset of Holocene coincides with change to more humid conditions. Changes in diatom composition reflect shallowing of lake)

Van der Kaars, S., X. Wang, A.P. Kershaw, F. Guichard & D.A. Setiabudi (2000)- A late Quaternary palaeoecological record from the Banda Sea, Indonesia; patterns of vegetation, climate and biomass burning in Indonesia and northern Australia. Palaeogeogr. Palaeoclim. Palaeoecology 155, p. 135-153. *(online at: https://www.academia.edu/17853126/A_Late_Quaternary_palaeoecological_record_from_the_Banda_Sea_Indonesia_patterns_of_vegetation_climate_and_biomass_burning_in_Indonesia_and_northern_Australia)*
(Banda Sea core SHI-9014 palynological and carbon isotope analyses provide a regional vegetation, fire and climate history for Banda Sea in last 180,000 years. During last two glacial periods drier climates in both E Indonesia and N Australia and lower montane forests expanded in E Indonesia indicating cooler climatic conditions. Before 37,000 yr BP. Dipterocarpaceae important part of lowland vegetation of E Indonesia. Subsequent demise likely related to increased human impact)

Van der Meer Mohr, J.C. (1927)- Note preliminaire sur la composition conchyliologique d'un amas prehistorique de coquilles pres de Medan (Sumatra). Miscellanea Zoologica Sumatrana, Medan, 18, p. 1-3.
('Preliminary note on shell composition of a prehistoric shell mound near Medan (Sumatra)'. Identifications of edible mollusc shells in 7m high and 50-60m long prehistoric shell midden/mound ('kjokkenmodding') in N Sumatra, at request of archeologist P.V. van Stein Callenfels. This mound is 'kitchen trash' from human settlements in coastal zone. Shells dominated by shallow marine-estuarine Meretrix meretrix (popularly known as 'Asiatic hard clam' from Venus group; 93%) and Ostrea (oyster; 6%), also some Arca and gastropods (NB: many original mounds destroyed by later mining activities for road cover and lime). Sumatra shell middens now 10-15 km inland from E coast of N Sumatra, in zone ~130 km long, and all located <5 m above sea level. Most recent estimate of age is ~6000 yr B.P., during mud-Holocene highstand. Similar shell mounds known from Malay Peninsula side of Malacca Straits) (Schurmann 1928, 1931, Van Heekeren 1972, Foo Shu Tien 2015))

Van Iperen, J.M., A.J van Bennekom & T.C.E. van Weering (1993)- Diatoms in surface sediments of the Indonesian Archipelago and their relation to hydrography. In: H. van Dam (ed.) Twelfth International Diatom Symposium, Renesse 1992, Hydrobiologia 269-270, 1, p. 113-128.
(Recent marine diatoms from 53 seafloor samples between 350-7200m water depth in Indonesian Archipelago, collected during Snellius II Expedition. Three assemblages, related to overlying water mass: (1) warm saline surface waters of Pacific and Indian Ocean origin (Thalassiosira oestrupii, Rhizosolenia bergonii); (2) low-salinity lobe in Makassar Strait ((Thalassionema frauenfeldii, Cyclotella striata); (3) seasonal upwelling areas in Arafura Sea and S of Java (Thalassionema nitzschioides, Chaetoceros resting spores). Also three groups of allochthonous species, indicators of productivity in littoral environment, bottom currents and river outflow)

Van Marle, L.J. (1988)- Bathymetric distribution of benthic foraminifera on the Australian- Irian Jaya continental margin, Eastern Indonesia. Marine Micropaleontology 13, 2, p. 97-152.

(Study of distribution of 164 species of benthic foraminifera in 35 seafloor samples from Australia- Irian Jaya continental margin between 60-2119 m water depth, along three transects across Banda Arc. Four fauna1 depth-zones and four subzones distinguished)

Van Marle, L.J. (1989)- Benthic foraminifera from the Banda Arc region, Indonesia, and their paleobathymetric significance for geologic interpretations of the Late Cenozoic sedimentary record. Ph.D. Thesis Vrije Universiteit, Free University Press, Amsterdam, p. 1-271.
(Collection of 11 papers, also published elsewhere, on modern foraminifera distribution and Neogene stratigraphy of E Indonesian islands)

Van Marle, L.J. (1991)- Eastern Indonesian, Late Cenozoic smaller benthic foraminifera. Verhandelingen Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, Afd. Natuurkunde 1, 34, p. 1-328.
(online at: https://dwc.knaw.nl/DL/publications/PU00011017.pdf)
(Key publication on taxonomy and distribution of Miocene- Recent deep water benthic foraminifera in E Indonesia, based mainly Recent seafloor sediments collected during the 1984-1985 Indonesian-Dutch Snellius II Expedition across basins of Eastern Indonesia, and Neogene ourcrop sediments from associated land expeditions)

Van Marle, L.J., J.E.van Hinte & A.J. Nederbragt (1987)- Plankton percentage of the foraminiferal fauna in seafloor samples from the Australian-Irian Jaya continental margin, Eastern Indonesia. Marine Geology 77, p. 151-156.
(Plankton percentage of foram fauna in 36 seafloor samples between 40-2119m depth from Australian-Irian Jaya continental margin increases with water depth. Percentage- Depth Transform derived from data set. Examples: around 100m water depth plankton% ~50%, below 500m >90%)

Van Steenis, C.G.G.J. (1934)- On the origin of the Malaysian mountain flora, part 1, Facts and statements of the problem. Bull. Jardin Botanique Buitenzorg, ser. 3, 13, p. 135-262.
(On distribution and origin of recent SE Asian mountain plants)

Van Steenis, C.G.G.J. (1935)- On the origin of the Malaysian mountain flora, part 2, Altitudinal zones, general consideration and renewed statement of the problem. Bull. Jardin Botanique Buitenzorg, ser. 3, 13, p. 289-290.

Van Steenis, C.G.G.J. (1936)- On the origin of the Malaysian mountain flora, part 3, Analysis of floristical relationships. Bull. Jardin Botanique Buitenzorg, ser. 3, 14, p. 56-72.
(Malesian (= Indonesian) mountain flora reached Indonesian archipelago along three migration routes, called Sumatra, Luzon and Papuan tracks)

Van Steenis, C.G.G.J. (1948)- Hoofdlijnen van de plantengeografie van de Indische archipel op grond van de verspreiding van de Phanerogamen geslachten. Tijdschrift Koninklijk Nederlands Aardrijkskundig Genootschap (2) 65, p. 193-208.
('The main lines of the plant geography of the Indies Archipelago based on the distribution of Phanerogams (seed plants) genera')

Van Steenis, C.G.G.J. (1950)- The delimitation of Malaysia and its main plant geographical divisions. Flora Malesiana, Ser. I, 1, p.
(see also Raes & Van Welzen, 2009)

Van Steenis, C.G.G.J. (1972)- The mountain flora of Java. E.J.Brill, Leiden, p. 1-89.
(online at: https://bsi.gov.in/uploads/userfiles/file/Rare%20Books/The%20Mountain%20Folra%20Of%20Java.pdf)
(Elegant review of modern Java mountain floras, with useful introductions on the volcanic mountain environments, climate, forest types and the works of early Java botanists, like Horsfield, Reinwardt, Kuhl & Van Hasselt, Blume, Junghuhn, Zollinger, Treub, Docters van Leeuwen, etc. Significant difference between flora of

W and E Java due to different climates, i.e. everwet in W Java (allowing for richer flora), seasonal dry in E part)

Van Steenis, C.G.G.J. (1979)- Plant-geography of East Malesia. Botanical J. Linnean Society 79, 2, p. 97-178.
(Modern plant distribution in E Indonesian archipelago (Lesser Sunda Islands; not Malaysia). Two main contacts between Malesian (Indonesian) and Australian floras must have occurred, (1) in Upper Cretaceous-Paleocene or earlier, and (2) before End-Miocene (abundant influx of Asian elements into Australian flora))

Van Waveren, I. (1989)- Pattern analysis of organic component abundances from deltaic and open marine deposits: palynofacies distribution (East Java, Indonesia). In: J.E. van Hinte et al. (eds.) Proc. Snellius II Symposium, Jakarta 1987, Netherlands J. of Sea Research 23, 4, p. 441-447.
(Eleven types of organic debris types in sea floor samples from Java Sea, off Solo River Delta, Porong Delta, etc. Mix of open marine (foraminifera, dinoflagellates) and land-derived material (spores-pollen, etc.))

Van Waveren, I. (1989)- Palynofacies analysis of surface sediments from the Northeastern Banda Sea (Indonesia). Proc. Snellius II Symposium, Jakarta 1987, Netherlands J. of Sea Research 24, 4, p. 501-509.
(Analysis of organic matter types in 31 deep water sea floor samples E of Seram)

Van Waveren, I. (1993)- Planktonic organic matter in surficial sediments of the Banda Sea (Indonesia); a palynological approach. Ph.D. Thesis University of Utrecht, Geologica Ultraiectina 104, p. 1-237.
(online at: https://dspace.library.uu.nl/handle/1874/274671)
(Collection of seven publications, six of which deal with Recent Banda Sea palynomorphs, organic matter, tintinnomorphs and dinoflagellate cysts)

Van Waveren, I. (1994)- Tintinnomorphs from deep-sea sediments of the Banda Sea (Indonesia). Scripta Geologica 105, p. 27-51.
(online at: https://repository.naturalis.nl/pub/317483)
(64 types of chitinous remains of tintinnomorph protozoans in (sub-)Recent sediments of Banda Sea. Presence and preservation may be related to effects of high productivity and high sedimentation rates in Banda Sea)

Van Waveren, I. & H. Visscher (1994)- Analysis of the composition and selective preservation of organic matter in surficial deep-sea sediments from a high-productivity area (Banda Sea, Indonesia). Palaeogeogr. Palaeoclim. Palaeoecology 112, 1-2, p. 85-111.
(online at: https://www.researchgate.net/publication/229095304_Analysis_of_the_composition_and_selective_preservation_of_organic_matter_in_surficial_deep-sea_sediments_from_a_highproductivity_area_Banda_Sea)
(Palynological analysis of box-core samples from deep-sea sediments along three transects in Banda Sea)

Van Zeist, W. (1984)- The prospect of palynology for the study of prehistoric man in Southeast Asia. Modern Quaternary Research in Southeast Asia 8, Balkema, Rotterdam, p. 1-15.
(Early review of use of palynology to identify paleo-vegetational changes in Quaternary sediments of Indonesia, with examples from Java, Sumatra, PNG)

Van Zeist, W., N.A. Polhaupessy & I.M. Stuijts (1979)- Two pollen diagrams from West Java, a preliminary report. Modern Quaternary Research in Southeast Asia 5, Balkema, Rotterdam, p. 43-56.
(Palynology of two shallow core holes in Holocene lake deposits at Situ Gunung and Telaga Patengan, SW Java)

Varol, O. (1985)- Distribution of calcareous nannoplankton in surface sediments from intertidal and shallow marine regimes of a marginal sea: Jason Bay, South China Sea. Marine Micropaleontology 9, p. 369-374.
(Sediments collected from intertidal and shallow marine (0-20m) parts of Jason Bay, S China Sea contain calcareous nannoplankton assemblages with 99% Gephyrocapsa oceanica and rare Helicosphaera carteri, Umbilicosphaera sibogae, Scapholithus fossilis, Cyclococcolithus leptoporus, Syracosphaera pulchrae. Nannoplankton species abundance increases with depth, becoming abundant below ~20m)

Vavra, V. (1906)- Ostracoden von Sumatra, Java, Siam, den Sandwich-lnseln und Japan (Reise von Dr. Walter Voltz). Zoologische Jahrbucher, Systematik Okologie und Geographie der Tiere 23, p. 413-436.
(online at: https://babel.hathitrust.org/cgi/pt?id=umn.31951002700313r&seq=513)
('Ostrocodes from Sumatra, Java, Thailand, the Hawai islands and Japan (journey of Dr. Walter Voltz)'. On Recent fresh-water ostracodes from swamps, lakes of these areas. With two plates)

Verheij, E. & P.L.A. Erftemeijer (1993)- Distribution of seagrasses and associated macroalgae in South Sulawesi, Indonesia. Blumea: Biodiversity, Evolution and Biogeography of Plants, 38, 1, p. 45-64.
(online at: https://repository.naturalis.nl/pub/524525/BLUM1993038001005.pdf)
(Eleven seagrass species found in Spermonde Archipelago, off South Sulawesi)

Victor, R. & C.H. Fernando (1979)- On some freshwater ostracod type specimens from Indonesia. Canadian Journal of Zoology 57, 1, p. 6-12.
(Re-description of some modern freshwater ostracod species from Sulawesi and Sumatra, originally described by Moniez (1892) (Strandesia, Hemicyprus, Cypretta))

Victor, R. & C.H. Fernando (1981)- Freshwater ostracods (Crustacea: Ostracoda) of the subfamily Cyprinotinae Bronstein, 1947 from Malaysia, Indonesia and the Philippines. Hydrobiologia 83, 1, p. 11-27.
(Recent Cyprinotus, Hemicypris and Heterocypris from ponds in W Indonesia, etc.)

Villain, J.M. (1995)- Modeles micropaleontologiques recents et stratigraphie sequentielle en Indonesie. In: M. Gayet et al. (eds.) First European Palaeontological Congress, Lyon 1993, Geobios, Memoire Special 18, p. 409-423.
('Recent micropaleontological models and sequence stratigraphy in Indonesia'. Foraminifera distribution on E Kalimantan shelf between Mahakam delta and Makassar Strait. Deltaic assemblages arranged according to salinity. Inner shelf with larger Rotaliidae towards euryhaline conditions. Operculina typical of shallow marine low-oxygen organic-rich clays; coarser seafloor rich in oxygen with Amphisteginids. Nodosariidae and planktonics bathymetric markers on slope. In bathyal areas mostly agglutinants. Model above valid only for highstand situations, comparable to present day. Sediments deposited during last lowstand period cored and correlated. Shelf microfaunas thin-walled, due to low oxygen and low carbonate concentrations; Rotaliidae indicate low salinities at shelf edge, where they coexist with Operculina and Amphistegina, close to deeper facies with planktonics, Buliminidae and Nodosariidae)

Von Martens, E. (1897)- Suss- und Brackwasser-Mollusken des Indischen Archipels. In: M. Weber (ed.), Zoologische Ergebnisse einer Reise in Niederlandisch Ost-Indien herausgegeben von Dr. Max Weber. Vierter Band, E.J. Brill, Leiden, p. 1-331.
(online at: https://www.zobodat.at/pdf/MON-Z-DIV_0063_0001-0453.pdf)
('Fresh- and brackish water molluscs of the Netherlands Indies Archipelago'. Described by German zoologist from Berlin, Prof. Eduard von Martens. Specimens presumably collected by Dr. Max Weber)

Vozenin-Serra, C. & C. Prive-Gill (1991)- Les terrasses alluviales pleistocenes du Mekong (Cambodge). I. Bois silicifies homoxyles recoltes entre Stung-Treng et Snoul. Review Palaeobotany Palynology 67, 1/2, p. 115-132.
('The Pleistocene alluvial terraces of the Mekong (Cambodia), I. Homoxyxl silified woods collected between Stung-Treng et Snoul'. Numerous wood remains from '+40m' Mekong Pleistocene terraces (Araucarioxylon spp., etc.))

Vozenin-Serra, C. & C. Prive-Gill (1991)- Les terrasses alluviales pleistocenes du Mekong (Cambodge). II. Bois silicifies heteroxyles recoltes entre Stung-Treng et Snoul. Review Palaeobotany Palynology 68, 1/2, p. 87-117.
('The Pleistocene alluvial terraces of the Mekong (Cambodia), II. Heteroxyxl silified woods collected between Stung-Treng et Snoul')

Vyverman, W. (1991)- Diatoms from Papua New Guinea. Bibliotheca Diatomologica 22, p. 1-224.
(Recent diatoms from lakes, etc., in PNG)

Vyverman W. & K. Sabbe (1995)- Diatom-temperature transfer functions based on the altitudinal zonation of diatom assemblages in Papua New Guinea: a possible tool in the reconstruction of regional palaeoclimatic changes. J. Paleolimnology 13, p. 65-77.

Wagey, G.A. (2002)- Ecology and physiology of phytoplankton in Ambon Bay, Indonesia. Ph.D. Thesis University of British Columbia, p. 1-185.
(105 phytoplankton species identified from Ambon Bay, including several dinoflagellates new to Indonesia)

Waller, H.O. (1960)- Foraminiferal biofacies off the South China Coast. Journal of Paleontology 34, 6, p. 1164-1182.
(Benthic foraminifera from shelf S of Taiwan and in Gulf of Tonkin. Four depth-related faunas: (1) Inner Shelf (65-150') Elphidium advenum, E. sagrum, Nonion japonius, Quinqueloculina; (2) C Shelf (151-275') Amphistegina lessonii, Hanzawaia nipponica, Streblus tepidus, Operculina bartschi; (3) Outer Shelf (276-400') Biloculinella labiata, Cassidulina neocarinata, Spiroloculina communis; (4) U Bathyal (401-656') Bolivina spathylata, Uvigerina auberiana, U. schwageri)

Wang, P. & J. Chappell (2001)- Foraminifera as Holocene environmental indicators in the South Alligator River, Northern Australia. Quaternary International 83-85, p. 47-62.
(Trends in foraminifera assemblages along 80 km length of macrotidal river E of Darwin, N coast of Australia. Due to tidal transport and resuspension, most foraminiferal thanatocoenoses in river contain many small marine forms, while % of large and heavy marine forams, like Quinqueloculina, decreases upstream)

Wang, P. & C. Samtleben (1983)- Calcareous nannoplankton in surface sediments of the East China Sea. Marine Micropaleontology 8, p. 249-259.
(28 species of coccoliths in surface sediments of East China Sea. Emiliania huxleyi and Gephyrocapsa oceanica together >90% of assemblages. Coccoliths very rare in water depths <50m). Coccolith species composition different between continental shelf and Okinawa Trough, reflecting different water masses)

Wang, R., A. Abelmann, B. Li & Q. Zhao (2000)- Abrupt variations of the radiolarian fauna at Mid-Pleistocene climate transition in the South China Sea. Chinese Science Bull. 45, 10, p. 952-955.
(Core 17957-2 from S China Sea shows distinct changes in radiolarian/foraminfera ratio and radiolarian assemblages that can be related to global climate cooling observed at M Pleistocene climate revolution at ~900 ka)

Wang, X., S. van der Kaars, P.P. Kershaw, M.I. Bird & F.A. Jansen (1999)- A record of fire, vegetation and climate through the last three glacial cycles from Lombok Ridge Core G6-4, Eastern Indian Ocean, Indonesia. Palaeogeogr. Palaeoclim. Palaeoecology 147, 3-4, p. 241-256.
(Deepsea core SW of Sumba with ~300,000 yr sediment record)

Watson, K.A. (1988)- The taxonomy and distribution of Recent reef Ostracoda from the Pulau Seribu, Java Sea. Doct. Dissertation, University of Wales, Aberystwyth, p. 1-434. *(Unpublished)*

Weber, M., (1902)- Der Indo-Australische Archipel und die Geschichte seiner Tierwelt. Summary of a presentation in Karslbald, 22 September 1902, Gustav Fisher Verlag, Jena, p. 1-46.
('The Indo-Australian Archipelago and the history of its animal world'. Intereseting early views on historcal geology and animal life in the Indonesian Archipelago. Calls Wallace's Line 'fiction, correctly characterizes Mesozoic radiolarian cherts as oceanic deposits, observed wide distribution of Triassic and Jurassic seas, etc.))

Weber- van Bosse, A. & M. Foslie (1904)- The Corallinaceae of the Siboga Expedition. In: M. Weber (ed.) Siboga Expeditie, Uitkomsten op zoologisch, botanisch, oceanographisch en geologisch gebied verzameld in Nederlandsch Oost-Indie 1899-1900, Monographie 61, Brill, Leiden, p. 1-110.

(Monograph on Recent marine coralline red algae collected by Siboga Expedition in 1899-1900. Lithothamnium group present at 55 stations, invariably in areas of strong tidal or other currents, thriving best from 10-30m, but also present on reef flats and deeper water down to ~120m)

Weber- van Bosse, A. (1932)- Resultats scientifiques du voyage aux Index Orientales Neerlandaises de LL. AA. RR. le Prince et la Princesse Leopold de Belgique, vol. 6, 1, Algues. Memoires Musee Royal d'histoire naturelle de Belgique, Brussels, Hors serie, p. 1-27.
(online at: https://ia601300.us.archive.org/14/items/resultatsscienti61stra/resultatsscienti61stra.pdf)
('Scientific results of the voyage of Prince and Princess Leopold of Belgium to the East Indies, vol. 6, 1, Algae'. Rel. brief report on Recent calcareous algae from localities across Indonesia)

Weinmann, A.E., D. Rodder, S. Lotters & M.R. Langer (2013)- Heading for new shores: projecting marine distribution ranges of selected larger foraminifera. PLoS One 8, 4, e62182, p. 1-14.
(online at: https://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0062182&type=printable)
(Modern symbiont-bearing larger foraminifera confined to tropical and subtropical shallow marine habitats, mainly between 30°N and 30°S) and minimum T limits e governed by 14-20°C isotherms. During times of extensive global warming (Eocene, Miocene), larger foraminifera found as far N as 50°N and 47°S (New Zealand). Species distribution models for Archaias angulatus, Calcarina spp., and Amphistegina spp. Amphistegina spp. widest potential distribution (T tolerance). Calcarina most tolerant to high T)

Wells, P., G. Wells, J. Calli & A. Chivas (1994)- Response of deep sea benthonic foraminifera to Late Quaternary climate changes, SE Indian Ocean, offshore Western Australia. Marine Micropaleontology 23, p. 185-229.
(Late Quaternary benthic foraminifera of four ODP 122 deep-sea cores off W Australia with two main assemblages (1) dominated by Uvigerina peregrina, (2) dominated by U. proboscidea. Episodes of high influx of particulate organic matter during glacial episodes, tied to upwelling episodes and to unusually low sea-surface paleotemperature indicated by planktic foraminifera)

Whatley, R.C. & R. Titterton (1981)- Some new Recent podocopid Ostracoda from the Solomon Islands, South-West Pacific. Revista Espanola Micropaleontologia 13, p. 157-170.

Whatley, R.C. & K. Watson (1988)- A preliminary account of the distribution of ostracoda in Recent reef and reef associated environments in the Pulau Seribu or Thousand Island Group, Java Sea. In: T. Hanai et al. (eds.) Evolutionary biology of ostracoda: its fundamentals and applications, Proc. 9th Int. Symposium on Ostracoda, Shizuoka, Elsevier, Developments in Palaeontology and Stratigraphy 11, p. 399-411.
(Samples on and around reef complex of Pulau Pari, Pulau Seribu, Java Sea, yielded 141 species of podocopid and platycopid Ostra coda. Bairdiidae maximum diversity on reef, Renaudcypris and Hansacypris mainly in intertidal zone. Loxoconcha, Xestoleberis and Ornatoleberis wider environmental tolerance)

Whatley, R.C. & Q. Zhao (1987)- Recent ostracoda of Malacca Straits (Part I). Revista Espanola Micropaleontologia 19, 3, p. 327-366.
(18 bottom samples of modern sediments from Malacca Straits over depth range of 10-100m contain 129 species of ostracodes (22 new). Faunas close affinity to South China Sea and Indonesia)

Whatley, R.C. & Q. Zhao (1988)- Recent ostracoda of Malacca Straits (Part II). Revista Espanola Micropaleontologia 20, 1, p. 5-37.

Whittaker, J.E. & R.L. Hodgkinson (1995)- The foraminifera of the Pitcairn Islands. In: T.G. Benton & T. Spencer (eds.) Pitcairn Islands: biogeography, ecology and prehistory, Biol. J. Linnean Society 56, p. 365-371.
(Recent foraminifera from Pitcairn Islands, Pacific Ocean. Living forams almost exclusively from phytal (attached or clinging) habitats. Foraminifera in sediment samples mainly thanatocoenoses. Fauna all calcareous, low diversity, dominated by large soritids (Marginopora, Amphisorus, Sorites) and Amphistegina, with small miliolids and small attached genera (discorbids, etc.). Apparent absence of Calcarina, small rotaliids, elphidiids and agglutinating species, common in W Pacific islands)

Wicaksono, S.A., J.M. Russell & S. Bijaksana (2015)- Compound-specific carbon isotope records of vegetation and hydrologic change in central Sulawesi, Indonesia, since 53,000 yr BP. Palaeogeogr. Palaeoclim. Palaeoecology 430, p. 47-56.
(Carbon isotopic composition (δ13C) of terrestrial leaf waxes in sediment cores from Lake Matano spanning 53 kyr. During Marine Isotope Stages 1 and 3, more negative δ13Cwax values indicate closed-canopy rainforests dominated in Sulawesi, in wetter, less seasonal climate. More abundant open canopy vegetation and possible expansion of C4 grasses between 29-14 ka BP, indicating more arid climate in Marine Isotope Stage 2 (incl. Last Glacial Maximum). Higher elevations maintain rainforest refugia during regionally arid time intervals when C4 savannas and grasslands expanded at lower elevations)

Wijono, S. (1991)- Distribusi foraminifera bentonik di daerah peraian P. Papateo, Kepulauan Seribu, Laut Jawa. Media Teknik (UGM) 13, 2, p. 119-131.
'Distribution of benthic foraminifera off Papateo Island, Pulau Seribu, Java Sea'. Forams from 18 samples from 21-30m depth)

Winantris (2011)- Fungal spore sedimen Resen delta front delta Mahakam Kalimantan Timur. Bull. Scientific Contribution (UNPAD) 9, 2, p. 107-120.
(online at: http://jurnal.unpad.ac.id/bsc/article/view/8267/3814)
('Fungal spores in Recent sediments of the Mahakam delta from, East Kalimantan'. Terrestrial and marine fungal spores present in delta front sediments, incl. Monoporisporites, Inapertisporites, Biporipsilonites)

Winantris (2012)- Kelimpahan polen dan spora endapan channel Delta Mahakam. Bull. Scientific Contribution (UNPAD) 10, 2, p. 89-95.
(online at: http://jurnal.unpad.ac.id/bsc/article/view/8280/3827)
('Abundance of pollen and spores in channels deposits of the Mahakam Delta'. Abundance of pollen higher in tidal channel deposits than in distributary channels. Oncosperma tigillarium and Nypa fruticans palmae pollen dominate in distributary channels, Rhizophora and Avicennia mangrove pollen dominant in tidal channels)

Winantris & L. Jurnaliah (2018)- Pollen and foraminifera approaches to identify sediment sources in the river mouth Mahakam, East Kalimantan. J. Geoscience Engineering Environment Technology (JGEET) 2, 4, p. 242-248.
(online at: http://journal.uir.ac.id/index.php/JGEET/article/view/689/627)
(Shallow core from Mahakam River before delta plain. Common pollen (incl. mangrove), rel. rare, simple foraminifera (Oolina, Ramulina, Stictogongylus vandiemenensis, etc.))

Winantris, A. Sudradjat, I. Syafri & A.T.Rahardjo (2014)- Diversitas polen Palmae pada endapan delta Mahakam Resen. Proc. 43rd Annual Conv. Indonesian Association Geologists (IAGI), Jakarta, PIT IAGI 2014-195, p. 1-5.
(online at: https://www.iagi.or.id/web/digital/6/2014_IAGI_Jakarta_Diversitas-Polen-Palmae.pdf)
('Diversity of Palmae pollen in Recent deposits of the Mahakam Delta'. Sediments of Mahakam Delta contain 28 palm pollen species in delta plain, 23 in delta front. Palm pollen can help differentiate between delta plain and delta front facies: (1) delta plain dominated by Oncosperma tiggilarium, Nypa fruticans, Eugessona insignis, Pinanga parvula and Cocos nucifera; (2) delta front dominated by Oncosperma tiggilarium, Nypa fruticans, Eugessona insignis, Phoenix paludosa and Licuala sp. Palm pollen abundance nearly 4x greater in delta plain than in delta front, also higher diversity)

Winantris, I. Syafri & A.T. Rahardjo (2012)- *Oncosperma tigillarium* merupakan bagian palino karakter delta plain di Delta Mahakam, Kalimantan. Bionatura-Jurnal Ilmu-ilmu Hayati dan Fisik 14, 3, p. 228-236.
(online at: http://jurnal.unpad.ac.id/bionatura/article/viewFile/7465/3426)
('Oncosperma tigillarium is part of palyno character of delta plain in Mahakam Delta, Kalimantan'. O. tigillarium pollen of coastal palm tree widespread in Recent delta plain samples, but absent in delta front, therefore good indicator of (upper) delta plain environment)

Woodroffe, S.A., B.P. Horton, P. Larcombe & J.E. Whittaker (2005)- Intertidal mangrove foraminifera from the central Great Barrier Reef shelf, Australia: implications for sea-level reconstruction. J. Foraminiferal Research 35, 3, p. 259-270.
(Foraminifera distribution in intertidal zone tied to elevation. Agglutinated foram assemblage of Miliammina fusca, Trochammina inflata, Ammotium and Haplophragmoides between just above Mean Low Water of Neap Tides to Highest Astronomical Tide level (vertical range 1.8 m). Ammonia aoteana- dominated assemblage between just below Mean Low Water of Neap Tides and Mean High Water of Neap Tides (vertical range 0.8 m))

Wu, R., Y. Gao, Q. Fang, C. Chen, B. Lan, L. Sun & D. Lan (2013)- Diatom assemblages in surface sediments from the South China Sea as environmental indicators. Chinese J. Oceanology Limnology 31, 1, p. 31-45.
(online at: http://dspace.xmu.edu.cn/bitstream/handle/2288/53977/Diatom_assemblages_in_surface_sediments_from_the_South_China_Sea_as_environmental_indicators.pdf)
(Diatoms in 62 surface sediment samples from depths from 101-4185m. 256 species, dominated by Coscinodiscus africanus, Coscinodiscus nodulifer, Cyclotella stylorum, Hemidiscus cuneiformis, Melosira sulcata, Nitzschia marina, Roperia tesselata, Thalassionema nitzschioides, etc. Seven zones)

Xu, J., P. Wang, B. Huang, Q. Li & Z. Jian (2005)- Response of planktonic foraminifera to glacial cycles: Mid-Pleistocene change in the southern South China Sea. Marine Micropaleontology 54, p. 89-105.
(online at: https://pxwang.tongji.edu.cn/__local/6/CE/0A/8C788E7E7091548CC36EC244794_3E Etc.)
(Planktonic foraminifera from ODP Site 1143 in S South China Sea show faunal response to glacial cycles in last 2.1 Ma. Globorotalia menardii rel. abundant in interglacials and low in glacials. Pulleniatina obliquiloculata low latitude species, relatively abundant during glacials and low during interglacials after ~850 ka)

Yahya, K., S. Shuib, F.I. Minhat, O. Ahmad & A. Talib (2014)- The distribution of benthic foraminiferal assemblages in the north-west coastal region of Malacca Straits, Malaysia. J. Coastal Life Medicine 2, 10, p. 784-790.
(online at: www.jclmm.com/qk/201410/5.pdf)
(Benthic foram assemblages from shallow marine environments around NW Penang island dominated by Ammonia (~80% of fauna), followed by Elphidium (~3%). Rare agglutinated taxa and Bolivina)

Yakzan, A.M., S. Jirin, S.S.M. Shah & R.J. Morley (2010)- The major trends of palynomorphs distribution in three fluvial systems, Peninsular Malaysia. Proc. Petroleum Geology Conference Exhibition (PGCE), Kuala Lumpur 2010, p. *(Extended Abstract)*
(Palynomorph distribution patterns in three fluvial systems on W (Klang-Langat River) and E (Pahang and Sedili Besar Rivers) coasts of Peninsular Malaysia. Ecological groups. mangrove (Rhizophora), back mangrove (Acrostichum, Nypa) and hinterland pollen. Pollen and spores redistributed by currents and less by wind. Sediments in offshore area contain pollen signals which approximately mirror vegetation character onshore)

Yakzan, A.M. & H. Kamaludin (2010)- Palynology of late Quaternary coastal sediments. Catena 30, 4, p. 391-406.
(Palynology of Late Pleistocene sediments from Pantai tin mine, W coast of Malay Peninsula, with freshwater Pandanus peat overlain by mangrove peat)

Yanagisawa, Y. (1987)- Age assignments of dredge and piston core samples based on diatom biostratigraphy. Coord. Comm. Joint Prospecting Mineral Resources in Asian Offshore Areas, CCOP, Technical Bull. 19, p. 73-87.
(online at: https://ncloud.dotdream.co.th/index.php/s/xCwHAtAFYxzBtC9)
(Latest Pliocene- Recent ages for sediments in piston cores from forearc area offshore E Java)

Yassini, I., B.G. Jones & M.R. Jones (1993)- Ostracods from the Gulf of Carpentaria, northeastern Australia. Senckenbergiana Lethaea 73, p. 375-406.

Yau, M.W.C. (1973)- Studies of the Recent foraminifera of the Sunda Shelf and the Java Sea. M.A. Thesis Rice University, Houston, p.
(online at: https://scholarship.rice.edu/handle/1911/104182)
(Foraminiferal assemblages on Sunda Shelf and Java Sea: (1) Inner shelf with Amphistegina radiata, Calcarina spengleri, Elphidium crispum, Operculina ammonoides and Quinqueloculina; (2) Central shelf Quinqueloculine-Rotalia spp., Asterorotalia trispinosa and Elphidium craticulatum (3) Outer shelf with Bolivina, Bulimina, Uvigerina spp., Cancris auriculus, etc. (4 l)ower middle bathyal with Bolivina-Uvigerina spp., Cassidulina subglobosa. High diversity correlates with possible relict assemblagesat lower sea-level)

Yin, J., C. Liu, J. Zhang, X. Yang, J. Wu, W. Oschmann, F.T. Fursich, B. Zhu & H. Zhang (2018)- Distribution and constraining factors of planktonic and benthic foraminifers in bottom sediments of the southern South China Sea. Palaeogeogr. Palaeoclim. Palaeoecology 502, p. 130-146.
(Water depth dominant factor controling foram assemblage composition and δ18O. Differences in proportion of agglutinated and porcelaneous tests in shallow-water zone controlled by terrestrial runoff from nearby river systems (Mekong and N Borneo rivers) and seasonal currents. Dominance of Melonis barleeanus at sites of active cold methane seepage in southern SCS)

Yordanova, E.K. & J. Hohenegger (2004)- Morphocoenoclines of living operculinid foraminifera. Micropaleontology 50, p. 149-177.
(Relations between water depth and shape in Operculina, Planoperculina and Planostegina in Ryuku islands. Thick Operculina with intensively coiled spirals predominate in shallow water (20 -40m); in deep euphotic zone (−120m) thin forms with weakly coiled spiral. Thin Planoperculina heterosteginoides restricted to deep euphotic zone (>80m) can extend distribution to just below euphotic zone, where it develops very thin tests)

Yulianto, E., A.T. Rahardjo, Dardji Noeradi, D.A. Siregar & K. Hirakawa (2005)- A Holocene pollen record of vegetation and coastal environmental changes in the coastal swamp forest at Batulicin, South Kalimantan, Indonesia. J. Asian Earth Sciences 25, 1, p. 1-8.
(Pollen analysis of coastal peat swamp core at Batulicin, SE Kalimantan, representing 9100 BP, showing Rhizophora mangrove forest since early Holocene. From ~6000 BP gradual change from mangrove forest to peat swamp forest due to higher precipitation and progradation. Human influence recognized from ~1600 BP)

Yulianto, E. & W.S. Sukapti (1998)- Perubahan iklim selama rentang Plistosen atas hingga Holosen di Indonesia berdasarkan rekaman data palinologi. Proc. 27th Annual Conv. Indonesian Association Geologists (IAGI), 2 (Sedimentology Paleontology Stratigraphy), Yogyakarta, p. 66-71.
(online at: https://www.iagi.or.id/web/digital/56/33.pdf)
('Late Pleistocene-Holocene climate change in Indonesia based on palynological data records')

Yulianto, E., W.S. Sukapti & K.T. Dewi (2019)- Late Holocene pollen record of environmental changes in Karimata Strait, Sunda Shelf region. Indonesian J. on Geoscience (IJOG) 6, 1, p. 41-55.
(online at: http://ijog.bgl.esdm.go.id/index.php/IJOG/article/view/291/277)
(Pollen analysis on 90 cm gravity core taken from Karimata Strait, 170km NW of Bangka, reveals Late Holocene environmental changes in central Sunda Shelf region. Deposition of core started at 1800 yr BP, with 50-70% mangrove pollen. Surrounding mangroves terminated by catastrophic event, possibly 535 AD Krakatau eruption)

Yulianto, E., W.S. Sukapti & Praptisih (2019)- Perubahan lingkungan dan evolusi mangrove di Pulau Belitung selama Holosen akhir berdasarkan Rekaman Polen. J. Geologi Kelautan 17, 2, p. 123-139.
(online at: http://ejournal.mgi.esdm.go.id/ejournal/index.php/jgk/article/view/578)
('Pollen record of Late Holocene environmental changes and mangrove evolution from Belitung Island, Indonesia'. Palynology data reflects coastal progradation in last ca.1500 years in N Belitung. Brief changes may be interpreted as 535-536 AD and 1883 AD Krakatau tsunamis)

Yulianto, E., W.S. Sukapti, A.T. Rahardjo, Dardji Noeradi, D.A. Siregar, P. Suparan & K. Hirakawa (2004)- Mangrove shoreline responses to Holocene environmental change, Makassar Strait, Indonesia. Review Palaeobotany Palynology 131, p. 251-268.
(Pollen analyses of two near-coastal sites at Batulicin, S Kalimantan and Pare-Pare, S Sulawesi. Mangroves developed at Batulicin in mid-Holocene, persisting to present at Batulicin. Mangrove development commenced at Pare-Pare in early Holocene, but since mid Holocene fluvial/floodplain deposition)

Zallesa, S., K.T. Dewi, N.C.D. Aryanto & R. Rahardiawan (2014)- The correlation between benthic foraminifera and sediment types of South Makassar Strait. Bulletin of the Marine Geology 29, 2, p. 53-59.
(online at: http://ejournal.mgi.esdm.go.id/index.php/bomg/article/view/65/66)
(Core top samples from water depths 200-1500m in S Makassar Straits with 38 species of benthic foraminifera. Most common species Anomalinoides colligerus, Lenticulina suborbicularis, Planulina wuellerstorfi, Pseudonodosaria discrete, Bolivina spathulata. Highest abundance of benthic foraminifera in silty sand and sandy silt)

Zamoras, L.R. & P.J. Militante-Matias (1997)- Recent Foraminifera of Matabungkay and Talim Bays, Batangas. J. Geological Society of the Philippines 52, 2, p.

Zhang, P., R. Zuraida, Y. Rosenthal, A. Holbourn, W. Kuhnt & J. Xu (2019)- Geochemical characteristics from tests of four modern planktonic foraminiferal species in the Indonesian Throughflow region and their implications. Geoscience Frontiers 10, 2, p. 505-516.
(online at: https://www.sciencedirect.com/science/article/pii/S1674987118300471)
(d18O and Mg/Ca of Globigerinoides ruber, Gs sacculifer, Pulleniatina obliquiloculata and Neogloboquadrina dutertrei from 60 coretop sediment samples from Indonesian Throughflow region suggest calcification within mixed layer for G. ruber (0-50m) and G. sacculifer (20-75 m), and within thermocline (~75-125m) for P. obliquiloculata and N. dutertrei)

Zhao, Q. & P. Wang (1988)- Distribution of modern ostracoda in the shelf seas off China. In: T. Hanai et al. (eds.) Evolutionary biology of ostracoda: its fundamentals and applications, Proc. 9th International Symposium on Ostracoda, Shizuoka 1985, Elsevier, Developments in Palaeontology and Stratigraphy 11, p. 805-821.

Zhao, Q. & R.C. Whatley (1989)- Recent podocopid Ostracoda of the Sedili River and Jason Bay, southeastern Malay Peninsula. Micropaleontology 35, p. 168-187.
(Distribution of ostracodes in Jason Bay, off SE coast of Malay Peninsula. 101 species recorded. Dead assemblages higher diversity than live assemblages, due to postmortem transportation. One new genus and 13 new species. Freshwater Sedili River only two species (Darwinula stephensoni, ?Cytherissa.), in brackish estuary 4 species (Sinocythere superba, Hemicytheridea reticulata, Keijella multisulcus, Neocytherideis sp.) Forty-eight species live in open sea, most common: Atjehella semiplicata, Cushmanidea subjaponica, Hemicytheridea spp., Javanella kendengensis, Neomonoceratina delicata, Parakrithella pseudadonta, etc.)

Zhao, Q. & R.C. Whatley (1997)- Distribution of the ostracod genera *Krithe* and *Parakrithe* in bottom sediments of the East China and Yellow Seas. Marine Micropaleontology 32, 1, p. 195-207.
(Distribution closely linked to water masses)

Zheng, S.Y. (1979)- The Recent foraminifera of the Xisha Islands, Guangdong Province, China, Part II. Studia Marina Sinica 15, p. 101-232. *(In Chinese with English summary)*

Zheng, S.Y. (1980)- The Recent foraminifera of the Zhongsha Islands, Guangdong Province, China, Part I. Studia Marina Sinica 16, p. 143-182. *(In Chinese with English summary)*

Zhou, B. (1995)- Recent ostracode fauna in the Pacific off Southwest Japan. Memoirs Faculty of Science, Kyoto University, Ser. Geology and Mineralogy 57, 2, p. 21-98.
(online at: http://repository.kulib.kyoto-u.ac.jp/dspace/bitstream/2433/186675/1/mfskugm%20057002_021.pdf)

Zhou, B. & Q. Zhao (1999)- Allochthonous ostracods in the South China Sea and their significance in indicating downslope sediment contamination. Marine Geology 156, p. 187-195.
(Modern distribution pattern of allochthonous ostracods in South China Sea: limited to continental shelf and slope, and around reef islands, suggesting ostracod shells did not travel far from source areas. Turbidites probably principal agent responsible for downslope transport of ostracods)

Zobel, B. (1978)- Deep-water Quaternary foraminifera from short cores taken between Australia and southeast Indonesia. BMR J. Australian Geology Geophysics 3, p. 345-349.
(online at: www.ga.gov.au/corporate_data/80976/Jou1978_v3_n4_p345.pdf)
(Twenty short cores from N Scott Plateau (water depths ~3200 m) and S slope of the Java Trench (4950-5790m) taken by R.V. Valdivia. 55 species of benthic foraminifera identified on plateau, less in trench (incl. Eggerela bradyi, Kareriella bradyi, Bulimina spp, Sphaeroidina bulloides, Cibicides wuellerstorfi, Globocassidulina, Gyroidinoides, Melonis pompilioides, Oridorsalis tener, etc. Only deep-water forms. Planktonic foraminifera belong to tropical associations. Holocene-Pleistocene boundary at ~60 cm in cores, and carbonate solution more marked above than below boundary)

Zong, Y. & B.H. Kamaludin (2004)- Diatom assemblages from two mangrove tidal flats in Peninsular Malaysia. Diatom Research 19, p. 329-344.

X.2. Cenozoic

Adams, C.G. (1967)- Tertiary Foraminifera in the Tethyan, American and Indo Pacific Provinces. In: C.G. Adams & D.V. Ager (eds.) Aspects of Tethyan biogeography: A Symposium, Systematics Association, London, Special Publ. 7, p. 195-217.
(Tertiary larger foraminifera three major bioprovinces: Americas, Tethys, Indo-Pacific)

Adams, C.G. (1968)- A revision of the foraminiferal genus *Austrotrillina* Parr. Bull. British Museum (Natural History), Geology, 16, p. 71-97.
(Evolutionary changes in Late Oligocene- E Miocene Austrotrillina are of value in stratigraphy)

Adams, C.G. (1970)- A reconsideration of the East Indian Letter classification of the Tertiary. Bull. British Museum (Natural History), Geology, 19, 3, p. 87-137.
(online at: https://ia800700.us.archive.org/0/items/bulletinofbritis19geol/bulletinofbritis19geol.pdf)
(Classic review and update of the Late Paleocene- Recent larger foraminifera zonation, known as East Indian Letter Classification, first proposed by Van der Vlerk & Umbgrove (1927))

Adams, C.G. (1973)- Some Tertiary foraminifera. In: A. Hallam (ed.) Atlas of Palaeobiogeography. Elsevier, Amsterdam, p. 453-468.
(Notes on biogeography of Eocene- M Miocene larger foram genera)

Adams, C.G. (1976)- Larger foraminifera and the Late Cenozoic history of the Mediterranean region. Palaeogeogr. Palaeoclim. Palaeoecology 20, p. 47-66.
(Larger foram faunas similar between Mediterranean and Indo-West Pacific, but different from Americas. From M Miocene onward diverged rapidly)

Adams, C.G. (1981)- Larger foraminifera and the Paleogene/ Neogene boundary. Proc. 7th International Congress Mediterranean Neogene, Athens 1979, Annales Geologiques des Pays Helleniques, hors serie, IV, p. 145-151.
(No major changes in larger foram faunas distribution across Oligo-Miocene boundary. In Indonesia- W Pacific first appearance of Miogypsina best marker event)

Adams, C.G. (1983)- Speciation, phylogenesis, tectonism, climate and eustasy: factors in the evolution of Cenozoic larger foraminiferal bioprovinces. In: R.W. Sims et al. (eds.) The emergence of the biosphere, Syst. Assoc. Special Volume 23, Academic Press, London, p. 255-287.
(Review of evolutionary patterns of Cenozoic larger foraminifera (Miogypsina, Cycloclypeus, Lepidocyclina, etc.), related to climate changes and tectonism. Early Miocene disconnect between Mediterranean and Indian Ocean, etc.)

Adams, C.G. (1984)- Neogene larger foraminifera, evolutionary and geological events in the context of datum planes. In: N. Ikebe & R. Tsuchi (eds.) Pacific Neogene datum planes, Contributions to biostratigraphy and chronology, University of Tokyo Press, p. 47-67.
(Thorough discussion of Oligocene- Recent larger foraminifera 'datum planes', with updated range chart)

Adams, C.G. (1987)- On the classification of the Lepidocyclinidae (Foraminiferida) with redescriptions of the unrelated Palaeocene genera *Actinosiphon* and *Orbitosiphon*. Micropaleontology 33, p. 289-317.
(Three subgenera, Lepidocyclina (Lepidocyclina), L. (Eulepidina), and L. (Nephrolepidina), discriminated on nature and arrangement of peri-embryonic chambers. Two groups of species recognized within L. (Lepidocyclina) on basis of equatorial chamber shape)

Adams, C.G. (1989)- Foraminifera as indicators of geological events. Proc. Geologists Association, London, 100, 3, p. 297-311.

Adams C.G. (1992)- Larger foraminifera and the dating of Neogene events. In: R. Tsuchi & J.C. Ingle (eds.) Pacific Neogene, University of Tokyo Press, p. 221-235.

Adams, C.G. & D.J. Belford (1974)- Foraminiferal biostratigraphy of the Oligo-Miocene limestones of Christmas Island (Indian Ocean). Palaeontology 17, p. 475-506.
(Late Oligocene- M Miocene (lower Te- Lower Tf zones) LBF assemblages in ~190m thick limestones capping a truncated basaltic volcanic cone in Indian Ocean off SW Java. Basal part rich in Sproclypeus margaritatus 'globulus', Heterostegina barriei and Gypsina globulus (Lower Te), middle part with Spiroclypeus margaritatus, Miogypsinoides bantamensis and less common Lep. (Eulepidina) and Borelis pygmaeus (Lower Te). Youngest assemblages in thin zone with Austrotrillina howchina and Flosculinella bontangensis. Mention of Eocene limestone, but no details)

Adams, C.G., J. Butterlin & B.K. Samanta (1986)- Larger foraminifera and events at the Eocene-Oligocene boundary in the Indo West Pacific region. In: C. Pomerol & I. Premoli Silva (eds.) Terminal Eocene events, Elsevier, Amsterdam, p. 237-252.
(In most Indo-Pacific localities Eocene terminated by disconformities, with extinction of Discocyclina, Pellatispira, Spiroclypeus vermicularis, etc., as in localities worldwide. Possibly triggered by global sea level fall with climatic deterioration. Only at Melinau, Sarawak, Eocene and Oligocene limestones in apparently unbroken sequence)

Adams, C.G. & P. Frame (1979)- Observations on *Cycloclypeus (Cycloclypeus)* Carpenter and *Cycloclypeus (Katacycloclypeus)* Tan (Foraminifera). Bull. British Museum (Natural History), Geology 32, 1, p. 3-17.
(online at: www.archive.org/details/bulletinofbritis32geollond)
(Genus Katacycloclypeus limited to Middle Miocene (Lower Tf letter stage). Microspheric forms from Fiji up to 90 mm in diameter)

Adams, C.G., A.W. Gentry & P.J. Whybrow (1979)- Dating the terminal Tethyan event. Utrecht Micropaleontological Bulletin 30, p. 273-298.
(Geographic distribution of larger foraminifera shows continuous connection between Mediterranean and Indian Ocean closed by mid-Burdigalian)

Adams, C.G., D.E. Lee & B.R. Rosen (1990)- Conflicting isotopic and biotic evidence for tropical sea-surface temperatures during the Tertiary. Palaeogeogr. Palaeoclim. Palaeoecology 77, p. 289-313. *(online at: https://www.researchgate.net/profile/Brianroy-Rosen-2/publication/223283317_Conflicting_isotope_ Etc)*
(Paleotemperatures derived from some isotope studies are too low to account for distribution and diversity of many Tertiary tropical- subtropical taxa)

Adams, C.G., P. Rodda & R.J. Kiteley (1979)- The extinction of the foraminiferal genus *Lepidocyclina* and the Miocene-Pliocene boundary problem in Fiji. Marine Micropaleontology 4, 4, p. 319-339.
(Last surviving species of Lepidocyclina, L. radiata, becomes extinct at N18/N19 boundary, near top Miocene)

Adisaputra, Mimin K. (1987)- Notes on *Cycloclypeus (Katacycloclypeus)* Tan and *Cycloclypeus (Radiocloclypeus)* Tan. J. RISET Geologi dan Pertambangan (LIPI) 8, 1, p. 25-33.
(Discussion of larger foram Cycloclypeus subgenera and species. Age of Katacycloclypeus is M Miocene (planktonic foram zone N8-N12). Stellate Radiocycloclypeus may be N12 and younger)

Adisaputra, Mimin K. & Hartono (2007)- Phillipsite mineral in deep sea sediment from single core in Roo Rise, Indian Ocean. Indonesian Mining J. 10, 3, p. 39-43.
(online at: http://jurnal.tekmira.esdm.go.id/index.php/imj/article/view/612/474)
(Core MD982156 in Roo Rise, Indian Ocean/ Java Trench, S of East Jawa, 30m long in water depth 3884m. Upper part of core abundant planktonic foraminifera, lower part mainly phillipsite-rich sediment (± 40%). Late Miocene or older (Paleocene according to Adisaputra & Kusnida (2010))

Adisaputra, Mimin K., N. Hasjim & A. Djojosumarto (1995)- Sundaland Neogene biostratigraphic events. In: S. Nishimura & R. Tsuchi (eds.) Proc. Oji Seminar on Neogene evolution of Pacific Ocean Gateways, Kyoto, IGCP-355, p. 62-71.

Adisaputra, Mimin K. & M. Hendrizan (2008)- Hiatus pada kala Eosen-Miosen Tengah di Tinggian Roo, Samudra Hindia, berdasarkan biostratigrafi nannoplankton. J. Geologi Kelautan 6, 3, p. 154-166.
(online at: http://ejournal.mgi.esdm.go.id/index.php/jgk/article/view/159/149)
('Eocene- Middle Miocene hiatus at the Roo Rise, Indian Ocean S of East Java, based on nannoplankton biostratigraphy'. Late Miocene and Paleocene calcareous nannoplankton from Roo Rise piston core MD982156 from MD III-IMAGES IV Expedition)

Adisaputra, Mimin K. & D. Kusnida (2010)- Paleocene postgenetic accumulation of nannoplankton on the phillipsite minerals in Roo Rise, Indian Ocean. Jurnal Geologi Indonesia 5, 1, p. 49-56.
(online at: www.bgl.esdm.go.id/publication/index.php/dir/article_detail/264)
(Paleocene nannoplankton on phillipsite crystals in core from Roo Rise, 3880- 3914m below sea level)

Adisaputra, Mimin K. & L. Sarmili (1995)- Neogene events through biostratigraphic constrain in the Banda Sea. In: S. Nishimura & R. Tsuchi (eds.) Proc. Oji Seminar on Neogene evolution of Pacific Ocean Gateways, Kyoto, IGCP-355, p. 72-79.

Ashton, P.S., R.J. Morley, J. Heckenhauer & V. Prasad (2021)- The magnificent Dipterocarps: precis for an Epitaph? Kew Bulletin 76, p. 87-125.
(Magnificent dipterocarps, valued for camphor-bearing resin, once dominated lowland tropical rainforests of SE Asia, but overharvesting threatens their species with extinction. Group believed to have migrated into SE Asia from Gondwana in Eocene time)

Ayala, E.O. (1980)- Chronostratigraphic studies of Philippine *Cycloclypeus*. The Philippine Geologist (J. Geological Society of the Philippines) 34, 1, p.
(Cycloclypeus larger foram assemblages common in Miocene carbonates of Philippines. Mainly Cycloclypeus carpenteri)

Baggio, D. & S. Sartori (1996)- The Lyllean coefficients checked for Tethyan and Indonesian Cenozoic benthic molluscs; a comparison with the Mesozoic. In: Reports of Shallow Tethys 4, Albrechtsburg 1994, Annali Musei Civico Rovereto 11, Supplement, p. 333-356.
(Lyell's survivorship percentages of Cenozoic mollusc assemblages re-calculated for Indonesian area)

Bakx, L.A.J. (1932)- De genera *Fasciolites* en *Neoalveolina* in het Pacifische gebied. Verhandelingen Geologisch-Mijnbouwkundig Genootschap Nederland Kolonien, Geologische Serie 9, p. 205-266.
('The genera Fasciolites and Neoalveolina in the Pacific area'. Eocene Alveolina spp.and Oligocene Borelis pygmaeus (at that time all assigned to genus Fasciolites), from Timor, Borneo, Java, New Guinea, Sulawesi)

Bande, M.B. & U. Prakash (1986)- The Tertiary flora of Southeast Asia with remarks on its palaeoenvironment and phytogeography of the Indo-Malayan region. Review Palaeobotany Palynology 49, p. 203-233.
(Compilation of information on Paleogene and Neogene flora of SE Asia, with reconstruction of Tertiary environments of region. SE Asia flora compared with that of India and migration of various families and genera between these two areas is traced)

Banner, F.T. & J. Highton (1989)- On *Pseudotaberina malabarica* (Carter) (Foraminiferida). J. Micropalaeontology 8, 1, p. 113-129.
(online at: https://www.j-micropalaeontol.net/8/113/1989/jm-8-113-1989.pdf)
(On Archaias-type soritid species formerly assigned to Orbitolites. In SE Asia characterizes carbonate platform facies of zone Tf1 (late E Miocene). With meandrine, involute chambers in large microspheric specimens. May be same species as Archaias vandervlerki De Neve from E Kalimantan)

Banner, F.T. & R.L. Hodgkinson (1991)- A revision of the foraminiferal subfamily Heterostegininae. Revista Espanola Micropaleontologia 23, 2, p. 101-140.
(New genus names Tansinhokella for Eocene members of Spiroclypeus group and Vlerkina for involute Heterostegina)

Banner, F.T. & M.A. Samuel (1995)- *Alanlordia*, a new genus of acervuline foraminifera from the Neogene of Indonesia. J. Micropalaeontology 14, p. 107-117.
(online at: https://www.j-micropalaeontol.net/14/107/1995/jm-14-107-1995.pdf)
(Neogene limestones of Nias and Tuangku, W of Sumatra, with new acervulinid forma Alanlordia niasensis niasensis n.gen., n.sp., in Late Pliocene (looks like Maastrichtian Vanderbeekia of Middle East). Serravallian limestones contain ancestral A. niasensis primitiva, n. subsp., and deeper marine and simpler A. banyakensis n. sp. (looks like Eocene Wilfordia))

Bannink, D.D. (1950)- Een monografie van het genus *Operculina* d'Orbigny, 1826. Ph.D. Thesis University of Leiden, p. 1-159. *(Unpublished)*
(Monograph on foram genus Operculina, including descriptions of Tertiary material from Borneo and Sumatra. Genus comprises nine species)

Barre-De Cruz, C. (1982)- Etude palynologique du Tertiaire de Sud-Est Asiatique (Kalimantan: delta de la Mahakam, Mer de Chine: Permis de Beibu). Thesis Universite de Bordeaux III, vols. 1, 161p. and vol. 2, 61p.
('Palynological study of the Tertiary of SE Asia (Kalimantan/Mahakam Delta and S China Sea/ Beibu Permit))

Basov, I.A. & V.A. Krashenninnikov (1995)- Stratigraphy and foraminifera of Pliocene-Quarternary deposits of the Timor Trough. Izdatelstvo "Nauchniy Mir", Moscow, p. 1-110.

Bassi, D., M. Aftabuzzaman, M. Bolivar-Feriche, J.C. Braga, J. Aguirre, W. Renema, H. Takayanagi & Y. Iryu (2021)- Biostratigraphical and palaeobiogeographical patterns of the larger porcelaneous foraminifer *Austrotrillina* Parr, 1942. Marine Micropaleontology 169, 102058, p. 1-18.
(Review of Tethyan larger porcelaneous foram Austrotrillina, widespread in M-L Eocene- M Miocene of W Tethys and Indo-Pacific. Four species (A. brunni, A. eocaenica, A. howchini, A. striata). A. asmariensis Adams junior synonym of A. brunni Marie. A. eocaenica first appears in M-L Eocene of Iran. Rupelian descendants A. brunni and A. striata migrated from W Tethys into Indo-Pacific, reaching Indonesia and W Australia in Chattian. From Burdigalian, A. howchini only in IndoPacific, possibly due to closing of Tethyan Seaway)

Bassi, D., M. Bolivar-Feriche, W. Renema, J.C. Braga, J. Pignatti, G. Di Domenico, K. Fujita, J.H. Lipps, J. Reolid & Y. Iryu (2022)- Larger porcelaneous foraminifera with a common ancestor: the Neogene Indo-Pacific *Flosculinella* and *Alveolinella* (Alveolinoidea). Marine Micropaleontology 173, 102124, p. 1-19.
(Thorough review of Miocene-Recent complex Indo-Pacific alveolinids. Borelis philippinensis is common ancestor of Indo-Pacific late Oligocene- M Miocene Flosculinella and M Miocene- Recent Alveolinella. Speciation occurred in E-M Miocene in Coral Triangle of SE Asia. In E Kalimantan, Alveolinella borneensis appeared in E Tortonian, A. quoyi appeared in Late Tortonian)

Baumann, P. (1972)- Les faunes de foraminiferes de l'Eocene superieur a la base du Miocene dans le basin de Pasir, Sud de Kalimantan. Revue Institut Francais Petrole 27, 6, p. 817-829.
('The foraminifera of the Late Eocene to the base of the Miocene in the Pasir Basin, S Kalimantan.' Good documentation of Late Eocene- Oligocene planktonic foraminifera zonation in Pasir Basin, SE Kalimantan)

Beets, C. (1942)- Mollusken aus dem Tertiar des Ostindischen Archipels. Leidsche Geologische Mededelingen 13, 1, p. 218-254.
(online at: https://repository.naturalis.nl/pub/505999/LGM1942013001013.pdf)
('Molluscs from the Tertiary of the East Indies Archipelago'. Three short papers on molluscs from collections in The Netherlands: (1) gastropod Buccinum in E Indies, (2) Notes on some interesting molluscs from E Indies, (3) Observations on small Neogene mollusc fauna from E Kalimantan (collected by Rutten))

Beets, C (1943)- Die Gattung *Galeodea* Link im Tertiar von Insulinde. Verhandelingen Geologisch-Mijnbouwkundig Genootschap Nederland Kolonien, Geologische Serie 13, p. 435-443.
('The genus Galeodea (gastropod) in the Tertiary of Indonesia')

Beets, C. (1949)- On the occurrence of *Biplanispira* in the uppermost Eocene (Kyet-U-Bok Band) of Burma. Geologie en Mijnbouw, N.S., 11, 7, p. 229-232.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0WFlhSElBcHp0ZFE/view)
(Late Eocene Biplanispira miriabilis in thin section in British Museum from Kyet-u-bok limestone band between shaly U Eocene Yaw stage and Oligo-Miocene Pegu System. Associated with Nummulits and Discocyclina)

Beets, C. (1950)- Revised determinations of East Indian and related fossil mollusca. Verhandelingen Kon. Nederlands Geologisch Mijnbouwkundig Genootschap, Geologische Serie 15, 2, p. 329-341.

Beets, C. (1950)- On an East-Indian representative of the rare gastropod genus *Trochocerithium*. Verhandelingen Kon. Nederlands Geologisch Mijnbouwkundig Genootschap, Geologische Serie 15, 2, p. 342-348.
(Incl. Trochocerithium gloriamaris n.sp.)

Beets, C. (1950)- On fossil brachyuran crabs from the East Indies. Verhandelingen Kon. Nederlands Geologisch Mijnbouwkundig Genootschap, Geologische Serie 15, 2, p. 349-354.
(Three species of Neogene fossil crabs from Java and Madura)

Beets, C. (1986)- Notes on *Buccinulum* (Gastropoda, Buccinidae), a reappraisal. Scripta Geologica 82, p. 83-100.
(online at: http://repository.naturalis.nl/document/148781)
(Review of Eocene- Recent gastropod genus Buccinulum in Indo- West Pacific region, mainly from W Indonesia. New subgenus Samudra proposed, with type species B. djocdjocartae)

Belford, D.J. (1982)- Redescription of *Miogypsina neodispansa* (Jones & Chapman), Foraminiferida, Christmas Island, Indian Ocean. BMR J. Australian Geology Geophysics 7, 4, p. 315-320.
(online at: https://d28rz98at9flks.cloudfront.net/81127/Jou1982_v7_n4_p315.pdf)
(Miogypsina neodispansa (Jones & Chapman, 1900) s redescribed from Christmas Island type locality, Indian Ocean. It is referred to subgenus Lepidosemicyclina and is senior synonym of M. droogeri Mohan & Tewari. Age probably Letter zone Tf1, late Early Miocene)

Beu, A.G. (2005)- Neogene fossil tonnoidean gastropods of Indonesia. Scripta Geologica 130, p. 1-186.
(online at: https://repository.naturalis.nl/pub/210121)
(Review of Neogene gastropods from Java, Nias, etc. in Naturalis, Leiden collections)

Bhaumik, A.K., A.K. Gupta, M.S. Raj, K. Mohan, S. De & S. Sarkar (2007)- Paleoceanographic evolution of the northeastern Indian Ocean during the Miocene: evidence from deep-sea foraminifera (DSDP Hole 216A). Indian J. Marine Science 36, 4, p. 332-341.
(online at: http://nopr.niscair.res.in/bitstream/123456789/57/1/IJMS%2036(4)%20(2007)%20332-341.pdf)
(Deepwater benthic forams at ODP Site 216A suggest shift in deep-sea ventilation at 15-14 Ma, coinciding with M Miocene cooling)

Bidgood, M.D., M.D. Simmons & C.G. Thomas (2000)- Agglutinated foraminifera from Miocene sediments of northwest Borneo. In: M.B. Hart et al. (eds.) Proc. 5th International Workshop on Agglutinated Foraminifera, Plymouth 1997, Grzybowski Foundation, Krakow, Special Publ. 7, p. 41-58.
(online at: http://gf.tmsoc.org/Documents/IWAF-5/Bidgood+Simmons+Thomas-IWAF5-1997.pdf)
(Miocene sediments from Brunei and Sarawak often rich in agglutinated forams. Assemblages characteristic of different depositional environments, including Distal turbidite (often deformed 'Trochammina/ Recurvoides'), Proximal turbidite (fine-grained and thin-walled Trochammina, Cyclammina, Haplophragmoides), Tidal flat/

tidal channel (coarse-grained Trochammina, Recurvoides and occasional Ammobaculites), Lagoon/ distributary channel margin (mangrove swamps with Trematophragmoides, Miliammina fusca), etc.)

Billman, H.G., L. Hottinger & H. Oesterle (1980)- Neogene to Recent rotaliid foraminifera from the Indo-Pacific Ocean; their canal system, their classification and their stratigraphic use. Schweizerische Palaontologische Abhandlungen 101, p. 71-113.
(Species and zonation of rotalid foraminifera successfully used for biozonation of E Kalimantan deltaic series. New species incl. Challengerella brady, Challengerella persica, Asterorotalia gaimardii inermis, Pseudorotalia schroeteriana angusta)

Billman, H.G. & M.E. Scrutton (1976)- Stratigraphic correlation in Indonesia. In: Offshore South East Asia Conference 1976, Singapore, SE Asia Petroleum Exploration Society (SEAPEX), Paper 9, p. 1-13.
(Elegant overview of Cenozoic biostratigraphic zonations in Indonesia)

Blow, W.H. (1969)- Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy. Proc. First Int. Conf. Planktonic Microfossils, Geneva 1967, 1, Brill, Leiden, p. 199-422.
(Classic text on Eocene- Recent planktonic foraminifera zonations, using the N and P-numbered zones, widely used in Indonesia. Parts of this work are based on classic Indonesian sections like Bojonegoro 1 well, Java, etc.)

Blow, W.H. (1979)- The Cainozoic Globigerinida. A study of the morphology, taxonomy, evolutionary relationships and the stratigraphical distribution of some Globigerinida. Brill, Leiden, 3 vols., p. 1-1413.
(Monumental study of Tertiary- Recent planktonic foraminifera (expansion of Blow (1969) book))

Bolli, H.M. & J.B. Saunders (1985)- Oligocene to Holocene low latitude planktic foraminifera. In: H.M. Bolli, J.B. Saunders & K. Perch-Nielsen (eds.) Plankton Stratigraphy, Cambridge University Press, Cambridge, p. 155-262.
(Comprehensive review of Oligocene- Recent planktonic foraminifera and zonations)

Boltovskoy, E. (1974)- Neogene planktonic foraminifera of the Indian Ocean (DSDP Leg 26). In: T. Davies & B.P. Luyendyk (eds.) Initial Reports of Deep Sea Drilling Project 26, Washington, p. 675-741.
(online at: www.deepseadrilling.org/26/volume/dsdp26_30.pdf)
(Neogene planktonic foraminifera from Leg 26 sites in Indian Ocean between W Australia and Madagascar)

Boudagher-Fadel, M.K. (2008)- Chapter 6- The Cenozoic larger benthic foraminifera: The Palaeogene. In: Evolution and geological significance of larger benthic foraminifera. Developments in Palaeontology and Stratigraphy 21, Elsevier, p. 297-418.

Boudagher-Fadel, M.K. (2008)- Chapter 7- The Cenozoic larger benthic foraminifera: The Neogene. In: Evolution and geological significance of larger benthic foraminifera. Developments in Palaeontology and Stratigraphy 21, Elsevier, p. 419-548.

Boudagher-Fadel, M.K. (2015)- Chapter 5- The Cenozoic planktonic foraminifera: The Paleogene. In: Biostratigraphic and geological significance of planktonic foraminifera, updated 2nd Edition, UCL Press (University College London), p. 141-201.
(online at: http://discovery.ucl.ac.uk/1404017/3/9781910634264_OnlinePDF.pdf)

Boudagher-Fadel, M.K. (2015)- Chapter 6- The Cenozoic planktonic foraminifera: The Neogene. In Biostratigraphic and geological significance of planktonic foraminifera, updated 2nd Edition, UCLPress (University College London), p. 203-269.
(online at: http://discovery.ucl.ac.uk/1404017/3/9781910634264_OnlinePDF.pdf)
(Elegant review of Miocene- Holocene planktonic foraminifera, with many of the species illustrated from Indonesia)

Boudagher-Fadel, M.K. & F.T. Banner (1997)- The revision of some genus-group names in Tethyan Lepidocyclininae. Paleopelagos, 7, p. 3-16.

Boudagher-Fadel, M.K. & F.T. Banner (1999)- Revision of the stratigraphic significance of the Oligocene-Miocene "Letter-Stages". Revue de Micropaleontologie 42, p. 93-97.
(online at: https://www.researchgate.net/publication/240915410_Revision_of_the_stratigraphic_significance_of_the_Oligocene-Miocene_'Letter-Stages')
(Review of the Oligo-Miocene part of the classic Indo-Pacific larger foram Letter zonation of Van der Vlerk and Umbgrove (1927) and others. 'New' correlation between Far East Letter Stages and Oligo-Miocene planktonic foram stages)

Boudagher-Fadel, M.K. & A.R. Lord (2000)- The evolution of *Lepidocyclina (L.) isolepidinoides, L. (Nephrolepidina) nephrolepidinoides, L. (N.) brouweri* in the Late Oligocene-Miocene of the Far East. J. Foraminiferal Research 30, 1, p. 71-76.
(online at: https://www.researchgate.net/publication/250084155_The_evolution_of_Lepidocyclina_L_Isolepidinoides_L_Nephrolepidina_Nephrolepidinoides_sp_nov_L_N_Brouweri_and_L_N_Ferreroi_ Etc)
(Re-description of the well-known evolution from Lepidocyclina (L) to Lepidocyclina (N) around the Oligo-Miocene boundary in material from NE and SE Kalimantan and Nias, off Sumatra)

Boudagher-Fadel, M.K. & G.D. Price (2010)- Evolution and paleogeographic distribution of the lepidocyclinids. J. Foraminiferal Research 40, 1, p. 79-108.
(online at: https://www.academia.edu/6772371/EVOLUTION_AND_PALEOGEOGRAPHIC_DISTRIBUTION_OF_THE_LEPIDOCYCLINIDS)
(Review of Eocene- Miocene lepidocyclinid foraminifera in the American, Mediterranean and Indo-Pacific bioprovinces. Earliest lepidocyclinids in Indo-Pacific Province migrated from Mediterranean in end Rupelian)

Boudagher-Fadel, M.K. & G.D. Price (2013)- The phylogenetic and palaeogeographic evolution of the miogypsinid larger benthic foraminifera. J. Geological Society, London, 170, p. 185-208.
(online at: http://discovery.ucl.ac.uk/1383038/)
(Extensive review of Oligocene- Miocene miogypsinid foraminifera, mainly in Indo-Pacific and Mediterranean provinces)

Boudagher-Fadel, M.K. & G.D. Price (2014)- The phylogenetic and palaeogeographic evolution of the nummulitoid larger benthic foraminifera. Micropalaeontology 60, 6, p. 483-508.
(online at: http://discovery.ucl.ac.uk/1426612/)

Boudagher-Fadel, M.K. & G.D. Price (2021)- The geographic, environmental and phylogenetic evolution of the Alveolinoidea from the Cretaceous to the present day. UCL Open Environment, 2021, 2, 03, p. 1-34.
(online at: https://journals.uclpress.co.uk/ucloe/article/id/440/)
(Alveolinoids larger forger foraminiferae found globally from Cretaceous to Recent. Exhibit provincialism with characteristic genera often confined to one of American, Tethyan or Indo-Pacific provinces. Alveolinoids originated in Cretaceous in Tethyan province, but went extinct at K-T boundaty. Late Paleocene alveolinoid forms migrated E-ward from Americas to Tethys, and from there to Indo-Pacific province. Alveolinoids still exist today (cosmopolitan Borelis and Indo-Pacific Alveolinella). Indo-Pacific alveolinids range from latest Ypresian (P9) to Bartonian (end M Eocene extinction event at 37.8Ma; same time as extinction of large Nummulites and Assilina). In Neogene of Far East evolution of Borelis-Flosculinella- Alveolinella lineage)

Bramlette, M.N. & W.R. Riedel (1954)- Stratigraphic value of discoasters and some other microfossils related to Recent coccolithophores. Journal of Paleontology 28, p. 385-403.
(Incl. emended re-description of Discoaster brouweri Tan Sin Hok 1927 and new Late Eocene species Discoaster tani and D. tani nodifer)

Bromfield, K. (2013)- Neogene corals from the Indo-Pacific, Indonesia, Papua New Guinea and Fiji. Bull. American Paleontology 387, p. 1-130.

(Descriptions of 155 M Miocene- E Pleistocene coral species (22 new, extinct) from (1) Salayar Lst in S Sulawesi, Yalam Lst in New Britain, PNG, and (3) Tokelau Lst, Fiji)

Bronnimann, P. & J. Resig (1971)- A Neogene Globigerinacean biochronologic time-scale of the Southwestern Pacific. Initial Reports Deep Sea Drilling Project (DSDP) 7, 2, p. 1235-1469.
(online at: http://deepseadrilling.org/07/volume/dsdp07pt2_28.pdf)
(Extensively documented M Miocene- Recent planktonic foram zonation of DSDP holes of Ontong Java Plateua and East Caroline Basin N of PNG, SW Pacific)

Burckle, L.H. (1978)- Early Miocene to Pliocene diatom datum levels for the equatorial Pacific. In: Biostratigraphic datum-planes of the Pacific Neogene, IGCP Project 114, Proc. Second Working group meeting, 1977, Geological Research Development Centre (GRDC), Bandung, Special Publ. 1, p. 25-44.
(Over 40 Early Miocene- Pliocene diatom datum levels in Equatorial Pacific)

Chandra, B.Y. (2010)- Characterization of clastic sediment: a palynofacies approach. Proc. 34th Annual Conv. Indonesian Petroleum Association, Jakarta, IPA10-G-075, p. 1-7.
(General paper describing palynofacies as tool for identification of depositional environment and sequences)

Chaproniere, G.C.H. (1975)- Palaeoecology of Oligo-Miocene larger foraminiferida, Australia. Alcheringa 1, p. 37-58.
(Eight large foram assemblages distinguished in E Miocene of W Australia North-West Cape area, each representing specific environment)

Chaproniere, G.C.H. (1980)- Biometrical studies of Early Neogene larger foraminiferida from Australia and New Zealand. Alcheringa 4, p. 153-181.
(Statistical study of latest Oligocene- M Miocene Cycloclypeus, Heterostegina, Miogypsina and Lepidocyclina from N Australia, New Zealand)

Chaproniere, G.C.H. (1980)- Influence of plate tectonics on the distribution of Late Palaeogene to Early Neogene larger foraminiferids in the Australasian region. Palaeogeogr. Palaeoclim. Palaeoecology 31, p. 299-317.
(Paleogeographic distribution of Oligocene-Miocene larger forams in SE Asia- Australia. Cycloclypeus, Miogypsina and Lepidocyclina invaded Australasian region in M Oligocene)

Chaproniere, G.C.H. (1981)- Australasian mid-Tertiary larger foraminiferal associations and their bearing on the East Indian Letter Classification. BMR J. Australian Geology Geophysics 6, p. 145-151.
(online at: https://d28rz98at9flks.cloudfront.net/81070/Jou1981_v6_n2_p145.pdf)
(Eight latest Oligocene- M Miocene larger foram associations in NW Cape Range, off Exmouth Gulf, NW Australia (LF1-LF8; Te- Lower Tf). No Spiroclypeus or Eulepidina observed (but Heterostegina borneensis in LF2. LF1 assemblage with Lacazinella, probably reworked from underlying Eocene Giralia Limestone)

Chaproniere, G.C.H. (1983)- Tertiary larger foraminiferids from the northwestern margin of the Queensland Plateau, Australia. Paleontological papers 1983, Bureau Mineral Resources Geology Geophysics, Bull. 217, p. 31-57.
(online at: https://d28rz98at9flks.cloudfront.net/12/Bull_217.pdf)
(M Eocene Ta3 with Asterocyclina and Latest Oligocene- E Miocene (around Lower Te/ upper Te boundary) larger foram assemblages in dredge samples from~1500-2500m water depth on Queensland Plateau, off Great Barrier Reef, NE Australia)

Chaproniere, G.C.H. (1984)- Oligocene and Miocene larger foraminiferida from Australia and New Zealand. Bureau Mineral Resources Geology Geophysics, Canberra, Bull. 188, p. 1-98.
(online at: https://www.ga.gov.au/products/servlet/controller?event=GEOCAT_DETAILS&catno=15)
(Larger foraminifera from Late Oligocene-M Miocene outcrops in Carnarvon Basin and W Australia; also Ashmore Reef 1 well in Bonaparte Gulf, Gage Roads 2 well in Perth Basin; Batesford and Bochara Lst in

Victoria; Wreck Island 1 well in Queensland; and various localities in New Zealand. Two new subspecies of Lepidocyclina (N): Lepidocyclina (N) howchini praehowchini and Lepidocyclina (N) orakeiensis waikukuensis)

Chaproniere, G.C.H. (1984)- The Neogene larger foraminiferal sequence in the Australian and New Zealand regions and its relevance to the East Indies Letter Stage classification. Palaeogeogr. Palaeoclim. Palaeoecology 46, p. 25-35.
(Neogene larger foraminifera in N Australia range from Late Oligocene-earliest M Miocene, in S Australia restricted to late E Miocene. Longer range in New Zealand: Late Oligocene- mid-M Miocene)

Chaproniere, G.C.H. (1992)- The distribution and development of Late Oligocene and Early Miocene reticulate globigerines in Australia. Marine Micropaleontology 18, p. 279-305.
(Globigerinoides represented by both hispid (Gs. primordius) and reticulate (Gs. quadrilobatus) forms at levels below first appearance of Globoquadrina dehiscens (within Zone N4A in N Australia). Globigerinoides quadrilobatus warm, shallow water dweller. Globigerina bulloides group temperate biogeographic range, but typical of upwelling zones in subtropical and tropical areas)

Cheng , Z., J. Wu, C. Luo, Z. Liu, E. Huang, H. Zhao, L. Dai & C. Weng (2023)- Coexistence of savanna and rainforest on the ice-age Sunda Shelf revealed by pollen records from southern South China Sea. Quaternary Science Reviews 301, 107947, p.
(Marine pollen records from southern South China Sea focusing on Late Pleistocene Last Glacial Maximum (LGM) time show that sites in front of Sunda Shelf paleo-rivers contain substantially more Poaceae (grass) pollen than site offshore North Borneo. Co-occurrence of grass and rainforest pollen suggests longitudinal climate gradient with increasing aridity toward interior of glacial Sundaland. In Borneo rainforest persisted during glacial periods, but vegetation records support hypothesis of a 'savanna corridor' over LGM Sundaland (see also Morley & Morley 2022, Hamilton et al. 2023))

Clarke, W.J. & W.H. Blow (1969)- The inter-relationship of some Late Eocene, Oligocene and Miocene larger foraminifera and planktonic biostratigraphic indices. In: P. Bronnimann & H.H. Renz (eds.) Proc. First Int. Conference on Planktonic Microfossils, Geneva 1967, Brill, Leiden, 2, p. 82-97.
(Key paper from the illustrious 1960s BP paleo lab. One of first papers calibrating the Eocene- Recent larger foraminifera Letter 'Letter Stages' with the planktonic foraminifera zonation (generally difficult, as they are rarely found together due to different facies). Includes chapter on occurrences of Miogypsinidae in Indonesia (W Java, Sumatra))

Cole, W.S. (1939)- Large foraminifera from Guam. Journal of Paleontology 13, 2, p. 183-189.
(Limestones from Guam with larger foraminifera, Spiroclypeus higginsi n. sp., primitive Miogypsinoides dehaartii var. formosensis and Lepidocyclina (N.) parva, Borelis pygmaeus. Age basal Miocene, zone Lower Te (more likely latest Oligocene?; JTvG))

Cole, W.S. (1945)- Larger foraminifera of Lau, Fiji. In: H.S. Ladd & J.E. Hoffmeister (eds.) Geology of Lau, Fiji. Bernice P. Bishop Museum Bull. 181, p. 272-297.

Cole, W.S. (1950)- Larger foraminifera from the Palau Islands. U.S. Geological Survey (USGS) Professional Paper 221-B, p. 21-31.
(online at: http://pubs.usgs.gov/pp/0221b/report.pdf)
(Description of larger foramifera from Palau islands, Micronesia, SW Pacific: Late Eocene (Tab) with Pellatispira and M Miocene (Tf) with Katacycloclypeus, Lepidocyclina rutteni, Lepidocyclina palauensis n.sp. (latter re-assigned to Lepidocyclina radiata by Cole (1963))

Cole, W.S. (1954)- Larger Foraminifera and smaller diagnostic Foraminifera from Bikini drill holes. U.S. Geological Survey (USGS) Professional Paper, 260-O, p. 569-608.
(online at: http://pubs.usgs.gov/pp/0260m/report.pdf)
(Description of 37 Oligocene- Recent foram species from two wells (2556') of Bikini Atoll)

Cole, W.S. (1957)- Larger foraminifera from Eniwetok Atoll drill holes. U.S. Geological Survey (USGS) Professional Paper, 260-V, p. 743-784.
(online at: http://pubs.usgs.gov/pp/0260v/report.pdf
(62 species of Late Eocene (Asterocyclina, Nummulites, Biplanispira, Pellatispira), Late Oligocene (Eulepidina, Heterostegina borneensis, Borelis, Miogypsinoides, Spiroclypeus), Miocene (Miogypsina, Flosculinella) and Pliocene- Recent (Calcarina, Marginopora, Sorites) larger foraminifera from three Eniwetok Atoll drill holes. Deeper water genus Cycloclypeus rare, suggesting continuous shallow marine facies)

Cole, W.S. (1957)- Geology of Saipan, Mariana Islands, Part 3 Paleontology, Larger Foraminifera. U.S. Geological Survey (USGS) Professional Paper 280-I, p. 321-360.
(online at: http://pubs.usgs.gov/pp/0280e-j/report.pdf)
(Larger foram assemblages for Saipan island: Late Eocene Tb (20 species; Pellatispira, Nummulites, Asterocyclina, etc.), Late Oligocene- E Miocene Te (35 species; incl. Miogypsinoides, Heterostegina borneensis) and Pleistocene (7 species))

Cole, W.S. (1960)- Upper Eocene and Oligocene larger foraminifera from Viti Levu, Fiji. U.S. Geological Survey (USGS) Professional Paper 374-A, p. 1-7.
(online at: http://pubs.usgs.gov/pp/0374a/report.pdf)
(Descriptions of Late Eocene (Nummullites, Pellatispira, Spiroclypeus, Discocyclina, Asterocyclina) and Early Oligocene (Nummulites fichteli, Gypsina discus) larger foraminifera from main island of Fiji)

Cole, W.S. (1960)- Problems of the geographic and stratigraphic distribution of certain Tertiary larger foraminifera. In: Hanzawa Memorial Volume, Science Reports Tohoku University, ser. 2 (Geology), Special Volume 4, p. 9-18.

Cole, W.S. (1963)- Analysis of *Lepidocyclina radiata* (Martin). Bull. American Paleontology 46, 208, p. 157-185.
(Study of Lepidocyclina from Tf (Miocene) of Futuna Lst of Lau, Fiji Islands, assigned to Lepidocyclina radiata)

Cole, W.S. (1963)- Tertiary larger foraminifera from Guam. U.S. Geological Survey (USGS) Professional Paper, 403-E, p. 1-28.
(online at: http://pubs.usgs.gov/pp/0403e/report.pdf)
(Late Eocene (Asterocyclina, Nummulites, Pellatispira, Biplanispira, Halkyardia), Oligocene (Nummulites fichteli), E-M Miocene (Miogypsinoides dehaartii, Katacycloclypeus annulatus) and Pleistocene (Calcarina, Baculogypsina, Cycloclypeus carpenteri) larger forams from outcrops on Guam)

Cole, W.S. (1969)- Larger foraminifera from deep sea drill holes on Midway Atoll. U.S. Geological Survey (USGS) Professional Paper, 680-C, p. 1-15.
(online at: http://pubs.usgs.gov/pp/0680c/report.pdf)
(Early Miocene Te larger forams (Miogypsinoides dehaartii, Spiroclypeus, Austrotrillina striata) in deeper part of 1261' deep well)

Cole, W.S. (1970)- Larger foraminifera of Late Eocene age from Eua, Tonga. U.S. Geological Survey (USGS) Professional Paper, 640-B, p. 1-17.
(online at: http://pubs.usgs.gov/pp/0640b/report.pdf)
(Eocene sample with 8 species of larger foraminifera characteristic of Upper Eocene (Tb) from E side of Eua Island, Tonga. Incl. Pellatispira madaraszi, Discocyclina omphala, Nummulites pengaronensis, Asterocyclina matanzensis, Spiroclypeus vermicularis, Heterostegina saipanensis, 'Biplanispira' fulgeria)

Cole, W.S. (1975)- Concordant age determinations by larger and planktonic foraminifera in the Tertiary of the Indo-Pacific region. J. Foraminiferal Research 5, p. 21-39.
(online at: http://jfr.geoscienceworld.org/content/5/1/21.full.pdf)

(Good agreement in age determinations based on larger and planktoic foraminifera from 5 widely separated localities: (1) Sentolo Fm, C Java late E Miocene Lower Tf LBF and Globigerinoides sicanus-Globigerinatella insueta (N8) planktonic foram zone; (2,3) Larat (Moluccas) and Solomon Islands: early E Miocene Te LBF zone and Globigerinita dissimilis zone planktonics, etc.)

Cole, W.S. & J. Bridge (1953)- Geology and larger Foraminifera of Saipan Island. U.S. Geological Survey (USGS) Professional Paper 253, p. 1-45.
(online at: http://pubs.usgs.gov/pp/0253/report.pdf)
(38 species of larger foraminifera from limestone outcrops on Saipan. Most species are Early Miocene forms, but 4 samples contain Eocene fauna and 3 have Plio-Pleistocene faunas)

Conesa, G.A.R., E. Favre, P. Munch, H. Dalmasso & C. Chaix (2005)- Biosedimentary and paleoenvironmental evolution of the Southern Marion Platform from the Middle to Late Miocene (northeast Australia, ODP Leg 194, Sites 1196 and 1199). In: F.S. Anselmetti et al. (eds.) Proc. Ocean Drilling Program (ODP), Scientific Results, 194, p. 1-38.
(Online at: www-odp.tamu.edu/ publications/194_SR/)

Cook, P.L. & R. Lagaaij (1976)- Some Tertiary and Recent conescharelliniform bryozoa. Bull. British Museum (Natural History), Zoology, 29, p. 317-376.
(Includes descriptions of Miocene bryozoa from Madura (3 new species of Lacrimula, L. asymmetrica, L. similis, L. grunaui, from E Miocene Globigerinatella insueta Zone)

Cosijn, A.J. (1938)- Statistical studies on the phylogeny of some foraminifera: *Cycloclypeus* and *Lepidocyclina* from Spain, *Globorotalia* from the East Indies. Doct. Thesis Technical University Delft, p. 1-66.
(online at: https://repository.tudelft.nl/islandora/object/uuid%3A75768866-49a4-43b7-9279-2521a01ffb28
(Not-overly-useful? sets of measurements on Cycloclypeus from Spain and Mio-Pliocene Globorotalia menardii and Gr. tumida from BPM Bojonegoro 1 well, E Java. Thesis supervised by J.H.F. Umbgrove)

Crespin, I. (1956)- Papers on Tertiary micropalaeontology. Bureau Mineral Resources Geology Geophysics, Canberra, Report 25, p. 1-77.
(online at: www.ga.gov.au/products-services/legacy-publications/reports.html)
(Seven papers on foraminifera from Australia, Phillippines, PNG, Bougainville)

Cui, X., Xin Li, J.C. Aitchison & H. Luo (2022)- Paleobiogeographic significance of Tethyan Lower Cretaceous *Cecrops septemporatus* Zone radiolarian faunas. Palaeogeogr. Palaeoclim. Palaeoecology 589, 110836, p.
(Valanginian to lower Hauterivian Cecrops septemporatus Zone radiolaria assemblages widespread across low-latitude Tethys, from Italy to Gonggui section in S Tibet (Purang ophiolitic melange in W part of Yarlung Tsangpo suture zone = likely distal parts of N India continental margin) to W-Pacific (in Indonesia known from Barru Complex of S Sulawesi; Munasri, 2013)

Cui, X., X. Li, J.C. Aitchison & H. Luo (2022)- Early Cretaceous monsoonal upwelling along the northern margin of the Gondwana continent: Evidence from radiolarian cherts. Marine Micropaleontology 181,1, 102247, p.
(Radiolarian-bearing bedded cherts exposed in volcanic-sedimentary sequences of Purang ophiolitic complex in W Yarlung Tsangpo suture zone, Tibet. Diverse Lower Cretaceous (Valanginian–Hauterivian) radiolarian assemblages of Cecrops septemporatus Zone, with 55-five species, including Archaeodictyomitra apiarium, Cecrops septemporatus, Hemicryptocapsa capita Tan, Pseudodictyomitra carpatica and Svinitzium pseudopuga. Geochemistry of cherts indicative of deposition in oceanic basin close to continental margin (in this case N margin of Indian continent, with enhanced upwelling activity(

De Beaufort, L.F. (1931)- Pisces, Reptilia and Aves. In: B.G. Escher et al. (eds.) De palaeontologie en stratigraphie van Nederlandsch Oost-Indie, Leidsche Geologische Mededelingen 5 (K. Martin memorial volume), p. 461-470.

(online at: www.repository.naturalis.nl/document/549718)
(Summary of fossil fish, reptiles and birds known from Indonesia as of 1931. No figures)

De Bock, J.F. (1973)- Embryonal structures of *Miogypsina*. Scripta Geologica, Leiden, 18, p. 1-15.
(online at: https://repository.naturalis.nl/pub/317395/SG1973018001.pdf)
(Detailed study of embryonal features of well-preserved bispiral E-M Miocene Miogypsina, based on material from NE Borneo, E Java and Madura)

De Bock, J.F. (1976)- Studies on some *Miogypsinoides-Miogypsina* s.s. associations with special reference to morphological features. Scripta Geologica, Leiden, 36, p. 1-137.
(online at: https://repository.naturalis.nl/pub/317461/SG1976036001.pdf)
(Detailed study of well-preserved Miocene Miogypsina and Miogypsinoides, partly based on material from Madura and Larat (Kai islands))

Dekkers, A.M., V. Liverani, S. Coric, S.J. Maxwell, & B.M. Landau (2020)- A new genus for Indo-Pacific fossil strombids, and two new species from the Miocene of Java and Borneo (Caenogastropoda, Strombidae). Basteria 84, p. 1-9.
(online at: https://natuurtijdschriften.nl/pub/1001530/BAST2020084001001.pdf)
(New Strombus species from early M Miocene of Wonosari, Gunung Kidul, C Java, in Natural History Museum, Vienna, collections and from E Kalimantan)

Dollfus, G.F. (1908)- Sur quelques polypiers fossiles des Indes neerlandaises. Jaarboek Mijnwezen Nederlandsch Oost-Indie 37 (Verbeek Moluccas Report), p. 676-686.
('On some fossil corals from the Netherlands Indies'. Brief description of five Late Tertiary corals collected by Verbeek from C Timor, E Seram and Daweloo island near Babar)

Donovan, S.K. & R.A. Helwerda (2017)- Neogene crinoids of southeast Asia: preservation, systematics and significance. Alcheringa 41, 2, p. 215-221.
(Cenozoic record of fossil crinoids is poor. Two new, small crinoid faunas reported of moderate diversity from Pliocene of E Kalimantan and Philippines. With Metacrinus?, Democrinus? sp.)

Donovan, S.K., W. Renema, C.A. Pinnington & C.J. Veltkamp (2012)- Significance of diadematid echinoid ossicles in micropalaeontological samples, Miocene-Pliocene of Indonesia. Alcheringa 36, 1, p. 99-105.
(On fragments of diadematoid echinoids from Miocene-Pliocene of Java, Kalimantan and Sulawesi. First report of such fossils from Neogene of the region)

Douville, H. (1923)- Sur quelques foraminiferes des Moluques orientales et de la Nouvelle Guinee. Jaarboek Mijnwezen Nederlandsch-Indie 50 (1921), Verhandelingen 2, p. 107-116.
('On some foraminifera from the eastern Moluccas and from New Guinea'. Brief description of Eocene larger forams in samples collected by Brouwer in Halmahera (Nummulites, Discocyclina, Alveolina), Roti (large Nummulites, Discocyclina), Seram (E Miocene Lepidocyclina in breccia with reworked angular clasts of Upper Cretaceous pelagic limestone), New Guinea, Kai Besar (rounded fragments of Eocene Lacazina in quartz sandstone, etc. No location info)

Douville, H. (1924)- Revision des Lepidocyclines. Memoires Societe Geologique France, N.S., 2, p. 5-49 and (1925) part 2, p. 51-123.
(Revision of lepidocyclinid Tertiary larger foraminifera, including material from various parts of Indonesia)

Doweld, A.B. (2014)- Proposals to conserve the name *Discoaster* against *Eu-discoaster*, *Helio-discoaster* and *Hemi-discoaster*, and the names *Heliodiscoaster* and *Hemidiscoaster* with those spellings (fossil Prymnesiophyta (Algae) vel Haptomonada (Protista)). Taxon 63, 1, p. 195-197.
(Commonly used calcareous nannofossil genus name Discoaster, as first described by Tan Sin Hok (1927), was an informal genus name that lacked a generic description and designation of type species. It combines 3 groups of species of formally described genera Eu-discoaster Tan (star-shaped discosterids; type species Discoaster

brouweri Tan), Helio-discoaster Tan (rosette-shaped discoasterids; type species Discoaster barbadiensis Tan and Hemi-discoaster Tan (star-shaped discoasterids with arms welded together; type species Discoaster molengraaffii Tan). This is a proposal to legitimize the common use of Discoaster)

Drooger, C.W. (1953)- Some Indonesian Miogysininae. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B56, 1, p. 104-123.
(Revision of eight miogypsinid species described from Indonesia by Rutten and Van der Vlerk, before papers of Tan Sin Hok. Four considered valid. Miogypsinella Hanzawa is junior synonym of Miogypsinoides Yabe and Hanzawa. No reason to maintain genus Conomiogypsinoides Tan Sin Hok)

Drooger, C.W. (1955)- Remarks on *Cycloclypeus*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B58, p. 415-433.
(Measurements on Cycloclypeus eidae from Tf1 (Burdigalian) of E Borneo, 40 km N of Balikpapan. No peaks of of Tan Sin Hok's 1932 'elementary species' found. Samples represent single populations)

Drooger, C.W. (1963)- Evolutionary trends in the Miogypsinidae. In: R. von Koenigswald (ed.) Evolutionary trends in foraminifera, Elsevier, Amsterdam, p. 315-349.
(Review of evolutionary patterns in Late Oligocene- M Miocene miogypsinid larger foraminifera. Includes some Indonesian/ Indo-Pacific material)

Drooger, C.W. (1984)- Evolutionary patterns in lineages of orbitoidal foraminifera. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, ser. B, 87, p. 103-130.

Drooger, C.W. (1993)- Radial foraminifera; morphometrics and evolution. Verhandelingen Koninklijke Nederlandse Akademie van Wetenschappen, Natuurkunde, I, 41, p. 1-242.
(online at: https://dwc.knaw.nl/DL/publications/PU00011102.pdf)
(Overview of evolution of Cretaceous and Tertiary larger foraminifera)

Dutta, S., S.M. Tripathi, M. Mallick, R.P. Mathews, P.F. Greenwood, M.R. Rao & R.E. Summons (2011)- Eocene out-of-India dispersal of Asian dipterocarps. Review Palaeobotany Palynology 166, p. 63-68.
(Dipterocarpaceae angiosperm trees common in rain forests of SE Asia (>470 species). Earliest dipterocarps in SE Asia from Oligocene of Borneo (Muller 1981). In W India dipterocarp resin and palynomorphs in ~53 Ma old Eocene sediments, suggesting Asian dipterocarps originated in Gondwana and dispersed from India into Asia once land connection between Indian-Asian plates established in M Eocene (49-41 Ma). Many angiosperms did not originate in SE Asian region, but dispersed into area from W Gondwanaland)

Eames, F.E., F.T. Banner, W.H. Blow & W.J. Clarke (1962)- Fundamentals of mid-Tertiary stratigraphical correlation. Cambridge University Press, p. 1-163.
(Classic study on correlations between mid-Tertiary larger and planktonic foraminifera zonations)

Eames, F.E., F.T. Banner, W.H. Blow, W.J. Clarke & A.H. Smout (1962)- Morphology, taxonomy, and stratigraphic occurrence of the Lepidocyclininae. Micropaleontology 8, 3, p. 289-322.

Evans, D., W. Muller, S. Oron & W. Renema (2013)- Eocene seasonality and seawater alkaline Earth reconstruction using shallow-dwelling large benthic foraminifera. Earth Planetary Science Letters 381, p. 104-115.
(Reconstruction of M Eocene tropical sea surface T seasonality using Nummulites djokdjokartae from Nanggulan, C Java (~39-37 Ma). Results indicate 5-6 °C annual T range, implying greater seasonality in M-Eocene (Bartonian) than in modern time (do not seem to mention estimated absolute temperature range; JTvg). Seasonal surface ocean cooling facilitated by enhanced Eocene tropical cyclone-induced upper ocean mixing, as suggested by recent modelling results)

Evans, D., N. Sagoo, W. Renema, L.J. Cotton, W. Muller, J.A. Todd, P.K. Saraswati et al. 2018. Eocene greenhouse climate revealed by coupled clumped isotope-Mg/Ca thermometry. Proc. National Academy of Sciences USA (PNAS) 115, 6, p. 1174-1179.
(online at: https://www.pnas.org/doi/pdf/10.1073/pnas.1714744115)
(Study of Mg/Ca measurements of foraminifera from globally distributed sites in tropics and mid-latitudes (including Eocene Nummulies from Nanggulan, S Java). Dataset constrains range in Eocene tropical Sea Surface Temperatures to 30-36 °C (present-day median tropical SST ~27 ° C))

Fleury, J.J., G. Bignot, A. Blondeau & A. Poignant (1985)- Biogeographie de foraminifers benthiques tethysiens du Senonien a l'Eocene superieur. Bull. Societe Geologique France (8) 1, 5, p. 757-770.
(online at: https://www.researchgate.net/publication/293108139_Biogeographie_de_Foraminiferes_benthiques_tethysiens_du_Senonien_a_l'Eocene_superieur)
(Includes global geographic distributions of Tethyan Eocene larger foram assemblages)

Fornasiero, M. (1996)- Mathematical model in benthic paleobiogeography for the Indonesian Tethyan and Post-Tethyan molluscs. In: Reports of Shallow Tethys 4, Albrechtsburg 1994, Annali Musei Civico Rovereto 11, Supplement, p. 375-386.

Frost, G.A. (1925)- Description of fish otoliths from the Tertiary formations of Atcheen, North Sumatra. Wetenschappelijke Mededeelingen Dienst Mijnbouw Nederlandsch-Indie 2, p. 1-28.
(37 species of fish otoliths from Neogene of oil field terrains of N Sumatra. Of limited biostratigraphic value?)

Germeraad, J.H., C.A. Hopping & J. Muller (1968)- Palynology of Tertiary sediments from tropical areas. Review Palaeobotany Palynology 6, p. 189-348.
(Classic paper on tropical Tertiary palynology by Shell on sections from Venezuela, Nigeria and Borneo. There are very few pantropical marker species, most are of significance only within a single botanical province)

Gerth, H. (1925)- Jungtertiare Korallen von Nias, Java und Borneo, nebst einer Uebersicht uber die aus dem Kainozoikum des Indischen Archipels bekannten Arten. Leidsche Geologische Mededelingen 1, 1, p. 22-82.
(online at: www.repository.naturalis.nl/document/549803)
('Late Tertiary corals from Nias, Java and Borneo, with an overview of the Cenozoic species known from the Indies Archipelago'. Substantial paper with descriptions of Neogene corals from localities in N and SE Nias, Tegal residency of C Java and E Kalimantan)

Gerth, H. (1931)- Coelenterata. In:B.G. Escher et al. (eds.) Onze palaeontologische kennis van Nederlandsch Oost Indie, Leidsche Geologische Mededelingen 5 (Feestbundel K. Martin), p. 120-151.
(online at: www.repository.naturalis.nl/document/549311)
(Coelenterata chapter in 'Our paleontological knowledge of the Netherlands East Indies'. Overview of coral species and occurrences in Indonesia: Devonian (New Guinea), Carboniferous (Sumatra), Permian (Timor mostly endemic solitary species), Triassic (Timor, Ceram, Misool, Sumatra), Jurassic- Cretaceous (very rare; Sumatra only), Tertiary (rel. widespread). No figures)

Gerth, H. (1935)- The distribution and evolution of the larger foraminifera in the Tertiary sediments. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 38, 4, p. 455-460.
(online at: https://dwc.knaw.nl/DL/publications/PU00016716.pdf)
(Table of Eocene- Miocene larger foram zonation on Java and comparisons with India, Europe and Americas)

Ghose, B.K. (1972)- The morphology of *Pellatispira glabra* with comments on the taxonomy, distribution and evolution of the genus. Revue de Micropaleontologie 15, 3, p. 149-162.
(Monograph on Late Eocene Indo-Pacific larger foram Pellatispira. (rrroneously accepts occurrence of Pellatispira in Australia, based on misidentified P. glabra by Chapman & Crespin (1935); JTvG))

Ghose, B.K. (1977)- Paleoecology of the Cenozoic reefal foraminifers and algae- a brief review. Palaeogeogr. Palaeoclim. Palaeoecology 22, p. 231-256.

(Review of ecologic distribution of Cenozoic larger foraminifers: (1) common 'Alveolina' in back-reef near reef core; (2) Orbitolites and Marginopora in sheltered waters on reef-flat and back-reef; (3) nummulitids and Discocyclina in both fore- and back-reef shoal areas; (4) Heterostegina in quieter waters of back-reef lagoons and reef-fiat pools; (5) Pellatispira typical fore-reef form. Also on calcareous algae)

Ghosh, M. & P.K. Saraswati (2002)- Biostratigraphic reliability of the grade of enclosure of Neogene *Lepidocyclina (Nephrolepidina)*. Indian J. Petroleum Geology 11, 2, p. 85-92.
(Grade of enclosure of protoconch by deuteroconch good estimate of relative age, but rel. high variation within sample requires statistically sufficient number of measurements. Calibrations may be different for different biogeographic provinces)

Glaessner, M.F. (1939)- Field guide to the study of larger foraminifera. Australasian Petroleum Co. Ltd., p. 1-17. *(Unpublished)*
(Larger foraminifera manual for field geologists to help identify genera in field using only a handlens)

Glaessner, M.F. (1943)- Problems of stratigraphic correlation in the Indo-Pacific Region. Proc. Royal Society of Victoria, New Series, 55, 1, p. 41-80.
(online at: https://www.biodiversitylibrary.org/partpdf/302413)
(Extensive discussion of post-Triassic statigraphy and paleontology of E Indonesia and New Guinea, in terms of geosynclinal, etc. deposits. Jurassic ammonites useful in New Guinea and the Sula Islands and are similar to Spiti shales of the Himalayas. Marine Cretaceous known from Borneo and Sumatra, Misool, Timor- E Celebes, etc. Lower Cretaceous sediments W of Timor- E Celebes syncline are mostly rich in Orbitolina and Nerinea and show Mediterranean affinities. No Orbitolina in similar-age beds in New Guinea, New Caledonia and Australia. Late Cretaceous Globotruncana limestones more widespread across all regions. Tertiary correlations largely based on Letter zonation of Van der Vlerk & Umbgrove (1927 and Tan Sin Hok. As noted by Umbgrove (1938) there are no known Paleogene deep marine basins in the Indo-Pacific region. The Mesozoic Timor- E Celebes geosyncline is characterized by strong intra-Miocene folding. Widespread Plio-Pleistocene folding. Etc.)

Glaessner, M.F. (1959)- Tertiary stratigraphic correlation in the Indo-Pacific region and Australia. J. Geological Society India 1, p. 53-67.
(Correlation of Tertiary foraminifera biozonations of India, Indonesia, New Zealand and Australia, largely based on larger foraminifera. Includes record of zone Te limestones with Spiroclypeus in Portuguese Timor (= 'Cablac Limestone' of Audley-Charles; JTvG?), associated with Aquitanian Globorotalia kugleri zone planktonic foraminifera in interbedded shales (Eames et al. 1962))

Govindan, A. (2003)- Tertiary larger foraminifera in Indian basins: a tie up with standard planktic zones and 'Letter Stages'. In: P. Kundal (ed.) Proc. XVIII Indian Colloquium on Micropaleontology and Stratigraphy, Nagpur, Gondwana Geological Magazine, Special Volume 6, p. 45-78.

Gramann, F. (1975)- Ostracoda from Tertiary sediments of Burma with reference to living species. Geologisches Jahrbuch, B, 14, p. 1-46.

Guernet, C. (1993)- Ostracodes du plateau d'Exmouth (Ocean Indien): remarques systmatiques et evolution des environnements oceaniques profonds au cours du Cenozoique. Geobios 26, 3, p. 345-360.
('Ostracodes of the Exmouth Plateau (Indian Ocean): systematic observations and deep sea environmental evolution during the Cenozoic'. Eocene- Pleistocene ostracodes from Indian Ocean DSDP wells 762 and 763. Assemblages relatively diverse until E Miocene and poor in M Miocene- Pleistocene. More than 30 species, mostly long-ranging)

Gupta, A.K. & M.S. Srinivasan (1992)- *Uvigerina proboscidea* abundances and paleoeeanography of the northern Indian Ocean DSDP Site 214 during the Late Neogene. Marine Micropaleontology 19, p. 355-367.
(Peaks of abundance of Uvigerina proboscidea inferred to represent times of high-surface productivity, related to intensified trade winds SW Indian monsoons, causing upwelling along equatorial divergemce in Indian

Ocean. Increases at ~8.5-7.5 Ma and ~6.2 Ma, highest between 5.8 and 5.1 Ma (coinciding with greatest development of Antarctic ice sheets and strong SW monsoons))

Haak, R. & J.A. Postuma (1975)- The relation between the tropical planktonic foraminiferal zonation and the Tertiary Far East Letter Classification. Geologie en Mijnbouw 54, 3-4, p. 195-198.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0U09RUV9Da09LTEE/view)
(Pioneering calibration between Tertiary planktonic foram zones and larger foram 'East Indies Letter Classification' by Shell micropaleontologists)

Hadiwisastra, S. (1994)- Penelitian nannoplangton dan biostratigrafi Tersier di Indonesia. Pros. Tridawarsa Puslitbang Geoteknologi LIPI, 2, p. 46-63.
('Studies of nannoplankton and Tertiary biostratigraphy in Indonesia'. Summary of analyses of nannoplankton from samples from (1) C Java Karangsambung, Kulunprogo and Nanggulan areas (Eocene- Miocene), (2) C Timor (Batu Putih Fm near Soe): CN12a-CN14b, Late Pliocene- Pleistocene, (3) Yamdena (Tanimbar Islands): Pliocene Batimafudi and Batilembuti Fms; NN5- NN15M Miocene- Pliocene, and Tungustuban Fm sand-shale in Oktofan area, Wermatang; Late Eocene?)

Haig, D.W., M.G. Smith, R. Riera & J.H. Parker (2020)- Widespread seagrass meadows during the early Miocene (Burdigalian) in Southwestern Australia paralleled modern seagrass distributions. Palaeogeogr. Palaeoclim. Palaeoecology 555, 109846, p. 1-25.
(Trealla Limestone in S Carnarvon Platform, contains oldest evidence for seagrass meadows in W Australia. Larger foraminifera of Lower Tf1 Letter Stage (Burdigalian, 19-16 Ma) (incl. Austrotrillina, Flosculinella). Some levels with abundant miliolid foraminifera and no corals probably deposited in and around dense seagrass meadows; other levels (e.g. coral floatstone) deposited under normal-marine conditions.Deposited during Burdigalian sea-level rise, and during part of Miocene climate optimum recognized elsewhere)

Hallock, P. & E.C. Glenn (1986)- Larger foraminifera: a tool for paleoenvironmental analysis of Cenozoic carbonate depositional facies. Palaios 1, p. 55-64.
(Modern larger foram facies distribution and Philippines Miocene comparison)

Hamilton, R. N. Amano, C.J.A. Bradshaw, F. Saltre, R. Patalano, D. Penny, J. Stevenson et al. (2023)- Forest mosaics, not savanna corridors, dominated in Southeast Asia during the Last Glacial Maximum. Proc. National Academy of Sciences USA (PNAS) 121, 1, e2311280120, p. 1-8.
(online at: https://www.pnas.org/doi/epdf/10.1073/pnas.2311280120)
(Contribution to ongoing debate whether large tracts of SE Asia's lowland rainforests were replaced with a 'savanna corridor' during cooler, more seasonal climates of Pleistocene Glacials. Late Glacial pollen records suggest expansion of montane forest in uplands and replacement of rainforest with seasonally dry tropical forest in lowlands (but resistance to savannization). Datasets indicate 'early' onset of LGM in SE Asia from ~30 ka)

Hanagata, S. & T. Nobuhara (2015)- Illustrated guide to Pliocene foraminifera from Miyakojima, Ryukyu island arc, with comments on biostratigraphy. Palaeontologia Electronica 18.1.3A, p. 1-140.
(online at: http://palaeo-electronica.org/content/pdfs/444.pdf)
(Comprehensive descriptions and illustrations of Pliocene open marine benthic and planktonic foraminifera from outcrop samples of SW Pacific island E of Taiwan)

Hanzawa, S. (1964)- The phylomorphogesis of the Tertiary foraminiferal families Lepidocyclinidae and Miogypsinidae. Science. Reports Tohoku University, ser. 2 (Geology), 35, 3, p. 295-318.
(online at: http://ir.library.tohoku.ac.jp/re/bitstream/10097/28776/1/KJ00004219393.pdf)
(Classic paper on Oligo-Miocene larger foraminifera)

Hanzawa, S. (1965)- Notes on some Discocyclinid and Nummulitid foraminifera from Java and Saipan. Science Reports Tohoku University, ser.2 (Geology), 37, 1, p. 41-47.
(online at: http://ci.nii.ac.jp/naid/110004646751/en)

(Includes re-study of Discocyclina spp. and Nummulites acutus from Nanggulan and Jiwo Hills, Java)

Harsono Pringgoprawiro, D. Kadar & S.K. Skwarko (1998)- Foraminifera in Indonesian stratigraphy, Vol. 1: Biostratigraphy, Cenozoic planktonic foraminifera. Geological Research Development Centre (GRDC), Bandung, p. 1-283.
(Major contribution/ compilation of foraminifera described from Indonesia, in 3 volumes, but very limited-distribution. Reviewed by W.A. Berggren in Micropaleontology 44, 2, p. 213-214)

Harsono Pringgoprawiro, D. Kadar & S.K. Skwarko (1998)- Foraminifera in Indonesian stratigraphy, Vol. 2: Cenozoic benthonic foraminifera. Geological Research Development Centre (GRDC), Bandung, p. 1-824.
(Listings and illustrations of 2200 species of benthic foraminifera described from Cenozoic of Indonesia)

Harsono Pringgoprawiro, D. Kadar & S.K. Skwarko (1998)- Foraminifera in Indonesian stratigraphy, Vol. 3: Palaeozoic and Mesozoic foraminifera. Geological Research Development Centre (GRDC), Bandung, p. 1-150.
(Listings and illustrations of Carboniferous- Cretaceous foraminifera described from Indonesia)

Hartono (1961)- Collection of smaller foraminifera at the Paleontological Section of the Geological Survey of Indonesia. Direktorat Geologi Indonesia, Publikasi Teknik- Seri Paleontologi 1, p. 9-19.

Hartono, H.M.S. (1964)- Coiling direction of *Pulleniatina obliquiloculata trochospira* n.var. and *Globorotalia menardii*. Geological Survey Indonesia Bull. 1, 1, p. 5-12.
(New variety of Pulleniatina obliquiloculata from Late Miocene- Pliocene at ~200m depth in well near Benoa-Denpasar road, S Bali)

Hasibuan, F. (1995)- Biostratigrafi moluska di Indonesia. Geological Research Development Centre (GRDC), Bandung, Special Publ. 16, p. 60-79.
(Molluscan biostratigraphy in Indonesia')

Hayward, B.W. & M.A. Buzas (1979)- Taxonomy and paleoecology of Early Miocene benthic foraminifera of northern New Zealand and the North Tasman Sea. Smithsonian Contr. Paleobiology 36, p. 1-154.
(online at: www.sil.si.edu/smithsoniancontributions/Paleobiology/pdf_lo/SCtP-0036.pdf)

Hayward, B.W., A.T. Sabaa, E. Thomas, S. Kawagata, R. Nomura, C. Schroder-Adams, A.K. Gupta & K. Johnson (2010)- Cenozoic record of elongate, cylindrical deep-sea foraminifera in the Indian Ocean (ODP Sites 722, 738, 744, 758 and 763). J. Foraminiferal Research 40, p. 113-133.
(Group of ~100 species of elongate, cylindrical deep-sea benthic foraminifera (families Stilostomellidae, Pleurostomellidae, Nodosariidae) from Indian Ocean ODP cores, which became extinct during cold glacial periods in Late Pliocene- mid-Pleistocene Climate Transition (~2.6- 0.6 Ma))

Ho Kiam Fui (1976)- Morphogenetic trend of *Lepidocyclina* and its application in time stratigraphy. Geologie en Mijnbouw 55, 3-4, p. 147-158.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0ZlpZYlRsbzN6b1U/view)
(Correlation between Lepidocyclina degree of curvature and planktonic foram zonation in E-M Miocene of C Luconia wells, Sarawak)

Hottinger, L. (1983)- Processes determining the distribution of larger Foraminifera in space and time. Utrecht Micropaleontological Bulletin 30, p. 239-253.
(online at: https://dspace.library.uu.nl/bitstream/1874/205889/1/Utrecht+Micropaleontological+Bulletins-30-Meulenkamp.pdf)
(Depth distribution of larger foraminifera controlled by temperature, light, symbionts, substrate, etc. Three major biogeographic provinces: (1) Caribbean, (2) Mediterranean and (3) Indo-Pacific (with Cycloclypeus, Operculinoids, Marginopora, Alveolinella))

Hottinger, L. (1997)- Shallow benthic foraminiferal assemblages as signals for depth of their deposition and their limitations. Bull. Societe Geologique France 168, p. 491-505.

Hottinger, L., E. Halicz & Z. Reiss (1991)- The foraminiferal genera *Pararotalia*, *Neorotalia* and *Calcarina*: taxonomic revision. Journal of Paleontology 65, p. 18-33.
(Oligocene smaller benthic foram Rotalia mexicana/ mecatepecensis commonly placed in genus Pararotalia, but should be Neorotalia)

Hottinger, L., J. Romero & E. Caus (2001)- Architecture and revision of the Pellatispirines, planispiral canaliferous foraminifera from the Late Eocene Tethys. Micropaleontology 47, p. 35-77.
(Detailed study of morphology of Late Eocene Pellatispira- Biplanispira- Vacuolispira larger foraminifera group, partly based on well-preserved specimens from Kalimantan. Widespread distribution across Neotethys)

Huang, C.Y. (1979)- Biometric study of *Lepidocyclina* in the Kungkuan Tuff of northern Taiwan. Acta Geologica Taiwanica 20, p. 41-51.
(Thin limestone in Kungkuan Tuff with Miogypsina, Miogypsinoides formosensis and Lepidocyclina (N). Degree of curvature indicates Aquitanian age; associated planktonic forams zone N5)

Huang, T. (1964)- '*Rotalia*' group from the Upper Cenozoic of Taiwan. Micropaleontology 10, 1, p. 49-62.
(On stratigraphic ranges of 24 species of Neogene Rotalia group (Ammonia, Asterorotalia, Pararotalia, Pseudorotalia))

Huang, T. (1983)- Significant datum planes of the Western Taiwan Neogene. Bull. Geological Society Malaysia 16, p. 47-60.
(online at: www.gsm.org.my/products/702001-101176-PDF.pdf)
(Twenty planktonic and 11 benthic foraminiferal (incl. Pseudorotalia, Asterorotalia, Pararotalia) datum planes useful for correlation recognized in marine Neogene of W Taiwan)

Huang, T. (1995)- Time and spatial distribution of some Neogene rotaliid foraminifera in the Southwest Pacific. In: S. Nishimura & R. Tsuchi (eds.) Proc. Oji Seminar on Neogene evolution of Pacific Ocean Gateways, Kyoto, IGCP-355, p. 23-33.

Hughes, G.W. (1986)- New Early Miocene occurrences of *Abyssamina quadrata* Schnitker and Tjalsma from Southeast Asia. J. Foraminiferal Research 16, 1, p. 34-35.
(online at: http://jfr.geoscienceworld.org/content/16/1/34.full.pdf)
(Abyssamina quadrata, previously recorded from Paleocene- Eocene of Atlantic Ocean, found in upper bathyal sediments of NE Kalimantan (zone N5, E Miocene) and in zones N1-N4 (Late Oligocene) of Malajon 1 well, offshore Palawan Island, Philippines. In both localities associated with upper bathyal benthic species)

Ikebe, N., M. Chiji & T. Huang (1981)- Important datum-planes of the western Pacific Neogene. Bull. Osaka Museum of Natural History 34, p. 79-86.
(online at: www.mus-nh.city.osaka.jp/publication/bulletin/bulletin/34/34-008.pdf)
(Neogene planktonic foraminifera zonation and correlation to other microfossil zonations, Indonesian Letter Stages and standard stratigraphy of Japan)

Iryu, Y., D. Bassi & W.J. Woelkerling (2009)- Typification and reassessment of seventeen species of coralline red algae (Corallinales and Sporolithales, Rhodophyta) described by W. Ishijima during 1954-1978. Palaeontology 52, 2, p. 401-427.

Ishijima, W. (1978)- Calcareous algae from the Philippines, Malaysia and Indonesia. In: T. Kobayashi & R. Toriyama (eds.) Geology and Palaeontology of Southeast Asia, University of Tokyo Press, 19, p. 167-190.
(Incl. Sporolithon borneoensis (= Archaeolithothamnium borneoense) and presumably Cretaceous-age coralline algae from W Kalimantan)

Isnaniawardhani, V. (2015)- Biostratigraphy: basics and biostratigraphic zones. Pustaka Reka Cipta, Bandung, p. 1-118.

Isnaniawardhani, V. (2017)- Prinsip dan aplikasi biostratigrafi. UNPAD Press, Bandung, p. 1-137.
('Principles and applications of biostratigraphy'. Introductory text to biostratigraphic tools in Tertiary sediments of Indonesia)

Jacques, F.M.B., G. Shi, T. Su & Z. Zhou (2015)- A tropical forest of the middle Miocene of Fujian (SE China) reveals Sino-Indian biogeographic affinities. Review Palaeobotany Palynology 216, p. 76-91.
(M Miocene Fotan flora of S Fujian, just above basalt with 14.8 ± 0.6 Ma radiometric age, considered to represent tropical rainforest based on occurrence of Dipterocarpaceae and other tropical-subtropical elements, Closer affinities to Indian Neogene than other Chinese paleofloras. During M Miocene Climatic Optimum tropical and subtropical vegetation moved N to S Fujian)

Jeannet, A. & R. Martin (1937)- Ueber Neozoische Echinoidea aus dem Niederlandisch-Indischen Archipel. Leidsche Geologische Mededelingen 8, 2, p. 215-308.
(online at: www.repository.naturalis.nl/document/549527)
('On Neozoic echinoids from the Netherlands Indies Archipelago'. Mainly paleontological descriptions of Neogene echinoid fossils from Java, Madura, Kalimantan, Timor, Ceram, N Sumatra, etc., from various European University collections)

Johnson, J.H. & B.J. Ferris (1949)- Tertiary coralline algae from the Dutch East Indies. Journal of Paleontology 23, 2, p. 193-198.
(Tertiary coralline algae, including five new species (Lithothamnium borneoense, L. nanosporum, Mesophyllum javaense, Lithophyllum parricellum, Corallina delicatula) from Eocene and Miocene of E Kalimantan and W Java, collected by LeRoy. Most common species is Lithoporella melobesioides)

Johnson, K., B.W. Hayward & A. Holbourn (2011)- Impact of the Middle Miocene climate transition on elongate, cylindrical foraminifera in the subtropical Pacific. Marine Micropaleontology 78, p. 50-64.
(58 species of elongate, cylindrical benthic foraminifera of 'Extinction Group' (Nodosariidae, Stilostomellidae, Pleurostomella; extinct during M Pleistocene Climate Transition) in ODP Sites 1146, (S China Sea) and 1237 (SE Pacific) show no major changes during major cooling in M Miocene (14.0-13.7 Ma))

Jones, R.W. (1996)- Micropalaeontology in petroleum exploration. Oxford University Press, Oxford, p. 1-432.
(Thorough general review of applied micropaleontology/biostratigraphy)

Jones, R.W., M.D. Simmons & J.E. Whittaker (2006)- On the stratigraphical and palaeobiogeographical significance of *Borelis melo melo* (Fichtel & Moll, 1798) and *B. melo curdica* (Reichel, 1937) (Foraminifera, Miliolida, Alveolinidae). J. Micropalaeontology 25, p. 175-185.
(online at: https://www.j-micropalaeontol.net/25/175/2006/jm-25-175-2006.pdf)
(Borelis melo melo ranges throughout Miocene, B. melo curdica restricted to late E- M Miocene. Both sub-species occur only in Indo-Pacific Province in late E Miocene (Burdigalian), but also present in Mediterranean province in latest Early- early M Miocene (M Burdigalian- Langhian))

Jones, T.R. & F. Chapman (1900)- On the Foraminifera of the orbitoidal limestones and reef rocks of Christmas Island. In: C.A. Andrews (ed.) A monograph of Christmas Island (Indian Ocean), Bull. British Museum (Natural History), Geology, 13, p. 226-264.
(Historic description of larger foramifera content of Tertiary limestones (on Indian Ocean seamount?), mainly from Flying Fish Cove. Include Miocene Lepidocyclina spp. and Eocene Discocyclina)

Kadar, D. & S. Soeka (1984)- Biostratigraphy of selected Neogene sequences in Indonesia. In: N. Ikebe & R. Tsuchi (eds.) Pacific Neogene datum planes; contributions to biostratigraphy and chronology, University of Tokyo Press, p. 193-202.

(Review of Neogene stratigraphy and foram datum levels of NW Java, South Central Java (Progo Mts.). NE Java (Cepu area) and East Kalimantan (Kutai Basin; zonation based on rotalid benthic foraminifera))

Kanno, S. (1986)- Revision of the genus *Vicarya* (Gastropoda) from the Indo-Pacific region. Bull. Joetsu University of Education, Niigata, 5, 3, p. 31-57.
(online at: https://juen.repo.nii.ac.jp/record/7557/files/kiyo5_3-04.pdf)
(Revision of well-known marine gastropod genus Vicarya from Indo-Pacific region. Six species in Miocene (incl. Vicarya formosa Oostingh 1935 from Bojongmanik Fm of W Java), three in Eocene. Specimens described from Eocene of Nanggulan, C Java, as Vicarya jogjacartensis by Martin 1914 too poorly preserved for identification)

Kapid, R. (2003)- Nannofosil gampingan: pengenalan dan aplikasi biostratigrafi. ITB Press, Bandung, p. 1-82.
('Calcareous nannofossils: Introduction and biostratigraphy application')

Kase, T., F. Kitao, Y.M. Aguilar, Y. Kurihara & H. Pandita (2008)- Reconstruction of color markings in *Vicarya*, a Miocene potamidid gastropod (Mollusca) from SE Asia and Japan. Paleontological Research (Palaeontological Society of Japan) 12, 4, p. 345-353.
(Includes material of Vicarya verneuili, from M Miocene Nyalindung Fm, Ciangsana, SW Java)

Kase, T., Y. Kurihara, Y.M. Aguilar, H. Pandita, A.G.S. Fernando & H. Hayashi (2015)- A new cerithioidean genus *Megistocerithium* (Gastropoda; Mollusca) from the Miocene of Southeast Asia: a possible relict of Mesozoic "Eustomatidae". Paleontological Research (Palaeontological Society of Japan) 19, 4, p. 299-311.
(online at: https://www.academia.edu/29141101/A_new_cerithioidean_genus_Megistocerithium_Gastropoda_Mollusca_from_the_Miocene_of_Southeast_Asia_a_possible_relict_of_Mesozoic_Eustomatidae_)
(New, large cerithiform gastropod genus/ species Megistocerithium magoi described, based on specimens from M Miocene Nyalindung Fm of W Java and the Philippines. Intertidal sandy mudflat dweller (mangove grazer?). M. magoi possibly relict of Mesozoic Eustomatidae)

Kase, T., Y. Kurihara, H. Hayashi, H. Pandita & Y.M. Aguilar (2008)- Age refinement of the Sonde molluscan fauna, East Java, Indonesia. Memoirs National Museum Natural Science, Tokyo, 45, p. 127-138.
(online at: www.kahaku.go.jp/research/publication/memoir/download/45/4513.pdf)
(Klitik Mb of U Kalibeng Fm along Solo River in Sonde area, E Java, Indonesia contains diverse shallow marine molluscs that have been standards of Neogene mollusc sequences in Indonesia, and were vaguely dated as Late Pliocene. Planktonic foraminifera in Klitik Mb along Solo River at Bangun, 2 km W of Sonde, suggest age between 3.95 Ma- 3.58 Ma, ~mid Pliocene (NB: unusual mix of shallow marine molluscs and siltstones with abundant deeper planktonic forams, which may be reworked from older Kalibeng Fm? Klitik Beds younger than concluded here?; JTvG))

Kay, E.A. (1990)- Cypraeidae of the Indo-Pacific: Cenozoic fossil history and biogeography. Third Int. Symposium on Marine Biogeography and Evolution in the Pacific, Bull. of Marine Science, Rosenstiel School, University of Miami, 47, 1, p. 23-34. *(online at: https://www.ingentaconnect.com/contentone/umrsmas/bullmar/1990/00000047/00000001/art00004#)*
(Includes summary/discussion of Indonesian Miocene-Recent cowries, which bear legacy of Tethyan ancestry)

Keij, A.J. (1964)- Neogene to Recent Species of *Cytherelloidea* (Ostracoda) from northwestern Borneo. Micropaleontology 10, 4, p. 415-430.
(Species of Cytherelloidea generally dominant element of Neogene-Recent shallow marine ostracode assemblages in the Indo-Malayan region. Two species groups, Seven new species. Etc.)

Keij, A.J. (1966)- Southeast Asian Neogene and Recent species of *Paijenborchella* (Ostracoda). Micropalaeontology 12, 3, p. 343-354.
(Review of occurrences of four species of Miocene- Recent highly-ornamented ostracode genus Paijenborchella Kingma 1948 in Miocene- Recent of N Borneo, Philippines, E Java, etc. Recent P. malaiensis is deep-water species living in water depths of >100m)

Keij, A.J. (1974)- Review of the Indo-Pacific species of *Triebelina* (Ostracoda). Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B77, 4, p. 345-358.
(One of a series of a pioneering papers on Neogene-Recent ostracoda from the Indo-Pacific region in 1960s-1970s, many from North Borneo, by Brunei Shell Petroleum Co. micropaleontologist Adriaan Jan Keij)

Keij, A.J. (1975)- Note on three Holocene Indo-Malayan ostracod species. Proc. Koninklijke Nederlandse Akademie van Wetenschappen,B78, 3, p. 231-241.

Keij, A.J. (1978)- Remarks on the Indo-Pacific ostracode genus *Loxoconchella*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen B81, 2, p. 215-229.

Keij, A.J. (1979)- Review of the Indo-West Pacific Neogene to Holocene ostracode genus *Atjehella*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen B 82, 4, p. 449-464.

Keij, A.J. (1979)- Brief review of the type species of genera from the Kingma collection. Proc. 7th International Symposium on Ostracodes, Belgrade, 1, p. 59-63.

Kennett, J.P., G. Keller & M. Srinivasan (1985)- Miocene planktonic foraminiferal biogeography and paleoceanographic development of the Indo-Pacific region. In: J.P. Kennett (ed.) The Miocene Ocean: paleooceanography and biogeography, Geological Society of America (GSA) Memoir 163, p. 197-236.
(Suggest closure of Indonesian Seaway around 13-12 Ma)

Kennett, J.P. & M.S. Srinivasan (1983)- Neogene planktonic foraminifera, a phylogenetic atlas. Hutchinson Ross Publ. Co., Stroudsburg, p. 1-265.

Kiel, S. (2013)- Lucinid bivalves from ancient methane seeps. J. Molluscan Studies 79, 4, p. 346-363.
(online at: https://academic.oup.com/mollus/article/79/4/346/1014058/Lucinid-bivalves-from-ancient-methane-seeps)
(Lucina-type bivalves reported from late Jurassic- late Miocene methane-seep deposits worldwide. Elliptiolucina hetzeli (Martin 1933), associated with asphalt seeps in Late Miocene sediments of Buton, Indonesia (Martin, 1933, Beets 1942) and E Pliocene seep deposits of Leyte, Philippines (Kase et al. 2007, Majima et al. 2007))

Kingma, J.T. (1948)- Contributions to the knowledge of the Young-Caenozoic Ostracoda from the Malayan region. Doct. Thesis University Utrecht, Kemink, p. 1-119.
(online at: http://dspace.library.uu.nl/handle/1874/236218)
(Late Neogene ostracods from (1) outcrop samples in Aceh (N Sumatra), (2) S Kendeng zone (E Java; 31 species), (3) Miocene-Pliocene of Bojonegoro 1 well (E Java; 41 species), and (4) Recent forms from Snellius Expedition samples in E Java Sea (19 species). Six new genera (Hemicytheridea, Atjehella, Paijenborchella, Tanella, Javanella), 94 species including 40 new. Includes description of Caudites javana Kingma, a widespread species in Indo Pacific. Limited stratigraphy/ stratigraphic results)

Koch, R.E. (1935)- Namens-Aenderung einiger Tertiar-Foraminiferen aus Niederlandisch Ost-Indien. Eclogae Geologicae Helvetiae 28, p. 557-558.
(online at: https://www.e-periodica.ch/digbib/view?pid=egh-001%3A1935%3A28%3A%3A619)
('Name changes of some Tertiary foraminifera from Netherlands East Indies'. Brief note about name changes for species described earlier by Koch (1926): Globigerina aspera= G. binaiensis, Globigerina bulloides var. tripartita= G. tripartita, etc.)

Kocsis, L. (2024)- The fossil record of marine fishes in Southeast Asia since the Miocene. 22nd Swiss Geoscience Meeting, Basel 2024, Abstract Volume, 4.9, p. 162. *(Abstract only)*
(online at: https://geoscience-meeting.ch/sgm2024/wp-content/uploads/abstract_books/SGM_2024_Abstract_Volume.pdf)

(Brief review of Neogene fossil fish recordin SE Asia, mainly Java, Madura, Sulawesi, Kalimantan and Brunei)

Kooyman, R.M., R.J. Morley, D.M. Crayn, E.M. Joyce, M. Rossetto, J.W.F. Slik, J.S. Strijk, Tao Su et al. (2019)- Origins and assembly of Malesian rainforests. Annual Review Ecology Evolution Systematics 50, 119-143.
(online at: https://www.annualreviews.org/doi/pdf/10.1146/annurev-ecolsys-110218-024737)
(Malesian rainforest assembly shaped by sharply contrasting evolutionary origins and biogeographic histories)

Kosmoswska-Ceranowicz, B., M. Sachanbinski & B. Łydzba-Kopczynska (2017)- Analytical characterization of "Indonesian amber" deposits: evidence of formation from volcanic activity. Baltica 30, 1, p. 55-60
(online at: https://gamtostyrimai.lt/uploads/documents/lediniai/Baltica/Vol-30-1-2017/...)
(Indonesian amber or "Borneo amber" widespread and associated with Miocene brown coals, like in Nyalau Fm at Merit Pila coal mine, Sarawak, which yielded world's largest piece of amber (3.5 x 1-2 m). Amber is resin from trees of family Dipterocarpaceae (Dipterocarpus, Hopea, Vateria, Vatica). Intense resin production in Cenozoic trees believed to be induced by active volcanism, when acidic volcanic tuff deposited in peat land)

Kramer, K. (1974)- Die tertiaeren Holzer Sudost-Asiens (unter Ausschluss der Dipterocarpaceae), 1 Teil. Palaeontographica. Abt. B, Palaeophytologie, 144, 3-6, p. 45-181.
('The Tertiary woods of SE Asia (with exclusion of Dipterocarpaceae', Part 1. Includes wood from Sumatra collected by Posthumus, incl. Dammaroxylon kaurioides n.sp., possibly of E Permian age (Booi et al. 2014))

Kramer, K. (1974)- Die tertiaeren Holzer Sudost-Asiens (unter Ausschluss der Dipterocarpaceae), 2 Teil. Palaeontographica. Abt. B, Palaeophytologie, 145, p. 1-150.
('The Tertiary woods of Southeast Asia (with exclusion of Dipterocarpaceae', Part 2)

Krausel, R. (1925)- Der Stand unserer Kenntnisse von der Tertiarflora Niederlandisch-Indiens. Verhandelingen Geologisch-Mijnbouwkundig Genootschap Nederland Kolonien, Geologische Serie 8 (Gedenkboek Verbeek, memorial volume), p. 329-342.
(online at: https://books.google.com/books?id=Yy0RAAAAIAAJ&pg)
('The state of knowledge of the Tertiary flora of the Netherlands Indies'. Review of known fossil plant occurrences on Java, Borneo, Sumatra)

Krijnen, W.F. (1931)- Het genus *Spiroclypeus* in het Indo-Pacifische gebied. Verhandelingen Geologisch-Mijnbouwkundig Genootschap Nederland Kolonien, Geologische Serie 9, 2, p. 77-112.
('The genus Spiroclypeus in the Indo-Pacific region'. Early review of larger foram genus Spiroclypeus Douville 1905, which evolved from involute Heterostegina borneensis by development of lateral chambers. Five species recognized: S. margaritatus, S. pleurocentralis, S. tidoenganensis, S. leupoldi and S. orbitoideus. Main distribution in latest Oligocene- Early Miocene (zone Upper Te) of Tethyan- Indo-Pacific region (in 1937 Spiroclypeus vermicularis n.sp. found in Late Eocene beds by Tan Sin Hok, 1937; part of Spiroclypeus in 1991 re-attributed to genus Tansinhokella- JTvG))

Krijnen, W.F. (1931)- Annotations to the map of the more important fossil localities in the Netherlands East Indies. Leidsche Geologische Mededelingen 5 (K. Martin Memorial Volume), p. 509-551.
(online at: https://repository.naturalis.nl/pub/505607/LGM1931005001019.pdf)
(Map and chronological listing of 330 significant Paleozoic- Recent fossil localities in Indonesia)

Kumar, P. & K.S. Soodan (1976)- Early Palaeocene planktonic foraminifera from Baratang Formation, Middle Andaman Island. Proc. 6th Indian Colloquium on Micropalaeontology and Stratigraphy, Hyderabad, p. 145-150.
(Early Paleocene planktonic foraminifera in Baratang Fm (= Mithakhari Gp) of Middle Andaman Island)

Kupper, H. (1942)- Note on a new *Cycloclypeus* from Australia. De Ingenieur in Nederlandsch-Indie (IV) 9, 1, p. 1-4.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1942-01.pdf)

(Commentary on Crespin (1940) paper on a Cycloclypeus population in Victoria, SE Australia. New species C. victoriensis is probably same as C. indopacificus Tan))

Ladd, H. (1966)- Chitons and gastropods (Haliotidae through Adeorbidae) from the Western Pacific islands. U.S. Geological Survey (USGS) Professional Paper 531, p. 1-98.
(online at: http://pubs.usgs.gov/pp/0531/report.pdf)

Ladd, H. (1970)- Eocene molluscs from Eua, Tonga. U.S. Geological Survey (USGS) Professional Paper 640-C, p. 1-12.
(online at: http://pubs.usgs.gov/pp/0640c/report.pdf)
(Many of the W Pacific islands have Late Eocene shallow marine limestones. On Eua in Tonga Group Late Eocene limestone overlies plutonic- volcanic rocks. Mainly small gastropods, probably slightly deeper marine fauna (see also Storrs Cole (1970) for larger and Todd (1970) for smaller associated foraminifera))

Ladd, H. (1972)- Cenozoic fossil molluscs from Western Pacific islands; gastropods (Turritellidae through Strombidae). U.S. Geological Survey (USGS) Professional Paper 532, p. 1-79 + 20 plates.
(online at: http://pubs.usgs.gov/pp/0532/report.pdf)
(170 species of Eocene- Pliocene gastropods from W Pacific islands. Most molluscs appear reef-associated; many are from lagoonal beds. Molluscs are of Indo-Pacific aspect, with closer ties to Indonesia and N Australia than to Ryukyus and Japan)

Ladd, H. (1977)- Cenozoic fossil molluscs from Western Pacific islands; gastropods (Eratoidae through Harpidae). U.S. Geological Survey (USGS) Professional Paper 533, p. 1-75 + 23 plates.
(online at: http://pubs.usgs.gov/pp/0533/report.pdf)
(Indo-Pacific gastropods, mainly from Late Miocene-Pliocene of Fiji, Eniwetok, Guam, Palau, etc.)

Ladd, H. (1982)- Cenozoic fossil molluscs from Western Pacific islands; gastropods (Eulimidae and Volutidae through Terebridae). U.S. Geological Survey (USGS) Professional Paper 1171, p. 1-100 + 41 plates.
(online at: http://pubs.usgs.gov/pp/1171/report.pdf)

Lambert, J. & A. Jeannet (1935)- Contribution a la connaissance des Echinides tertiaires des iles de la Sonde. I. Echinides reguliers. Memoires Societe Paleontologique Suisse, Birkhauser, 56, 1, p. 1-62.
('Contribution to the knowledge of the Tertiary echinoids of the Sonde Islands (=Indonesia)'. On of very few publications on Miocene- Recent regular echinoderms from Indonesia (Java, Borneo, Sumbawa, Timor), based on material collected by Tobler, Leupold, Gerth, Weber, etc. Many new species (manuscript initially submitted in 1934 for publication in 'Wetenschappelijke Mededeelingen' Bandung, was refused due to lack of finances))

Lei, Z.Q. (1997)- Tertiary palynological sequence and the related problems in Pearl River mouth basin. In: P. Dheeradilok et al. (eds.) Proc. International Conference Stratigraphy and tectonic evolution of Southeast Asia and the South Pacific (GEOTHAI'97), Dept. Mineral Resources (DMR), Bangkok, 1, p. 218-222.
(online at: http://library.dmr.go.th/library/Proceedings-Yearbooks/M_1/1997/7641.pdf)
(Tertiary palynology zonation of Pearl River Mouth basin, northern South China Sea. Warmest paleoclimate in M Miocene (common tropical mangroves of Florschuetzia cf. levipoli- Dacrydiumites florinii assemblage), coldest in Late Oligocene (Alnipollenites- Pinuspoleenites assemblage). Tertiary biozonation similarities with SE Asia, but calibration of Oligo-Miocene Florschuetzia zones younger than Morley 1977)

Lelono, E.B. (2001)- Revisi zonasi polen Eosen. Lembaran Publikasi Lemigas 35, 1, p. 16-26.
('Revision of Eocene pollen zonation'. Palynology study of Nanggulan Fm in C Java shows seven Eocene biozones within former Proxapertites operculatus zone)

Lelono, E.B. (2001)- Climatic effect to the distribution of the Late Paleogene indicator of pollen *Meyeripollis naharkotensis* in Western Indonesia. Lemigas Scientific Contributions 24, 2, p. 12-16.

Lelono, E.B. (2005)- Penelitian palinologi pada sedimen Paleogen di kawasan Indonesia bagian Barat. Lembaran Publikasi Minyak dan Gas Bumi (Lemigas) 39, 2, p. 15-23.
(online at: https://journal.lemigas.esdm.go.id/index.php/LPMGB/article/view/734)
('Palynological investigations of Paleogene sediments in the western part of Indonesia'. Review of Eocene-Oligocene palynology)

Lelono, E.B. (2006)- Eocene- Oligocene climate based on palynological records. Lemigas Scientific Contributions 29, 2, p. 10-23.
(online at: https://journal.lemigas.esdm.go.id/index.php/SCOG/article/view/1023/812)
(Review of Eocene-Oligocene paleoclimate trends from W Indonesia palynomorphs. M Eocene 'peak' wet climate, Late Eocene- earliest Oligocene gradually drying. Late Oligocene wetter than E-M Oligocene. Etc.)

Lelono, E.B. (2007)- Zonasi polen Tersier Indonesia Timur. Lembaran Publikasi Minyak dan Gas Bumi (Lemigas) 41, 1, p. 1-8.
(online at: https://journal.lemigas.esdm.go.id/index.php/LPMGB/article/view/45
('Tertiary pollen zonation of East Indonesia'. Differences of pollen assemblages between Papua (Australian plate) and Sulawesi- Java (Sundaland/Asian plate) necessitate separate palynozonation for E Indonesia: (1) Spinizonocolpites baculatus (Paleo-Eocene). (2) Metroxyllon salamonense (M Miocene), (3) Foveosporites spp; (4) Nothofagidites emarcida (Late Miocene; 3 sub-zones), (5)(6) Malvacipollis diversus (Pliocene), (7) 'Garcinia cuspidata type' (Late Pliocene) (8) Proteacidites spp. (Pleistocene))

Lelono, E.B. (2009)- The dispersal route of the Australian elements of *Dacrydium* and *Casuarina* from its origin to SE Asia. Scientific Contributions Oil and Gas (SCOG) Lemigas, Jakarta, 32, 3, p. 157-161.
(online at: https://journal.lemigas.esdm.go.id/index.php/SCOG/article/view/846)
Australian pollen Dacrydium may have dispersed into SE Asia prior to E Oligocene via Ninety East Ridge and Indian plate, and subsequent distribution across Sunda region and Indochina was limited by paleoclimate, explaining why it is present in some areas of Sunda region, but not others. Dispersal of Casuarina remains unresolved; migration via India unlikely as it is not known from Indian subcontinent)

Lelono, E.B. (2012)- The migration pathway of some selected Australian palynomorphs from their origin to SE Asia. Scientific Contributions Oil and Gas (SCOG) Lemigas, 35, 2, p. 49-56.
(online at: https://journal.lemigas.esdm.go.id/index.php/SCOG/article/view/777/564)
(Proposes alternative dispersal route of Australian taxa Dacrydium and Casuarina to SE Asia. Previously thought to have migrated to Sunda region after collision of Australian and Asian plates, or arrival with Gondwanan fragment in Early Oligocene. Records of Dacrydium in Eocene of Ninety East Ridge and Indian subcontinent may support alternative dispersal route into SE Asia via Indian plate)

Lelono, E.B. (2017)- Pollen records from the Oligocene of Western Indonesia as the evidences of climate changes. Scientific Contributions Oil and Gas (SCOG) Lemigas, 40, 3, p. 107-115.
(online at: http://journal.lemigas.esdm.go.id/index.php/SCOG/article/view/46
(Oligocene of SE Asia characterized by dry/ seasonal climate. Samples from offshore wells in Java, W Natuna and C Sumatera. Abundant fresh water algae of Pediastrum and Bosedinia in E Oligocene indicates presence of lacustrine sediments; dry/ seasonal climate marked by rich grass pollen Monoporites annulatus and absent or rare rain forest elements. Late Oligocene with common brackish elements indicate shifting paleoenvironment into transition-shallow marine. More humid climate evidenced by high Dacrydium and Casuarina and other rain forest palynomorphs. Wettest climate in Java region)

Leloux, J. & W. Renema (2007)- Types and originals of fossil Porifera and Cnidaria of Indonesia in Naturalis. Nat. Natuurhistorisch Museum Technical Bull., Leiden, 10, p. 1-305.
(online at: https://repository.naturalis.nl/pub/270361/TheGerthCatalogueLR.pdf)
(Listings and colorillustrations of type material of 229 taxa in large collection (>3000 specimens) of Tertiary and Permian corals in Naturalis Natural History Museum, Leiden, mainly from K. Martin, H. Gerth and J.H.F. Umbgrove collections)

Leloux, J. & F.P. Wesselingh (2009)- Types of Cenozoic Mollusca from Java in the Martin collection of Naturalis. Nationaal Natuurhistorisch Museum (NNM) Technical Bulletin 11, p. 1-765.
(online at: www.repository.naturalis.nl/document/143887)
(Updated, expanded and illustrated version of Van den Hoek Ostende et al. (2002) of 912 type specimens of Tertiary bivalves, gastropods and scaphopods from Java in K. Martin collection at Naturalis Museum, Leiden. With listing of fossil localities and 289 color plates)

Leriche, M. (1954)- Les faunes ichthyologiques marines du Neogene des Indes Orientales. Memoires Societe Paleontologique Suisse, Birkhauser, 70, p. 1-21.
('The marine fish faunas from the Neogene of the East Indies'. Incl. new species of shark tooth Carcharodon megalodon indica)

LeRoy, L.W. (1940)- The ostracode genus *Cytherelloidea* from the Late Tertiary of the Netherlands East Indies. Natuurkundig Tijdschrift voor Nederlandsch-Indie 100, p. 179-196.
(Fourteen new species of ostracode Cytherelloidea from Indonesia, incl. 3 from Late Miocene of NE Kalimantan, 4 from Mio-Pliocene from Bantam, W Java and 7 species from Miocene of C Sumatra)

LeRoy, L.W. (1941)- The ostracode genus *Cytherelloidea* from the Tertiary of the Netherlands East Indies. Journal of Paleontology 15, 6, p. 612-621.
(Same paper as above: 14 new species of ostracode Cytherelloidea from Indonesia, incl. 3 from Late Miocene of NE Kalimantan, 4 from Mio-Pliocene from Bantam, W Java and 7 species from Miocene of C Sumatra)

LeRoy, L.W. (1945)- A contribution to ostracodal ontogeny. Journal of Paleontology 19, p. 81-86.
(Includes discussion and illustrations of growth stages of Cythereis holmani from Telisa Fm of C Sumatra)

LeRoy, L.W. (1948)- The foraminifer *Orbulina universa* d'Orbigny, a suggested middle Tertiary time indicator. Journal of Paleontology 22, 4, p. 500-508.
(Lowest stratigraphic occurrence of pelagic foraminifer Orbulina universa proposed to bea good mid-Miocene markerhorizon. With discussion of stratigraphy and faunas of Kassikan section near Aliantan, Sultanate of Siak, C Sumatra, where this event occurs near top of Telisa Fm)

LeRoy, L.W. (1964)- Smaller foraminifera from the Late Tertiary of Southern Okinawa. U.S. Geological Survey (USGS) Professional Paper 454-F, p. 1-58.
(online at: http://pubs.usgs.gov/pp/0454f/report.pdf)
(Good descriptions of typical Indo-Pacific shallow and shelfal marine smaller benthic foraminifera))

Less, G. & E. Ozcan (2008)- The Late Eocene evolution of nummulitid foraminifer *Spiroclypeus* in the Western Tethys. Acta Palaeontologica Polonica 53, p. 303-316.

Leupold, W. & I.M. van der Vlerk (1931)- The Tertiary. In: B.G. Escher et al. (eds.) Stratigraphie van Nederlandsch Oost-Indie, Feestbundel K. Martin, Leidsche Geologische Mededelingen 5, p. 611-648.
(online at: www.repository.naturalis.nl/document/549456)
(Overview of Tertiary stratigraphy across 'Netherlands East Indies' in the 1931Prof. K. Martin 80st birthday memorial volume. With distribution chart of larger foraminifera and a more detailed version of the 'Letter Classification' zonation of the Tertiary. Added stages/ zones Tg and Th, subdivided Tf into 3 zones, Te into 5 zones, Ta into 2 zones. With rel. extensive review of state of knowledge of Cenozoic stratigraphies of Indonesian basins, especially from Wolfgang Leupold's multi-year geological mapping areas of NE and E Kalimantan)

Lignac-Grutterink, L.H. (1943)- Some Tertiary Corallinaceae of the Malaysian Archipelago. Verhandelingen Geologisch-Mijnbouwkundig Genootschap Nederland Kolonien, Geologische Serie 13, p. 283-297.
(Brief descriptions of Tertiary calcareous algae from Borneo, Java, etc., in Leiden collection. No location/ stratigraphy info)

Lindley, I.D. (2004)- Some living and fossil echinoderms from the Bismarck Archipelago, Papua New Guinea, and two new echinoid species. Proc. Linnean Soc. New South Wales 125, p. 115-139.
(online at: https://www.biodiversitylibrary.org/item/132499#page/120/mode/1up)

Lloyd, A.R. (1974)- Time measurement of geological time and precision in correlation. Proc. SE Asia Petroleum Exploration Society (SEAPEX) 1, Singapore, p. 31-43.
(On stratigraphic correlations, with examples from SE Asia)

Lloyd, A.R. (1975)- Paleontology and its role on oil exploration. Proc. SE Asia Petroleum Exploration Society (SEAPEX) 2, Singapore, p. 152-159.
(Brief summary of importance of paleontology in oil exploration)

Ludbrook, N.H. (1965)- Tertiary fossils from Christmas Island (Indian Ocean). J. Geological Society of Australia, 12, p. 285-294.
(Christmas Island in Indian Ocean S of W Java is Tertiary atoll formed on volcanic cone rising more >13,000' from ocean floor. Algal limestones in outcrop of Late Eocene (Tb) and Early Miocene (Te-Tf) age. Upper Eocene limestone with Discocyclina, Nummulites, Heterostegina. Lower Miocene limestone lower part with Lepidocyclina (Eulepidina), followed by Miogypsinoides dehaarti, then Flosculinella bontangensis. No rocks younger than Burdigalian identified other than young fringing reef)

Lunt, P. (2003)- Biogeography of some Eocene larger foraminifera, and their application in distinguishing geological plates. Palaeontologia Electronica 6, 1, p. 1-22.
(online at: http://palaeoelectronica.org/paleo/2003_2/geo/issue2_03.htm)
(Eocene larger foram assemblages can distinguish between carbonates from Asian-Pacific-Mediterranean (Pellatispira-Assilina) or Australian- New Guinea (Lacazinella) realms)

Lunt, P. (2013)- Foraminiferal micropalaeontology in SE Asia. In: A.J. Bowden et al. (eds.) Landmarks in foraminiferal micropalaeontology: history and development, The Micropalaeontological Society, Special Publication, Geological Society, London, p. 193-206.
(Elegant account of history of foraminiferal micropaleontology in SE Asia (mainly Indonesia) since late 1800's)

Lunt, P. & T. Allan (2004)- A history and application of larger foraminifera in Indonesian biostratigraphy, calibrated to isotopic dating. 2004 Workshop on Micropaleontology, Geological Research Development Centre Museum, Bandung, p. 1-109.
(online at: http://w.nummulites.net/PC_carbonates/Books/Lunt_Allan_2004.pdf)
(Well-illustrated, modern overview of Indonesian Tertiary larger foraminifera and zonations)

Lunt, P. & X. Luan (2022)- SE Asian Cenozoic larger foraminifera: taxonomic questions, apparent radiation and abrupt extinctions. J. of Earth Science (China) 33, 6, p. 1378-1399.
(online at: http://en.earth-science.net/en/article/doi/10.1007/s12583-022-1614-4)
(From Mediterranean to Indo-Pacific, Latest Eocene-Miocene larger foraminifera more homogenous than previously believed. Lineages show morphological radiation interrupted by extinctions. Pattern superimposed on long-term decline in larger foraminiferal abundance. Classic "large" genera were most severely hit by series of M Eocene -M-Miocene extinction events. Three major and one minor larger foraminiferal extinction events since M Eocene coincide with abrupt tectonic events across SE Asia, but there are probably multiple causes for these extinctions, including global climatic changes)

Lunt, P. & W. Renema (2014)- On the *Heterostegina- Tansinhokella- Spiroclypeus* lineage(s) in SE Asia. Berita Sedimentologi 30, p. 6-31.
(online at: https://journal.iagi.or.id/index.php/FOSI/article/view/132)
(Detailed discussion of larger foraminifera evolutionary development from Heterostegina (Vlerkina) through Tansinhokella to Spiroclypeus, which can be observed twice in fossil record: (1) in Late Eocene (Tb) and (2) in mid- Oligocene- E Miocene (Te). Morphologically no reliable way to distinguish Late Eocene and Late Oligocene tests. Second evolutionary development of Tansinhokella and Spiroclypeus happened at same time in

three geographically separate areas. Tansinhokella important for biostratigraphic subdivision of Letter Stage Te)

MacGillavry, H.J. (1962)- Lineages in the genus *Cycloclypeus* Carpenter. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B65, 5, p. 429-458.
(Eight Eocene- Recent lineages in larger foram Cycloclypeus group distinguished)

MacGillavry, H.J. (1978)- Foraminifera and parallel evolution- how or why? Geologie en Mijnbouw 57, 3, p. 385-394.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0UE1PRmVLSE1GdHM/view)
(On evolutionary trends in larger foraminifera, by former Stanvac micropaleontologist H.J. MacGillavry. With appendix B and C summarizing larger foraminifera (Cycloclypeus, miogypsinids) from Indonesia and Tertiary stratigraphy near Baturaja, S Sumatra)

Madon, M.B., R.B.A. Karim & R.W.H. Fatt (1999)- Tertiary stratigraphy and correlation schemes. In: Petronas (ed.) The Petroleum Geology and Resources of Malaysia. Petronas, Kuala Lumpur, p. 113-137.
(Review of biozonations and stratigraphic nomenclature of Malay Basin, Penyu Basin, NW Borneo, Sabah)

Malz, H. (1981)- *Atjehella jacobi* n.sp., eine pliozane Ostracoden-Art von Java. Senckenbergiana Lethaea 62, p. 167-171.
(Atjehella jacobi n.sp., a Pliocene ostracod species from Java')

Mao, L. & Swee Yeok Foong (2013)- Tracing ancestral biogeography of *Sonneratia* based on fossil pollen and their probable modern analogues. Palaeoworld 22, p. 133-143.
(online at: www.researchgate.net/publication/257458965_Tracing_ancestral_biogeography_of_Sonneratia_based_on_fossil_pollen_and_their_probable_modern_analogues)
(Review of biostratigraphy and biogeography of tropical mangrove pollen Florschuetzia, which is ancestral to modern Sonneratia. Florschuetzia documented from Late Eocene- M Miocene in paleotropics around Tethyan region. Migrated from center of origin in SE Asia probably during E Eocene, and radiated and expanded to China, Japan, Australia and E Africa. Until warm early M Miocene (Langhian) Sonneratia had largest geographical range)

Martin, K. (1924)- Eenige opmerkingen over ouderdomsbepalingen van het Indische Tertiair. De Mijningenieur 5, 2, p. 15-19.
('Some remarks on the age determinations of the Indies Tertiary'. In the past, ages of rocks often assumed from lithologic similarities to other known formations, but paleontologic studies often proved them to be wrong (quotes examples from Timor, Nanggulan, etc.). Current problem with interpretation of Tertiary faunas of Indonesia is lack of similarities with European faunas and no clear index fossils identified yet)

Martini, E. (1971)- Standard Tertiary and Quaternary calcareous nannoplankton zonation. In: A. Farinacci (ed.) Proc. of the Second Planktonic Conference, Rome, 1970, Edizioni Tecnoscienza, p. 737-785.
(Classic early synthesis of Cenozoic calcareous nannofossils biozonation (NP and NN zones))

Martino, E., P.D. Taylor, A. Kudryavtsev & J.W. Schopf (2016)- Calcitization of aragonitic bryozoans in Cenozoic tropical carbonates from East Kalimantan, Indonesia. Facies 62, 11, p. 1-15.
(online at: www.researchgate.net/publication/293479447_Calcitization_of_aragonitic_bryozoans_in_C Etc.)
(Calcification of originally aragonitic skeletons of cheilostome bryozoans Reussirella and Reptadeonella in muddy reefs from Miocene of E Kalimantan)

Matsumaru, K. (1974)- The transition of the larger foraminiferal assemblages in the Western Pacific Ocean- especially from the Tertiary period. J. of Geography (Chigaku Zasshi), Tokyo, 83, 5, p. 281-301.
(online at: https://www.jstage.jst.go.jp/article/jgeography1889/83/5/83_5_281/_pdf/-char/en)
(In Japanese. Review of stratigraphic distribution of larger foraminifera in localities across SE Asia- W Pacific, the Letter zonation and its calibration to planktonic foram zones)

Matsumaru, K. (1980)- Cenozoic larger foraminiferal assemblages of Japan, Part 1. A comparison with Southeast Asia. In: T. Kobayashi et al. (eds.) Geology and Palaeontology of Southeast Asia, University of Tokyo Press, 21, p. 211-224.
(Review of Eocene- Recent larger foram occurrences and zonation in Japan and correlation with Indonesian letter zonation. Three abundance peaks: (1) M-L Eocene, (2) Late Oligocene; (3) latest E Miocene-M Miocene)

Matsumaru, K. (2011)- A new definition of the Letter Stages in the Philippine Archipelago. Stratigraphy 8, 4, p. 237-252.
(M Paleocene- Recent Letter Stages for Philippines re-defined in terms of 17 larger foram assemblage zones)

Matsumaru, K. (2012)- Miogypsinid foraminiferal biostratigraphy from the Oligocene to Miocene sedimentary rocks in the Tethys Region. In: I.A. Dar (ed.) Earth Sciences, InTech Open Science, 25, p. 619-648.
(online at: http://cdn.intechopen.com/pdfs/27609/..)
(Good review of late Oligocene- M Miocene miogypsinid larger foram evolution and biostratigraphy

Matsumaru, K. & K. Kimura (2012)- Larger foraminifera from the Eocene Shimizu and Miocene Misaki Formations in Tosa Shimizu City, Kochi Prefecture, Japan. Transactions Proceedings Paleontological Society of Japan 156, p. 255-269.
(online at: www.palaeo-soc-japan.jp/download/TPPSJ/TPPSJ_NS156.pdf)
(Larger foraminifers from Eocene Shimizu Fm (Asterocyclina, Orbitoclypeus. Discocyclina) and Miocene Misaki Fm (Nephrolepidina) in Tosa Shimizu City, Kochi Prefecture, Shikoku, Japan)

McGowran, B. (1986)- Cainozoic oceanic and climatic events: the Indo-Pacific foraminiferal biostratigraphic record. Palaeogeogr. Palaeoclim. Palaeoecology 55, p. 247-265.
(Overall Cenozoic climatic deterioration reversed in Eocene and in Miocene by short-lived, far-reaching, extratropical excursions by tropical-type foraminifera. Widespread oceanic hiatuses appear to correlate with episodes of global warming and transgression)

McGowran, B. (2005)- Biostratigraphy: microfossils and geological time. Cambridge University Press, p.1-459.
(Text book)

McGowran, B. (2013)- Martin Glaessner's foraminiferal micropaleontology. In: A.J. Bowden et al. (eds.) Landmarks in foraminiferal micropaleontology: history and development, The Micropaleontological Society, Geological Society, London, Special Publ., p. 227-250.
(Review of M.F. Glaessner' contributions in 1940s-1950s to foraminiferal micropaleontology, including Cenozoic larger foraminifera of Papua New Guinea and biostratigraphic correlations in the Indo-Pacific region)

McGowran, B. & Q. Li (2000)- Evolutionary palaeoecology of Cainozoic Foraminifera: Tethys, Indo-Pacific, Southern Australia. Historical Biology 15, p. 3-27.
(Tertiary larger foram extinctions and migrations into higher latitudes tied to major cooling/warming events)

Mohler, W.A. (1946)- Handleiding voor het determineeren van de groot foraminiferen-genera van Nederlandsch Oost Indie. Compiled by W.A. Mohler, Private publication, p. 1-46.
('Manual for the identification of larger foram genera of Indonesia'. Small atlas of key Eocene- Recent larger foraminifera genera of Indonesia, by Swiss BPM-Shell micropaleontologist Willi Mohler, who was based in Balikpapan before and after the Japanese occupation and in Bandung during the Japanese occupation)

Morgenroth, P., A.T. Rahardjo & K.A. Maryunani (2000)- Dinoflagellate cysts- an alternative stratigraphic tool in marine Tertiary strata in Indonesia. Proc. 29th Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 4, p. 99-108.
(Initial studies of dinoflagellates from Eocene- Miocene outcrops across Java)

Morley, R.J. (1977)- Palynology of Tertiary and Quaternary sediments in Southeast Asia. Proc. 6th Annual Conv. Indonesian Petroleum Association (IPA), Jakarta, 1, p. 255-276.
(Up to eight palynozones in Oligocene- Quaternary, mainly based on evolution of Florschuetzia species of mangrove pollen, building on Germeraad et al. 1968 work)

Morley, R.J. (1982)- Fossil pollen attributable to *Alangium* Lamarck (Alangiaceae) from the Tertiary of Malesia. Review Palaeobotany Palynology 36, p. 65-94.
(On four pollen-morphological evolutionary trends and Eocene-Pliocene fossil record of genus Alangium a flowering plant, in SE Asia)

Morley, R.J. (1991)- Tertiary stratigraphic palynology in Southeast Asia: current status and new directions. Bull. Geological Society Malaysia 28, p. 1-36.
(online at: https://gsm.org.my/products/702001-101072-PDF.pdf)
(Palynology is the only biostratigraphic tool for correlation of non-marine sediments and correlation across facies. Age-restricted palynomorphs are relatively few, so Tertiary palynology seems mainly useful in correlation rather than dating. Higher resolution requires quantitative palynological zonation schemes)

Morley, R.J. (1996)- Biostratigraphic characterization of systems tracts in Tertiary sedimentary basins. In: C.A. Caughey et al. (eds.) Proc. International Symposium on Sequence Stratigraphy in SE Asia, Jakarta 1995, Indonesian Petroleum Association (IPA), p. 49-70.
(On palynomorph distribution patterns in sequences/ systems tracts)

Morley, R.J. (1998)- Palynological evidence for Tertiary plant dispersals in the SE Asian region in relation to plate tectonics and climate. In: R. Hall & J.D. Holloway (eds.) Biogeography and geological evolution of SE Asia, Backhuys Publishers, Leiden, p. 211-234.
(online at; http://searg.rhul.ac.uk/publications/books/biogeography/biogeog_pdfs/Morley.pdf)
(Tertiary plant dispersals reflect tectonic and climatic evolution of SE Asia. Sunda Eocene flora stretched as far East as S arm of Sulawesi, and after Makassar Straits opening, part of this flora became stranded E of Wallace Line. Small number of plant taxa dispersed W across Wallace line since Miocene, at 17 Ma, 14, 9.5, 3.5 and ~1 Ma. Much of Sunda region moisture deficient in Oligocene- earliest Miocene, ever-wet rainforest becoming widespread at ~20 Ma, after which they repeatedly expanded and contracted. Greatest extent of rainforest at beginning of M Miocene. Quaternary 'glacial' periods with low sea levels and more seasonal climates, leading to more pine forests and savannah. New Guinea mountains formed in M Miocene allowing dispersal of Gondwana taxa from S. Some, like Podocarpus imbricatus, Phyllocladus subsequently dispersed into SE Asia)

Morley, R.J. (2000)- Tertiary history of the Malesian flora: a palynological perspective. In: L.G. Saw et al. (eds.) Taxonomy: the cornerstone of biodiversity, Forest Research Institute Malaysia, Kepong, p. 197-210.

Morley, R.J. (2000)- Origin and evolution of tropical rain forests. Hohn Wiley & Sons, Chichester, p. 1-362.
(Incl. SE Asia chapter describing Cenozoic vegetation response to plate tectonic evolution, as reflected in Indonesia palynology records. In M Eocene SW Sulawesi has Laurasian flora, and was attached to E Kalimantan. Makassar Straits became floral-faunal migration barrier in Late Eocene. First Australian- New Guinea floral elements (Casuarina, etc.) start appearing in W Java Sea around 22-21 Ma)

Morley, R.J. (2002)- Tertiary vegetation history of SE Asia, with emphasis on biogeographical relationships with Australia. In: P. Kershaw et al. (eds.) Bridging Wallace's Line: the environmental and cultural history of the SE Asian- Australian region, Advances in Geoecology 34, Catena Verlag, p. 2-28.

Morley, R.J. (2003)- Interplate dispersal paths for megathermal angiosperms. In: Perspectives in plant ecology, evolution and systematics 6, 1-2, Urban & Fischer Verlag, p. 5-20.
(Review of dispersal of megathermal angiosperms between tectonic plates in Cretaceous and Tertiary. Early Cretaceous radiation of angiosperms unrelated to formation of Tethys. Nine dispersal routes, some tied to Late Cretaceous- E Tertiary Gondwana break-up and routes formed since M Eocene phases of plate collision)

Morley, R.J. (2007)- Cretaceous and Tertiary climate change and the past distribution of megathermal rainforests. In: M.B. Bush, J.R. Flenley & W.D. Gosling (eds.) Tropical rainforest responses to climatic change, Chapter 1, Springer-Praxis, p. 1-31. *(see also 2nd Edition, 2011)*
(Review of long term climate patterns and distribution of tropical rainforests)

Morley, R.J. (2011)- Dispersal and paleoecology of tropical Podocarps. Smithsonian Contr. Botany 95, p. 21-41.
(online at: http://smithsonianrex.si.edu/index.php/scb/article/download/175/131)
(Tropical Podocarpaceae family appeared in Triassic of Gondwana and essentially remained southern family. Podocarpus s.l. dispersed into SE Asia in Late Eocene, explained by dispersal from India and possibly multiple dispersal events from Australia. Dacrydium reached SE Asia in E Oligocene and expanded range to Japan during M Miocene climatic optimum. Dacrycarpus and Phyllocladus dispersed into New Guinea as island emerged in Late Miocene, then island hopped to Borneo in M Pliocene. Dacrycarpus reached Sumatra and Malay Peninsula in Pleistocene)

Morley, R.J. (2012)- A review of the Cenozoic palaeoclimate history of Southeast Asia. In: D.J. Gower et al. (eds.) Biotic evolution and environmental change in Southeast Asia, The Systematics Association, Cambridge University Press, Chapter 6, p. 79-114.
(online at: https://www.researchgate.net/publication/285737329_A_review_of_the_Cenozoic_palaeoclimate_history_of_Southeast_Asia)
(Summary of Cenozoic climatic and environmental history of Sunda region, from Sulawesi to S Vietnam, based on palynological records, occurrence of coals (= everwet climate) and paleosols)

Morley, R.J. (2018)- Assembly and division of the South and South-East Asian flora in relation to tectonics and climate change. J. Tropical Ecology 34, 4, p. 209-234.
(online at: www.researchgate.net/publication/326628937_Assembly_and_division_of_the_South_ Etc.)
(On main phases of plant dispersal into and out of SE Asia. Late Cretaceous poorly understood. India's drift into perhumid low latitudes during Eocene brought dispersal into SE Asia of megathermal angiosperms which originated in W Gondwana, and became established since ~49 Ma, with terrestrial connection after ~41 Ma. Oligocene was time of seasonally dry climates except along E and S seaboard of Sundaland. At end of Oligocene widespread perhumid conditions across region. Uplift of Himalaya during M Miocene thermal maximum, created dispersal of E Asian evergreen taxa into N India. After Australasia- Sunda collision and uplift of New Guinea and islands of Wallacea, Gondwanan montane taxa also found their way into region)

Morley, R.J. & J.R. Flenley (1987)- Late Cainozoic vegetational and environmental changes in the Malay Archipelago. In: T.C. Whitmore (ed.) Biogeographic evolution of the Malay Archipelago, Oxford Monographs Biogeography 4, Clarendon Press, Oxford, p. 50-59.

Morley, R.J. & Jaizan H.M. Jais (2024)- Pollen and spore atlas for the Cenozoic of the Malay Basin, Malaysia. Geological Society of Malaysia, p. 1-272.
(online at: https://gsm.org.my/articles/pollen-and-spore-atlas-for-the-cenozoic-of-the-malay-basin-malaysia/)
(Milestone atlas with descriptions of 186 Cenozoic pollen, spore and algal form-taxa that are important in evaluation of palynostratigraphy and environment interpretation in Oligocene-Pliocene of Malay Basin. With 7 new genera, 55 new form-species, 4 new forms and 10 new combinations. With summaries of PR1-PR15 bio-zonation, vegetation types, basin evolution, paleogeography and extensive references list)

Morley, R.J., E.B. Lelono, L. Nugrahaningsih & Nur Hasjim (2000)- LEMIGAS Tertiary palynology project: aims, progress and preliminary results from the Middle Eocene to Pliocene of Sumatra and Java. Geological Research Development Centre (GRDC), Paleontology Series 10, Bandung, p. 27-47.
(Summary of palynology work in Java (Eocene of Nanggulan and Bayah), Sumatra (E Oligocene Pematang Fm, Late Oligocene Talang Akar Fm, E Miocene Gumai Fm, M Miocene Air Benakat Fm. No positive records of Paleocene- Early Eocene deposits in W Indonesia)

Morley, R.J. & H.P. Morley (2013)- Mid Cenozoic freshwater wetlands of the Sunda region. J. Limnology 72, 2s, p. 18-35.
(online at: www.jlimnol.it/index.php/jlimnol/article/view/704)
(On development of widespread rift lake systems in Oligocene of SE Asia and their eventual demise following marine transgression. Pollen and spores illustrate multitude of fresh and brackish water swamp communities around margins)

Morley, R.J. & H.P. Morley (2013)- Improving stratigraphic resolution in Indonesia using the method of sequence biostratigraphy. Proc. 47th Annual Conv. Indonesian Petroleum Association (IPA), Jakarta, IPA23-G-63, p. 1-17.
(online at: https://www.researchgate.net/publication/373546134_Improving_stratigraphic_resolution_in_Indonesia_using_the_method_of_sequence_biostratigraphy)
(About dating non-marine successions by analyzing cyclical patterns in palynological, etc., data that reflect astronomically-driven sea level changes. Palaeogene non-marine basins such as Cuu Long and Song Hong Basins in Vietnam exhibit depositional cycles reflecting 406 kyr eccentricity-driven astronomical cycles. In Mahakam Delta Miocene multiple sands within main reservoir interval driven by 106 ka eccentricity and 41 ka obliquity changes. Etc.)

Morley, R.J. & H.P. Morley (2022)- The prelude to the Holocene: tropical Asia during the Pleistocene. Holocene Climate Change and Environment 2022, p. 1-32.
(Review of climate and vegetation history for tropical Asia for last 3 Myrs. Pliocene witnessed demise of humid tropical forests across India, which were replaced at least in the north by open grasslands, which supported a diverse fauna, elements of which migrated to Java during E Pleistocene. Fauna did not migrate along a "savanna corridor"(no representatives of Indochina deciduous forests or savanna). During Pleistocene glaciation, rain forests remained along equatorial SE Asia, with savanna in Indochina and expansion of seasonal evergreen and deciduous forests across emergent Java Sea and Java)

Muller, J. (1972)- Palynological evidence for change in geomorphology, climate and vegetation in the Mio-Pliocene of Malesia. In: P.S. & M. Ashton (eds.) The Quaternary era in Malesia, University of Hull, Department of Geography Miscellaneous Series 13, p. 6-16.

Muller, J. (1978)- New observations on pollen morphology and fossil distribution of the genus *Sonneratia* (Sonneratiaceae). Review Palaeobotany Palynology 26, p. 277-300.
(Study of mangrove pollen Sonneratia, and fossil representatives Florschuetzia. May have originated in Eocene in Tethys area, but earliest reliable occurrence (Florschuetzia trilobata) from E Miocene in Sunda area)

Murgese, D.S. & P. De Deckker (2005)- The distribution of deep-sea benthic foraminifera in core tops from the eastern Indian Ocean. Marine Micropaleontology 56, p. 25-49.
(57 core tops between 700- 4335m from E Indian Ocean between Australia and Indonesia. Seven key-species useful for environments. Two species groups: (1) Oridorsalis tener umbonatus, Epistominella exigua and Pyrgo murrhina (cold, well-oxygenated, low carbon flux to sea floor) and (2) Nummoloculina irregularis and Cibicidoides pseudoungerianus (upper-bathyal). Uvigerina proboscidea mainly at low latitudes, with high carbon flux due to higher primary productivity at sea surface, and low oxygen levels due to organic matter oxidation and presence of oxygen-depleted Indonesian Intermediate Water and N Indian Intermediate Water)

Nishida, S. (1987)- Calcareous nannoplankton biostratigraphy in the Sunda Arc. Committee Co-ord. Joint Prospecting Mineral Resources in Asian Offshore Areas, CCOP, Technical Bull. 19, p. 69-72.
(online at: https://ncloud.dotdream.co.th/index.php/s/xCwHAtAFYxzBtC9)
(Bottom samples from S Java continental slope and both flanks of Sunda Trench contain modern nannofossil assemblages dominated by Umbellosphaera irregularis and Gephyrocapsa oceanica)

Novak, V. (2014)- Larger benthic foraminifera in Miocene carbonates of Indonesia. Ph.D. Thesis Universiteit Utrecht, Utrecht Studies in Earth Sciences 64, p. 1-211.
(online at: https://dspace.library.uu.nl/handle/1874/300544)

(Collection of papers on Miocene larger benthic foraminifera from Kutai Basin and Taballar Lst from Mangkalihat Peninsula (E Kalimantan), Bulu Fm (C Java))

Novak, V. & W. Renema (2018)- Ecological tolerances of Miocene larger benthic foraminifera from Indonesia. J. Asian Earth Sciences 151, p. 301-323.
(online at: https://www.sciencedirect.com/science/article/pii/S1367912017306235)
(Paleoenvironmental models for distribution of Miocene larger foraminifera, based on cluster analysis of assemblages from E-M Miocene mixed carbonate-clastic series in Kutai Basin, E Kalimantan, and Serravallian Bulu Fm of NE Java)

Nuttall, W.L.F. (1926)- A revision of the *Orbitoides* of Christmas Island (Indian Ocean). Quart. J. Geological Society, London, 82, p. 22-43.
(Eocene limestones with Discocyclina, and E Miocene limestones with Lepidocyclina, Miogypsina, Spiroclypeus)

O'Dogherty L., J.P. Caulet & P. Dumitrica & N. Suzuki, (2021)- Catalogue of Cenozoic radiolarian genera (Class Polycystinea). Geodiversitas 43, 21, p. 709-1185.
(online at: https://sciencepress.mnhn.fr/sites/default/files/articles/pdf/g2021v43a21.pdf)
(Extensively illustrated revision of Cenozoic polycystine radiolaria: 1683 revised genera, many originally described by Ehrenberg and Haeckel in the late 1800s)

O'Dogherty L., N. Suzuki, J.P. Caulet & P. Dumitrica (2022)- Inventory of Cenozoic radiolarian species (Class Polycystinea) - 1834-2020. Geodiversitas 44, 5, p. 75-205.
(online at: https://sciencepress.mnhn.fr/sites/default/files/articles/pdf/g2022v44a5.pdf
(Complete list of Cenozoic Polycystinea radiolaria, reported between 1834 and 2020 (6898 names))

O'Herne, L. (1972)- Secondary chamberlets in *Cycloclypeus*. Scripta Geologica, Leiden, 7, p. 1-35.
(Biometrical study of Miocene Cycloclypeus from Van der Vlerk collection from NE Java, Madura and Larat)

O'Herne, L. (1974)- A reconsideration of *Amphistegina lessonii* d'Orbigny, 1826, sensu Brady, 1884 (Foraminifera). Scripta Geologica, 26, p. 1-53.
(online at: www.repository.naturalis.nl/document/148755)
(Discussion of Oligocene-Recent Amphistegina species, mainly from Java)

O'Herne, L. (1976)- A comparison of *Lepidocyclina ferreroi* (Provale, 1909) and *Lepidocyclina multilobata* (Gerth, 1939). Scripta Geologica, Leiden, 35, p. 1-48.
(online at: www.repository.naturalis.nl/document/148824)
(Comparison of two stellate Lepidocyclina species from E-M Miocene of Java, Madura and Borneo)

O'Herne, L. & I.M. van der Vlerk (1971)- Geological age determinations on a biometrical basis (comparison of eight parameters). Boll. Societa Paleontologica Italiana 10, 1, p. 3-18.

Ong Goan Eng (1960)- The development of Shell micropaleontological studies in South Sumatra. Contributions Department of Geology, Institute of Technology Bandung (ITB) 47, p. 3-9.
(Very general paper on use of micropaleontology since 1930 by BPM/ Shell in S Sumatra. Discussion of biofacies versus lithostratigraphy and time. No details on faunas, stratigraphy, etc.)

Oostingh, C.H. (1938)- Mollusken als gidsfossielen voor het Neogeen in Nederlandsch Indie. Handelingen 8e Nederlandsch-Indisch Natuurwetenschappelijk Congres, Soerabaja 1938, p. 508-516.
(online at: https://resolver.kb.nl/resolve?urn=MMKB31:042287000:pdf)
('Molluscs as index fossils for the Neogene in the Netherlands Indies'. Crude succession of mollusc assemblages (no range charts), based on Tertiary collections from Banten (SW Java) by Koolhoven and Musper. Gastropod Vicarya callosa is M Miocene (Tf2-3) index species. Other potential marker species mainly

Turritella spp. No illustrations, range charts, etc. Very few of Neogene Java species known from elsewhere (unlike Eocene species; JTvG))

Osberger, R. (1956)- Korallen als Hilfsmittel der Tertiar und Quartar-Stratigraphie Indonesiens. Djawatan Geologi (Indonesia Geol. Survey), Publikasi Keilmuan 32, p. 1-79.
('The use of corals as a tool in Tertiary and Quaternary stratigraphy of Indonesia'. Elegant review with listings of all Eocene- Pliocene fossil coral faunas described from Indonesia. Percentage of living coral species increases from 0% in Eocene-Oligocene, 6-9% in Early Miocene, 15-30% in M Miocene, 30-60% in Late Miocene-Pliocene and ~80% in Pleistocene. With range charts)

Ozcan, E., A.O. Yucel, L.S. Erkızan, M.N. Gultekin, S. Kaygılı & S. Yurtsever (2022)- Atlas of the Tethyan Orthophragmines. Mediterranean Geoscience Reviews 4, p. 3-213.
(online at: https://link.springer.com/article/10.1007/s42990-022-00072-1)
(Brilliantly illusrrated review of larger foraminifera of the Paleo-Eocene Discocyclina-Asterocyclina-etc. group from the Tethys region. Not much information from Indonesian region, except to mention common occurrence in Borneo of Discocyclina omphalus Von Fritsch)

Pandita, H., Y. Zaim, Aswan & Y. Rizal (2012)- The evolution pattern of Turritelline shell morphology in Java; relationship with geochronological aspects. Proc. 1st Earth Science Int. Seminar, Yogyakarta, 1, p. 187-197.

Panuju (2010)- The high resolution Neogene and Quaternary nannoplankton zonation for Indonesian basins. Proc. 39th Annual Conv. Indonesian Association Geologists (IAGI), Lombok, PIT-IAGI-2010-265, p. 1-14.
(online at: https://www.iagi.or.id/web/digital/12/2010_IAGI_Lombok_The-High-Resolution-Neogene.pdf)
(New high resolution Neogene-Quaternary nannoplankton zonation for Indonesia, mostly based on material from NE Java, NW Java, Sumatra, Kutei, S Sulawesi, Salawati, Bintuni and Waipoga-Waropen Basins. The 21 standard zones of Martini (1971) can be further subdivided into 58 subzones)

Parker, F.L. (1967)- Late Tertiary biostratigraphy (planktonic foraminifera) of tropical Indo-Pacific deep-sea cores. Bull. American Paleontology. 52, 235, p. 115-208.
(Late Miocene- Pliocene planktonic foram zonation on samples from deep sea cores and Fiji outcrop samples)

Piccoli, G. (1984)- Cenozoic molluscan associations of Mediterranean and Southeast Asia: a comparison. Memorie di Scienze Geologiche, Padova, 36, p. 499-521.

Piccoli, G. (2002)- Tethyan exchange of benthic molluscs between SE Asia and Mediterranean in the Paleogene Memorie di Scienze Geologiche, Padova, 54, p. 1-8.

Piccoli, G., E. Robba, C. Chonglakmani, P. Dheeradilok & R. Ingavat (1984)- Paleogene Tethyan and Neogene Indo-Pacific shallow benthic molluscs: some thoughts. J. Geological Society Thailand 7, 1, p. 13-18.
(online at: https://library.dmr.go.th/multim/JGST/26.pdf)
(Eocene molluscs of Nanggulan, Java, many affinities with coeval European species (not endemic as usually suggested by K. Martin a.o.), pointing to Tethyan seaway connection). Most of Miocene molluscs from Java different from Mediterranean suggesting seaway closed by E Miocene (Neogene- Recent Indo-Pacific province). Also discussion of Quaternary molluscs from subsurface near Bangkok)

Posthumus, O. (1929)- On paleobotanical investigations in the Dutch East Indies and adjacent regions. Bull. Jardin Botanique Buitenzorg, ser. 3, 10, 3, p. 374-384.
(Historic, brief review of papers on plant fossils since Goppert (1854; Java), Heer (1874, 1879; Ombilin, Sumatra), Geyler (1877; S Borneo), etc. until 1927)

Postuma, J.A. (1971)- Manual of planktonic foraminifera. Elsevier Publishing Co., Amsterdam, p. 1-417.
('Classic' Shell Oil Company manual of Mid-Cretaceous- Recent planktonic foraminifera)

Poumot, C. (1989)- Palynological evidence for eustatic events in the Tropical Neogene. Bull. Centres Recherches Exploration-Production Elf-Aquitaine 13, 2, p. 437-453.

Prakash, U. (1971)- Fossil woods from the Tertiary of Burma. The Palaeobotanist 20, 1-3, p. 48-70.
(online at: https://jpsonline.co.in/index.php/jop/article/view/887/874)
(Fossil woods of genera Dipterocarpus and Shorea-Pentacme (Dipterocarpaceae), Gluta-Melanorrhoea (Anacardiaceae), Sterculia, Acacia, Cynometra, Cassia, etc, from Tertiary beds of Burma. Collected by unknown Burmese collector and sent to Prof. B. Sahni in India. Ages and locality uncertain)

Prakash, U. & M.B. Bande (1980)- Some more fossil woods from the Tertiary of Burma. Palaeobotany 26, 3, p. 261-78.
(online at: https://jpsonline.co.in/index.php/jop/article/view/1071/1058)
(Fossil woods comparable to Shorea ovata (Dipterocarpaceae), Swintonia, Albizia, Careya, Lagerstroemia and Araucaria-Agathis from Burma Neogene. Most species also in present day flora of Burma and nearby areas)

Prijosoesilo, P. (1972)- Calcareous nannoplankton, a new biostratigraphic tool in the oil industry with emphasis in Indonesia. Proc. 1st Annual Conv. Indonesian Petroleum Association (IPA), Jakarta, p. 43-56.
(Brief review of use of nannofossils in Cenozoic of SE Asia)

Prins B. (1971)- Speculations on relations, evolution, and stratigraphic distribution of discoasters. In: A. Farinacci (ed.) Proc. 2nd Planktonic Conference, Roma 1970, Ediz. Tecnoscienza, 2, p. 1017-1031.
(Discussion of legitimacy of calcareous nannofossil genus Discoaster Tan Sin Hok 1927)

Proto Decima, F. & C. Masotti (1981)- The genus *Gephyrocapsa* (Coccolithophorales) in the Plio-Pleistocene of the Timor Trough. Memorie di Scienze Geologiche, Padova, 34, p. 453-464.

Rachman, R.S. & Winantris (2023)- The significance relationship between palynomorph fossil preservation and grain size in rock: case study of Walat and Batuasih Formations, Sukabumi, West Java. Jurnal Geocelebes 7, 1, p. 17-28.
(online at: https://journal.unhas.ac.id/index.php/geocelebes/article/view/22746)
(Palynomorphs in Late Eocene-Oligocene Walat and Batuasih Fms, W Java, better preservation in fine-grained rocks. Large palynomorph grains better preserved than smaller ones)

Rahmawati, D., S.S. Surjono, D.H. Barianto & W. Rahardjo (2024)- Bioerosion in the Late Eocene *Discocyclina discus sowerbyi* (Nuttall, 1926) in Bayat area, Indonesia. J. Tropical Biodiversity and Biotechnology (JTBB) (UGM) 9, 4, jtbb93779. p. 1-16.
(online at: https://journal.ugm.ac.id/jtbb/article/view/93779/40078)
(Bioerosion by predatory gastropods, polychaete worms and parrotfish in Late Eocene Discocyclina discus sowerbyi tests in Gamping Beds, Bayah, southern C Java. Four bioerosional trace fossils)

Raju, D.S.N. (1973)- *Miogypsina indica*, a new species of Miogypsinidae from the Miocene of India. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B76, p. 140-142.

Raju, D.S.N. (1974)- Study of Indian Miogypsinidae. Utrecht Micropaleontological Bulletin 9, p. 1-148.
(online at: https://dspace.library.uu.nl/bitstream/1874/205813/1/Utrecht+Micropaleontological+Bulletins-9-Raju.pdf)
(Study of Miogypsinoides- Miogypsina evolution in India. Highly applicable to the latest Oligocene-Middle Miocene Miogypsinoides- Miogypsina lineage(s) in Indonesian carbonates)

Raju, D.S.N. (2011)- Oligo-Miocene larger foraminiferal zones of India and their importance in classification and correlation. Oil and Natural Gas Corporation (ONGC) Bulletin 46, p. 40–62.

Reich, S., V. Warter, F.P. Wesselingh, H. Zwaan, L. Lourens & W. Renema (2015)- Paleoecological significance of stable isotope ratios in Miocene tropical shallow marine habitats (Indonesia). Palaios 30, 1, p. 53-65.
(Aragonitic shells of Burdigalian and Tortonian molluscs from C Java and E Kalimantan analyzed for δ18O and δ13C ratios. Depleted δ18O and δ13C ratios in brackish water samples. Also chemosymbiotic species show depleted δ13C ratios. Seagrass communities yield comparatively enriched δ13C ratios. Stable isotope ratios may provide additional evidence for distinguishing paleoenvironments)

Renema, W. (2002)- Larger foraminifera as marine environmental indicators. Scripta Geologica 124, p. 1-260.
(online at: https://repository.naturalis.nl/pub/219196)

Renema, W. (2005)- The genus *Planorbulinella* (Foraminiferida) in Indonesia. Scripta Geologica 129, p. 137-146.
(online at: https://repository.naturalis.nl/document/41886)
(Late Oligocene- Miocene Planorbulinella from Java, SE Borneo, W Sulawesi. Two new species)

Renema, W. (2006)- Comment on "Significant Miocene larger foraminifera from South Central Java" by M.K. BouDagher-Fadel and S.W. Lokier. Revue de Paleobiologie, Geneve, 25, 1, p. 405-406.
(Argues for maintaining Lepidocyclina and Eulepidina as separate genera)

Renema, W. (2007)- Fauna development of larger benthic foraminifera in the Cenozoic of Southeast Asia. In: W. Renema (ed.) Biogeography, time, and place: distributions, barriers, and islands, Topics in Geobiology 29, Springer, p. 179-215.
(online at: www.researchgate.net/publication/225246778_Fauna_Development_of_Larger_Benthic_Foraminifera_in_the_Cenozoic_of_Southeast_Asia)
(Overview of Far East (Indo-West Pacific) Tertiary larger foraminifera zonations)

Renema, W. (2008)- Internal architecture of Miocene *Pseudotaberina* and its relation to Caribbean Archaiasins. Palaeontology 51, 1, p. 71-79.
(Soritid LF Pseudotaberina malabarica described from material collected by Martin in 1911 from Burdigalian (Tf1) Jonggrangan Fm near Yogyakarta, C Java. Also known from W Java (Tf2), E Kalimantan, PNG, etc.)

Renema, W. (2015)- Spatiotemporal variation in morphological evolution in the Oligocene-Recent larger benthic foraminifera genus *Cycloclypeus* reveals geographically undersampled speciation. GeoResJ 5, p. 12-22.
(online at: www.sciencedirect.com/science/article/pii/S2214242814000217)
(Genus Cycloclypeus ranges from Oligocene- Recent, first appearing in E Rupelian of Java and Kalimantan (C. koolhoveni). Late Oligocene- Recent C. eidae to C. carpenteri lineage in Mediterranean and Indo- West Pacific provinces. C. annulatus derived from C. eidae as separate lineage in late Early-M Miocene)

Renema, W. & L. Cotton (2015)- Three dimensional reconstructions of *Nummulites* tests reveal complex chamber shapes. PeerJ, 3:e1072, p. 1-14.
(online at: https://peerj.com/articles/1072/)
(Study of details of interior morphology of M Eocene (~38-40 Ma) Nummulites djokdjokartae from Nanggulan Fm at Kali Watupuruh, Nanggulan, C Java. At top of sampled interval abundant large Discocyclina indicating M Bartonian Ta-Tb boundary)

Renema, W., A. Racey & P. Lunt (2002)- Palaeogene nummulitid foraminifera from the Indonesian Archipelago: a review. Cainozoic Research 2, 1-2, p. 23-78.
(same paper in Renema 2002, Scripta Geologica 124, p. 110-165)
(online at: https://natuurtijdschriften.nl/pub/541685/CR2003002001004.pdf)
(60 species of Eocene-Oligocene Nummulites reported from Indonesia, only 7 believed to be valid. Sangiran mud volcano boulders of Nummulites-Pellatispira limestone with Nummulites gerthi/ N. pengaronensis and planktonic foraminifera (zone P15; around Middle-Late Eocene boundary. Samples from Miomaffo, Timor (= from allochthonous Sundaland-derived'Banda Terrane') with Nummulites and Pellatispira)

Renz, O. & H. Kupper (1946)- Uber morphogenetische Untersuchungen an Grossforaminiferen. Eclogae Geologicae Helvetiae 39, p. 317-342.
(online at: http://retro.seals.ch/digbib/view?pid=egh-001:1946:39::344)
('On morphogenetic investigations of larger foraminifera')

Resiwati, P., J.W. Farrell, T.R. Janecek, J.K. Weissel et al. (1992)- Calibration of Late Neogene calcareous nannofossil datum planes with the paleomagnetic record From Site 758, ODP Leg 121, Ninetyeast Ridge, Indian Ocean. Proc. 21st Annual Conv. Indonesian Asssoc. Geologists (IAGI), 2, p. 417-436.
(online at: https://www.iagi.or.id/web/digital/50/21th-Vol-II-(7-10-Des-1992)-47-66.pdf)
(Correlation between paleomagnetic reversal scale and calcareous nannofossil zonation at ODP Site 758 core on northern part of Ninetyeast Ridge in NW Indian Ocean for last 7 Myrs)

Riedel, W.R. & A. Sanfilippo (1974)- Radiolaria from the Southern Indian Ocean, DSDP Leg 26. In: B.P. Luyendyk & T.A. Davies (eds.) Initial Reports Deep Sea Drilling Project (DSDP) 26, Chapter 33, p. 771-813.
(online at: www.deepseadrilling.org/26/volume/dsdp26_33.pdf)
(Radiolaria from Leg 26 Sites 250-258 in Indian Ocean W of Australia. Dictyomitra lilyae Tan Sin Hok, 1927= Early Cretaceous (Albian?) age range. Pages 773-774: unpublished nannofossil work on Tan Sin Hok's Bebalain 150 sample from Roti island by Bukry: presence of Cretarhabdus crenulatus, Eiffellithus turriseiffeli, Manivitella pemmatoidea, Parhabdolithus embergeri, Vagalapilla matalosa, Watznaueria barnesae and W. biporta, indicating Aptian-Turonian age. Other samples of Tan Sin Hok from Roti (no. 149, 154, and 384) contain similar radiolarian assemblages and probably of same age)

Riedel, W.R. & A. Sanfilippo (1978)- Stratigraphy and evolution of tropical Cenozoic radiolarians. Micropaleontology 24, p. 61-96.
(Review of zonation and datum levels of low-latitude Cenozoic radiolarians)

Rogl, F. (1974)- The evolution of the *Globorotalia truncatulinoides* and *Globorotalia crassaformis* group in the Pliocene and Pleistocene of the Timor Trough, DSDP Leg 27, Site 262. In: J.J. Veevers et al. (eds.) Initial Reports Deep Sea Drilling Project (DSDP) 27, Washington, p. 769-771.
(online at: www.deepseadrilling.org/27/volume/dsdp27_37.pdf)
(DSDP Site 262 in Timor Trough SW of Timor Island with rel. thick U Pliocene-Pleistocene. Good record of evolutionary trends in Globorotalia crassaformis- G. tosaensis- G. truncatulinoides groups)

Rutten, L. (1924)- Kleine mededeelingen over foraminiferen uit Nederlandsch-Indie. Verslagen Koninklijke Akademie van Wetenschappen, Amsterdam, 23, 6, p. 539-544.
(see English version, Rutten (1924) (Brief communications on foraminifera from the Netherlands Indies'))

Rutten, L. (1924)- Some notes on foraminifera from the Dutch Indies. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 23, 6, p. 529-534.
(online at: https://dwc.knaw.nl/DL/publications/PU00015069.pdf)
(English version of Rutten (1924, above). Includes: 1. Linderina limestone on Bacan (probably Late Miocene or Pliocene calcareous sandstone rich in Planorbulinella); (2) Embryonal chambers of Lepidocyclina acuta from Balikpapan area, E Kalimantan, (3) Age of limestone on island Kamarang) in Pare Pare Bay, S Sulawesi (Late Neogene, common Operculina without Lepidocyclina, etc.; (4) Lepidocyclina cf. epigona in Globigerina-chert of Skru Island (W New Guinea) (confirmation of small, but true Lepidocyclina))

Saito, T. (1977)- Late Cenozoic planktonic foraminiferal datum levels: the present state of knowledge toward accomplishing Pan-Pacific stratigraphic correlation. Proc. First International Congress Pacific Neogene stratigraphy, Tokyo 1976, p. 61-80.

Saito, T., L.H. Burckle & J.D. Hays (1975)- Late Miocene to Pleistocene biostratigraphy of equatorial Pacific sediments. In: T. Saito & L.H. Burckle (eds.) Late Neogene Epoch boundaries, Micropaleontology Press, p. 226-244.

Samanta, B.K. (1984)- The genus *Biplanispira* Umbgrove (Foraminiferida) and its occurrence in India. Geological Magazine 121, 4, p. 311-318.
(Biplanispira closely related to Pellatispira, from which it probably evolved. Biplanispira is confined to region between E India and Eua, Tonga, and middle and upper parts of Upper Eocene zone Tb. In U Eocene of Garo Hills, E India, represented by B. hoffmeisteri (Whipple) and from S Andaman Island by B. mirabilis)

Sanfilippo, A., M.J. Westberg-Smith & W.R. Riedel (1985)- Cenozoic Radiolaria. In: H.M. Bolli, J.B. Saunders & K. Perch-Nielsen (eds.) Plankton Stratigraphy, Chapter 14, Cambridge University Press, p. 631-712.
(Standard work on Cenozoic radiolaria and zonations; more recent textbook on radiolaria by De Wever, Dumitrica et al. 2001. See also more recent milestone reviews by O'Dogherty (2009, 2021) and Baumgartner))

Scheffen, W. (1932)- Zur Lepidocyclinen-Bestimmung. De Mijningenieur 13, p. 97-99.
('On the identification of Lepidocyclina species' (Tertiary larger foraminifera))

Scheffen, W. (1932)- Zwei charakteristische lageniden von Java. Wetenschappelijke Mededeelingen, Dienst Mijnbouw Nederlandsch-Indie 19, p. 195-198.
(Two new Lagena-type smaller benthic foraminifera from W Java Miocene. One of them probably Vaginulopsis)

Scheffen, W. (1932)- Ostindische Lepidocyclinen, Teil I. Wetenschappelijke Mededeelingen, Dienst Mijnbouw Nederlandsch-Indie 21, p. 4-76.
('East Indies Lepidocyclinids'. Paleontological paper describing dozen new 'species' and varieties of Lepidocyclina. Little or nothing on localities, stratigraphic significance, etc.)

Scheffen, W. (1932)- Zur Morphologie und Morphogenese der 'Lepidocyclinen'. Palaontologische Zeitschrift 14, 4, p. 233-256.
('Morphology and morphogenesis of the Lepidocyclinids')

Schlumberger, C. (1896)- Note sur le genre *Tinoporus*. Memoires Societe Zoologique France 1896, 9, p. 87-90.
(Description of Recent species Baculogypsina floresiana n.sp. from S coast of Flores, and comparison with Baculogypsina sphaerulata, common in Indian and Pacific Oceans)

Schweitzer, H.J. (1958)- Die fossilen Dipterocarpaceen Holzer. Palaeontographica, B, 105, p. 1-66.
('The fossil dipterocarp woods'. Includes descriptions of new species of petrified wood from Miocene-Quaternary of Sumatra (Jambi, Palembang), Java (Bogor, Banten), E Kalimantan (Dryobalanoxylon musperi, D. bangkoense, D. borneense, D. sumatrense, etc.))

Serra-Kiel, J., L. Hottinger, E. Caus, K. Drobne, C. Fernandez, A.K. Jauhri, G. Less, R. Pavlovec, J. Pignatti et al. (1998)- Larger foraminiferal biostratigraphy of the Tethyan Paleocene and Eocene. Bull. Societe Geologique France 169, 2, p. 281-299.
(online at: https://www.researchgate.net/profile/Gyoergy-Less/publication/241772566_tethyan_Paleocene-Eocene_Larger_Foraminifera_Biostratigraphy_Shallow_Benthic_Zones_SBZ_Etc.))
(New standard zonation of Paleogene larger foraminifera. 20 Shallow Benthic Zones (SBZ) in Paleo-Eocene, based on alveolinids, nummulitids and orthophragminids. Correlation with Berggren et al. (1995) time scale based on magnetostratigraphic data from Pyrenean Basin and correlation with nannoplankton and planktonic foraminifera zonations in E and C parts of (Neo-)Tethys)

Shafik, S. & G.H. Chaproniere (1978)- Nannofossil and planktic foraminiferal biostratigraphy around the Oligocene-Miocene boundary in parts of the Indo-Pacific region. BMR J. Australian Geology Geophysics 3, p. 135-151.
(online at: www.ga.gov.au/corporate_data/80954/Jou1978_v3_n2_p135.pdf)

(Based on study of Ashmore Reef 1 (Australia NW shelf), outcrop material from 'Meleri Beds' of W Papua and Capricorn 1 well (off Queensland) a.o. find overlapping ranges of Globigerinoides primordius and Globorotalia opima in combined zone N3/N4. Globigerinoides first appearance datum is in Late Oligocene)

Sharma, V. & L.B. Devi (2007)- Neogene oceanographic and climatic changes in the northern Indian Ocean: evidence from radiolaria. Indian J. Marine Science 36, 4, p. 361-368.
(online at: nopr.niscair.res.in/bitstream/123456789/...)
(E Neogene radiolaria from N Indian Ocean and Andaman-Nicobar islands show influx of colder water immigrant radiolaria species in early Middle Miocene (~12.5- 16 Ma))

Sharma, V. & M.P. Ram (2003)- Early to Middle Miocene radiolarian assemblages and biostratigraphy, Andaman Islands, Northeast Indian Ocean. J. Palaeontological Society India 48, p. 1-39.
(online at: http://palaeontologicalsociety.in/vol48/v1.pdf)
(Ten low-latitude radiolarian events identified in E-M Miocene sections of Andaman- Nicobar islands)

Shin, C.P., M. Yasuhara, H. Iwatani, T. Kase, A.G.S. Fernando, H. Hayashi, Y. Kurihara & H. Pandita (2019)- Neogene marine ostracod diversity and faunal composition in Java, Indonesia: Indo-Australian Archipelago biodiversity hotspot and the Pliocene diversity jump. J. Crustacean Biology 39, 3, p. 244-252.
(online at: https://academic.oup.com/jcb/article/39/3/244/5304204)
(M Miocene- E Pliocene marine ostracods from W and C Java significant diversity increase from Late Miocene to Pliocene, supporting 'Pliocene Diversity Jump' hypothesis (but also different paleoenvironments?; JTvG))

Shuto, T. (1975)- Preliminary correlation of the Neogene molluscan faunas in Southeast Asia. In: T. Kobayashi & R. Toriyama (eds.) Geology and Palaeontology of Southeast Asia, University of Tokyo Press, 15, p. 289-301.

Shuto, T. (1977)- Correlation of Neogene formation of Southeast and South Asia by means of molluscan faunas. In: T. Saito & H. Ujiie (eds.) Proc. First International Congress on Pacific Neogene Stratigraphy, Tokyo 1976, p. 133-144.

Shuto, T. (1978)- On the genera *Siphonofusus* and *Euthria* of the Indo- West Pacific. Transactions Proceedings Paleontological Society of Japan, N.S., 111, p. 358-369.
(online at: www.palaeo-soc-japan.jp/download/TPPSJ/TPPSJ_NS111.pdf)
(On gastropod genera Buccinulum (E Miocene and younger), Siphonofusus (M Miocene and younger) and Euthria (Eocene and younger). With descriptions of species from Java, Kalimantan)

Shuto, T. (1978)- Molluscan biohorizons in the Indo- West Pacific Neogene. Proc. 2nd Working Group Meeeting Biostrat Datum Planes in the Pacific, Bandung 1977, IGCP Project 114, Geological Research Development Centre (GRDC), Bandung, Special Publ. 1, p. 99-110.

Shuto, T. (1984)- A revision of the Burmese Tertiary turrids. Memoirs Faculty of Science, Kyushu University, ser. D (Geology), 25, 2, p. 115-157.
(online at: https://catalog.lib.kyushu-u.ac.jp/opac_download_md/1546094/25_02_p115.pdf)
(Revision of Burmese turrid gastropods, mainly based on material from Vredenburg (1921). No figures)

Siemon, F. (1929)- Jungtertiare Molluskenfauna aus Niederlandisch Ost-Indien. Berichte Naturforschenden Gesellschaft Freiburg 29, p. 309-369.
(online at: www.zobodat.at/pdf/Berichte-naturf-Ges-Freiburg-Br_29_0309-0369.pdf)
('Late Tertiary mollusc fauna from the Netherlands East Indies'. Mio-Pliocene? bivalves and gastropods from (1) S West Papua (Noordwest, Bibis and Noord Rivers, S of Central Range; from 1907 and 1909/1910 New Guinea expeditions) and (2) W Java (Cirebon area), collected by G. Kemmerling. Also shark tooth Carcharias gangeticus. One small plate)

Sieverts, H. (1934)- Fossile Brechiten (Aspergillen), besonders aus dem ostindischen Tertiar. Palaontologische Zeitschrift 16, p. 263-275.

('Fossil Brechites (Aspergillen), particularly from the East Indies Tertiary'. Descriptions of tube-dwelling molluscs, including fossils from Pliocene of Muna Island and Miocene of Brunei, collected by mining engineer A.C.D. Bothe)

Singh, R.K. & A.K. Gupta (2010)- Deep-sea benthic foraminiferal changes in the eastern Indian Ocean (ODP Hole 757B): their links to deep Indonesian (Pacific) flow and high latitude glaciation during the Neogene. Episodes 33, 2, p. 74-81.
(online at: www.episodes.org/index.php/epi/article/download/62122/48423)
(ODP Hole 757B, Ninetyeast Ridge, SE Indian Ocean, with well-oxygenated deep-sea benthic foraminifera (Cibicides cicatricosus, C. pseudoungerianus, Oridorsalis umbonatus) dominant in late Oligocene- E Miocene, but declining through M Miocene as Site 757 became under influence of Indonesian Throughflow with water masses from Pacific Ocean. Nuttallides umbonifera major increase at ~11 Ma, coinciding with increase in Nd isotope values, indicating substantial transport of deep Pacific water to Indian Ocean through Indonesian seaway. N. umbonifera decreases drastically during 3-2.8 Ma, coinciding with closure of Indonesian seaway and switch in shallow ITF source from warm, saline S Pacific to cool, fresh N Pacific thermocline water)

Skwarko, S.K. (1994)- Mollusca in Indonesian Cenozoic biostratigraphy (a computerised compilation), 1. Introductory chapters, Amphineura, Cephalopoda, Scaphopoda. Geological Research Development Centre (GRDC), Bandung, p. 1-125.

Skwarko, S.K. & E. Sufiati (1994)- Mollusca in Indonesian Cenozoic biostratigraphy (a computerised compilation), 3. Gastropoda. Geological Research Development Centre (GRDC), Paleont. Section, p. 1-796.

Skwarko, S.K., E. Sufiati & A. Limbong (1994)- Mollusca in Indonesian Cenozoic biostratigraphy (a computerised compilation), 2. Bivalvia. Geological Research Development Centre (GRDC), Paleont. Section, p. 1-326.

Srinivasan, M.S. & S.N. Chaturvedi (1992)- Neogene planktonic foraminiferal biochronology of the DSDP sites along the Ninetyeast Ridge, Northern Indian Ocean. In: K. Ishizaki & T. Saito (eds.) Centenary of Japanese micropaleontology, Terra Scientific Publishing, Tokyo, p. 175-188.
(online at: http://svr4.terrapub.co.jp/e-library/cjm/pdf/0175.pdf)
(Neogene planktonic foram biostratigraphy of Ninety East Ridge, N Indian Ocean, DSDP sites. Major epoch boundaries marked: by Top Globigerinoides fistulosus (Plio-Pleistocene), Base Globorotalia tumida (base Pliocene), Base Globoquadrina dehiscens (Base Miocene). Top Globoquadrina binaiensis is useful marker close to N5-N6 boundary in tropical Indian Ocean and Indo-Pacific region. Absence of Pulleniatina spectabilis suggests effective closing of Indonesian Seaway in M Miocene, etc.)

Srinivasan, M.S. & J.P. Kennett (1981)- A review of Neogene planktonic foraminiferal biostratigraphy applications in the equatorial and South Pacific. In: J.E. Warme et al. (eds.) The Deep Sea Drilling Project, a decade of progress, Society for Sedimentary Geology (SEPM) Special Publ. 32, p. 395-432.

Srinivasan, M.S. & D.K. Sinha (1992)- Late Neogene planktonic foraminiferal events of the Southwest Pacific and Indian Ocean: a comparison. In: R. Tsuchi & J. Ingle (eds.) Pacific Neogene: environment, evolution and events. University of Tokyo Press, Tokyo, p. 203-220.

Stainforth, R.M., J.L. Lamb, H.P. Luterbacher, J.H. Beard & R.M. Jeffords (1975)- Cenozoic planktonic foraminifera zonation and characteristics of index forms. University of Kansas Paleontological Contributions 62, p. 1-425.
(Classic mid-1970s work on Paleogene-Neogene planktonic foraminifera)

Suhartati, M.N. (1984)- Biometri statistik rasio panjang lebar pada genus *Asterorotalia* di sumur bor Cengkareng. J. RISET Geologi dan Pertambangan (LIPI) 5, 2, p. 17-24.
(online at: http://pustaka.geotek.lipi.go.id/wp-content/uploads/2016/02/Riset-Vol.5-No.2-2.pdf)

('Statistical biometry of the length ratio in the genus Asterorotalia in wellbore Cengkareng'. Length-width ratios of 3 Asterorotalia species (A. inspinosa, A.multispinosa, A. trispinosa) in 200m deep Cengkareng well, NW Java)

Takayama, T. (1984)- Coccolith biostratigraphy in Southeast Asia. In: T. Kobayashi, R. Toriyama & W. Hashimoto (eds.) Geology and Palaeontology of Southeast Asia, University of Tokyo Press, 25, p. 303-304.
(Very brief review of literature on calcareous nannoplankton in SE Asia. Very little work done in Indonesia)

Tan Sin Hok (1927)- Over de samenstelling en het ontstaan van krijt- en mergel-gesteenten van de Molukken. Jaarboek Mijnwezen Nederlandsch-Indie 55 (1926), Verhandelingen 3, p. 5-165. *(also Ph.D. Thesis, Technische Hogeschool Delft, p. 1-165)*
(online at: https://www.delpher.nl/nl/boeken/view?identifier=MMKB24:063568000:00001)
('On the composition and origin of chalks and marls of the Moluccas'. Pioneering study of radiolarians and calcareous nannoplankton (Discoaster) from deep water sediments of Timor, Roti, Yamdena, Halmahera, etc. Very little stratigraphic context of samples. ((N.B. Radiolaria described from Roti are not of Late Tertiary age as assumed by TSH, but are of Early Cretaceous age (e.g. Eucyrtidium (now Archaeodictyomitra) brouweri; Baumgartner 1992, Jasin & Haile 1996, O'Dogherty 2009). Many of the new 'Late Tertiary' radiolarian species from Roti described by TSH are also present in Early Cretaceous of SW Sulawesi; Munasri 2013))

Tan Sin Hok (1927)- Discoasteridae Incertae Sedis. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 30, 3, p. 411-419.
(online at: https://dwc.knaw.nl/DL/publications/PU00015461.pdf)
(First publication on 'aragonite stars' (calcareous nannofossils/coccoliths) from E Indonesia, including new Late Pliocene species of Discoaster: D. brouweri, D. molengraaffi, D. pentaradiatus, etc.)

Tan Sin Hok (1930)- Enkele opmerkingen over de stratigraphische verspreiding van *Trybliolepidina* v.d. Vlerk. De Mijningenieur 11, p. 144-146.
('Some remarks on the stratigraphic distibution of Trybliolepidina'. This most evolved Lepidocyclina subgenus regarded by Van der Vlerk as limited to upper Tertiary f, but questioned by TSH)

Tan Sin Hok (1930)- Over *Spiroclypeus* met opmerkingen over zijn stratigraphische verspreiding. De Mijningenieur 11, 9, p. 180-184.
('On Spiroclypeus and its stratigraphic distribution'. Larger foram Spiroclypeus evolved from Heterostegina in middle part of Tertiary e= Late Oligocene)

Tan Sin Hok (1930)- On *Cycloclypeus* its phylogeny and signification for the biostratigraphy in general and for the stratigraphy of the Tertiary of the Indo-Pacific region. Handelingen 6e Nederlandsch-Indisch Natuurwetenschappelijk Congres, Bandung,1930, p. 641-644.

Tan Sin Hok (1930)- Over *Cycloclypeus*: voorlopige resultaten eener biostratigraphische studie. De Mijningenieur 11, 12, p. 233-242.
('On Cycloclypeus: preliminary results of a biostratigraphic study'. Larger foram genus known from Early Oligocene- Recent. Reported in more detail in 1932))

Tan Sin Hok (1931)- Discoasteridae, Coccolithinae and Radiolaria. In: B.G. Escher et al. (eds.) De palaeontologie en stratigraphie van Nederlandsch Oost-Indie, Feestbundel K. Martin, Leidsche Geologische Mededelingen 5, p. 92-114.
(Listings of calcareous nannoplanton and radiolaria species reported from the Indonesian region by 1931)

Tan Sin Hok (1932)- On the genus *Cycloclypeus* Carpenter, Part 1 and an appendix on the Heterostegines of Tjimanggoe, S. Bantam, Java. Wetenschappelijke Mededeelingen Dienst Mijnbouw Nederlandsch-Indie, 19, p. 1-194.
('About the genus Cycloclypeus Carpenter, part 1 with an appendix on the heterosteginids of Cimanggu, South Banten, Java'. Classic paper on evolution and species of Early Oligocene- Recent Cycloclypeus in West Java)

Tan Sin Hok (1935)- Die peri-embryonalen Aquatorialkammern bei einigen Orbitoiden. De Ingenieur in Nederlandsch-Indie (IV, Mijnbouw en Geologie) 2, 12, p. IV.113- IV.126.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1935-12.pdf)
('The peri-embryonal chambers in some orbtoidal foraminifera'. General discussion of initial chamber morphologies in Cretaceous and Tertiary orbitoidal foraminifera (not focused on Indonesian material))

Tan Sin Hok (1936)- Beitrag zur Kenntnis der Lepidocycliniden. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 39, 8, p. 990-999.
('Contribution to the knowledge of the Lepidocyclinids'. First (Eocene) Polylepidina-type Lepidocyclina from Indo-Pacific, from tributary of Mahakam River, E Borneo)

Tan Sin Hok (1936)- Zur Kenntnis der Lepidocycliniden. Natuurkundig Tijdschrift voor Nederlandsch-Indie 96, p. 235-280.
('About the knowledge of the Lepidocyclinids'. Mainly critical review of Barker & Grimsdale (1936) paper on American lepidocyclinids. No figures or data)

Tan Sin Hok (1936)- Zur Kenntnis der Miogypsiniden. De Ingenieur in Nederlandsch-Indie (IV) 3, 3, p. 45-61.
('On the knowledge of Miogypsinids'. First of series of five papers on miogypsinid evolution and stages/species in Indonesia. Miogypsinids probably evolved from Rotalia. Comprehensive dission of the five main stages in Late Oligocene-Middle Miocene miogypsinids: Miogypsinoides complanata, Miogypsina borneensis, M. ecuadorensis, M. indonesiensis and M. bifida. Mainly based on samples from East Kalimantan)

Tan Sin Hok (1936)- Zur Kenntnis der Miogypsiniden. I Fortzetsung. De Ingenieur in Nederlandsch-Indie (IV) 3, 5, p. IV.84- IV.98.
('On the knowledge of the Miogypsinids- First continuation'. Discussion of more 'obscure' Miogypsina species and details of chamber patterns and stolons)

Tan Sin Hok (1936)- Zur Kenntnis der Miogypsiniden. II Fortzetsung und Schlusz. De Ingenieur in Nederlandsch-Indie (IV) 3, 7, p. IV.109- IV.123.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1936-07.pdf)
('On the knowledge of the Miogypsinids- Second continuation and End'. Discussion of growth patterns of miogypsinid and other larger forams and remarks on stratigraphic distribution and interregional correlations. No illustrations)

Tan Sin Hok (1936)- Over verschillende paleontologische criteria voor de geleding van het Tertiair. De Ingenieur in Nederlandsch-Indie (IV) 3, 9, p. IV.173- IV.179.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1935-09.pdf
('On the different paleontological criteria for the subdivision of the Tertiary'. Discussion of value of various larger foram genera for biostratigraphic subdivision of the Tertiary. Proposes to use the new genus name Vacuolispira for the Pellatispira inflata Umbgrove 1928 group)

Tan Sin Hok (1937)- Note on *Miogypsina kotoi* Hanzawa. De Ingenieur in Nederlandsch-Indie (IV) 4, 2, p. IV.31-IV.33.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1937-02.pdf)
(Description of material of Miocene Miogypsina kotoi Hanzawa from Honsyu island, Japan)

Tan Sin Hok (1937)- Weitere Untersuchungen uber die Miogypsiniden I. De Ingenieur in Nederlandsch-Indie (IV) 4, 3, p. IV.35- IV.45.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1937-03.pdf)
('Further investigations on the Miogypsinids- I'. Miogypsinid population from the late Aquitanian of Madura. Contains Miogypsina kotoi, M. kotoi bispiralis, M. thecidaeformis, M. aff. Polymorpha spiralis and Miolepidocyclina excentrica n.sp. (with excellent illustrations; JTvG))

Tan Sin Hok (1937)- Weitere Untersuchungen uber die Miogypsiniden II. De Ingenieur in Nederlandsch-Indie (IV) 4, 6, p. IV.87- IV.111.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1937-06.pdf)
('Further investigations on the Miogypsinids- II'. Mainly on Miogypsina indonesiensis group, an advanced member of the Miogypsina lineage, (Tf; early M Miocene), here reclassified as subspecies of M. cushmani)

Tan Sin Hok (1937)- On the genus *Spiroclypeus* Douville with a description of the Eocene *Spiroclypeus vermicularis* nov. sp. from Koetai in East Borneo. De Ingenieur in Nederlandsch-Indie (IV) 4, 10, p. 177-193.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1937-10.pdf)
(Review of larger foram genus Spiroclypeus. Stratigraphic range of Spiroclypeus not only Late Oligocene- E Miocene (zone Upper Te), but also in Late Eocene (Tb). On p. 179: mention of Biplanispira in Wani series of Buton)

Tan Sin Hok (1939)- The results of phylomorphogenetic studies of some larger foraminifera (a review). De Ingenieur in Nederlandsch-Indie (IV) 6, 7, p. IV.93- IV.97.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1939-07.pdf)
(Brief general review on evolutionary studies on larger foraminifera lineages, incl.the Cycloclypeus, Miogypsina and Lepidocyclina groups)

Tan Sin Hok (1939)- Remarks on the "Letter classification" of the East Indian Tertiary. De Ingenieur in Nederlandsch-Indie (IV) 6, 7, p. IV.98- IV.101.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1939-07.pdf)
(Brief paper with comments on larger foram 'letter zonation'. Miogypsinoides appears in Late Oligocene, etc.)

Termier G. & A.F. Poignant (1982)- Une symbiose algue rouge-spongiaire dans le Miocene inferieur de l'Indonesie. Comptes Rendus hebdomadaires Seances de l'Academie Sciences Paris, Serie 2, 294, D, p. 349-353.
(online at: https://gallica.bnf.fr/ark:/12148/bpt6k6327319s/f363.item.r=indonesie)
('A red algae-sponge symbiosis from the Lower Miocene of Indonesia'. Basal Miocene from offshore West Java (= Baturaja Fm?; JTvG) with encrusting calcareous red algae Lithophyllum johnsoni and Solenomeris sp., in apparent symbiotic relationship with sponge))

Thalmann, H.E. (1933)- Zwei neue Vertreter der Foraminiferen-Gattung *Rotalia* Lamarck 1804: *R. cubana* nom. nov. und *R. trispinosa* nom. nov. Eclogae Geologicae Helvetiae 26, 2, p. 248-251.
(online at: https://www.e-periodica.ch/digbib/view?pid=egh-001:1933:26#301)
('Two new representatives of the genus Rotalia Lamarck: R. cubana new name and R. trispinosa'. Short paper. New name Rotalia trispinosa for Rotalia pulchella, described by Brady (1884) from Bangka Straits)

Thalmann, H.E. (1934)- Mitteilungen uber Foraminiferen I. Eclogae Geologicae Helvetiae 27, 2, p. 428-440.
(online at: http://retro.seals.ch/cntmng?type=pdf&rid=egh-001:1934:27::628&subp=hires)
('Communications on foraminifera- I'. Brief review of Miocene- Pleistocene Pseudorotalia species from Indonesia. Includes chapters 1 on Rotalia gaimardi, 2 on Rotalia conoides from Cepu area, E Java, and 4 on two new species from Plio-Pleistocene of Java, Rotalia catilliformis and Rotalia alveiformis)

Thalmann, H.E. (1935)- Mitteilungen uber Foraminiferen II. Eclogae Geologicae Helvetiae 28, 2, p. 592-606.
(online at: https://www.e-periodica.ch/digbib/view?pid=egh-001%3A1935%3A28%3A%3A667)
('Communications on foraminifera- II'. Includes chapter 8, brief description of Rotalia Pseudorotalia indopacifica n.sp. from Late Tertiary and E Quaternary of N Java (no figures))

Thalmann, H.E. (1937)- Mitteilungen uber Foraminiferen III. Eclogae Geologicae Helvetiae 30, 2, p. 337-356.
(online at: http://retro.seals.ch/cntmng?type=pdf&rid=egh-001:1937:30::563&subp=hires)
('Communications on foraminifera- III'. Nothing on SE Asia)

Thalmann, H.E. (1938)- Wert und Bedeutung morphogenetischer Untersuchungen an Gross-Foraminiferen fur die Stratigraphie. Eclogae Geologicae Helvetiae 31, 2, p. 333-337.
(online at: http://retro.seals.ch/digbib/view?pid=egh-001:1938:31::352)
('Value and meaning of morphogenetic investigations on larger foraminifera for stratigraphy'. Mainly a brief review in the importance of work by Tan Sin Hok (1932-1937) on the 'morphogenetic' evolution of Cenozoic larger foraminifera Lepidocyclina, Spiroclypeus, Miogypsina, Cycloclypeus, etc. from Indonesia)

Thalmann, H.E. (1942)- Occurrence of the genus *Lacazina* Munier-Chalmas in the East Indies. Geological Society of America (GSA) Bull. 53, 12, p. 1838-1839. *(Abstract only)*
(Eocene limestones with larger foram Lacazina (= Lacazinella) often associated with Nummulites, Discocyclina and Alveolina. Known only from E half of East Indies Archipelago: E Sulawesi (Boealemo Peninsula), W Papua (Pisang Island E of Misool; Onin Island, Dramai Island; S and E of Triton Bay; Setawka River in SW New Guinea; Birds Head between Rumberpon and Horna, and Sungei Ingsiim; on Wilhelmina Peak of Central Range; S and E of Paniai Lake), PNG (Chimu aerodrome), Kai Besar Island (between Riamroe and Yamtimur))

Thalmann, H.E. (1946)- New occurrences of the foraminiferal genus *Hantkenina* in Europe and Asia. Geological Society of America (GSA) Bull. 57, 12, 2, p. 1236-1237. *(Meeting abstract only)*
(Incl. presence of Eocene planktonic foraminifera of genus Hantkenina in E Kalimantan (many localities along upper reaches and tributaries of Mahakam River; Long Iram area), and C Seram (Wai Tali, Cape Pasanea; Germeraad, 1946))

Theodoridis, S. (1983)- On the legitimacy of the generic name *Discoaster* Tan Sin Hok, 1927 ex Tan Sin Hok, 1931. Int. Nannoplankton Assoc. (INA) Newsletter 5, 1, p. 15-21.
(online at: http://ina.tmsoc.org/JNR/NINA/INANewsl5(1).pdf)
(Commonly used genus name Discoaster is technically invalid because at the time of first description no type species was designated by Tan Sin Hok (instead spp. assigned to 'subgenera' Eu-Discoaster and Helio-Discoaster)

Theodoridis, S. (1984)- Calcareous nannofossil biozonation of the Miocene and revision of the Helicoliths and Discoasters. Utrecht Micropaleontological Bulletin 32, p. 1-271.
(online at: http://dspace.library.uu.nl/handle/1874/205891)
(Miocene nannofossil zonation, partly based on Miocene material of Solo River section, Kendeng zone, E Java. Returns to emended definitions of Helio-discoaster Tan and Eu-discoaster Tan for Discoaster species)

Tobler, A. (1925)- Uber eine ostindische *Lepidocyclina* mit mehrkammeriger nucleoconch. Eclogae Geologicae Helvetiae 19, p. 269-274.
(online at: https://www.e-periodica.ch/digbib/view?pid=egh-001:1925-1926:19#284)
('On an East Indies Lepidocyclina with multi-chambered embryon'. Lepidocyclina (probably Eulepidina) from E Miocene at Sungi Tjengal, N margin of Gumai Mts, S Sumatra, with multi-chambered embryon (not unusual aberrant growth in orbitoidal foraminifera, with no apparent ecological or biostratigraphic significance; JTvG))

Tobler, A. (1927)- Verkalkung der Lateralkammern bei *Miogypsina*. Eclogae Geologicae Helvetiae 20, 2, p. 323-330.
(online at: https://www.e-periodica.ch/digbib/view?pid=egh-001%3A1926%3A20%3A%3A338)
('Calcification of the lateral chambers in Miogypsina'. Incl. new species Miogypsina tuberosan.sp. from Popoh in Kediri Residency in SE Java (Coll. W. Hotz) and M. abunensis n.sp. from limestone in Sungei Abun in Sarawak (Coll. M. Muhlberg)

Umbgrove, J.H.F. (1928)- Het genus *Pellatispira* in het Indo-Pacifische gebied. Dienst Mijnbouw Nederlandsch-Indie, Wetenschappelijke Mededeelingen 10, p. 43-71.
(online at: https://resolver.kb.nl/resolve?urn=MMKB24:065531000:pdf)
('The genus Pellatispira in the Indo-Pacific area'. Review of Late Eocene (Ta-Tb) larger foram genus Pellatispira in Indonesian region. Seven species, five of which new)

Umbgrove, J.H.F. (1930)- Tertiary sea connections between Europe and the Indo-Pacific area. Proc. Fourth Pacific Science Congress, Java 1929, vol. IIA, p. 91-104.
(Describing difficulties in Indonesia- Europe biostratigraphic correlations due to faunal provincialism)

Umbgrove, J.H.F. (1931)- Tertiary foraminifera. In: B.G. Escher et al. (eds.) De palaeontologie en stratigraphie van Nederlandsch Oost-Indie, Feestbundel K. Martin, Leidsche Geologische Mededelingen 5, p. 36-91.
(online at: http://dare.uva.nl/cgi/arno/show.cgi?fid=549641)
(Chapter in K. Martin memorial volume, listing >640 foraminifera species reported from Tertiary of Indonesia)

Umbgrove, J.H.F. (1954)- Tertiary stratigraphy of the Far East; a protest. American Journal of Science 252, p. 503-504.
(online at: https://ajsonline.org/article/58514.pdf)
(Umbgrove objects to use in a publication of the 'East Indies Letter Classification' without properly referencing the main person who first developed it in 1925 (subsequent refinements followed by Van der Vlerk & Umbgrove (1927) and Leupold and Van der Vlerk (1931). No detail data)

Van Cappelle, H. (1885)- Het karakter van de Nederlandsch--Indische Tertiaire fauna. Doct. Thesis Rijksuniversiteit Leiden, p. 1-198.
('The nature of the Tertiary fauna of the Netherlands Indies'. Early interpretation of nature of Indonesian Tertiary faunas by student of K. Martin, with comparisons of Recent and Tertiary fossil faunas)

Van der Vlerk, I.M. (1923)- Een overgangsvorm tusschen *Orthophragmina* en *Lepidocyclina* uit het Tertair van Java. Verhandelingen Geologisch-Mijnbouwkundig Genootschap Nederland Kolonien, Geologische Serie 7, 2, p. 91-98.
('A transitional form between Orthophragmina and Lepidocyclina from the Tertiary of Java'. Description of new genus and species Orthocyclina soeroeanensis from Kali Soeroean, Bagelen area, C Java. Looks like an advanced M-L Miocene radiate Lepidocyclina (Trybliolepidina). Names never used by later workers; JTvG)

Van der Vlerk, I.M. (1924)- *Miogypsina Dehaartii* nov. spec. de Larat (Moluques). Eclogae Geologicae Helvetiae 18, p. 429-431.
(online at: http://retro.seals.ch/cntmng?type=pdf&rid=egh-001:1923-1924:18::764&subp=hires)
(New miogypsinid larger foram species from Larat, an island off SW coast of New Guinea, collected by BPM geologist De Haart. No locality info. No lateral chambers, so should be assigned to genus Miogypsinoides. See also Van der Vlerk 1966)

Van der Vlerk, I.M. (1924)- De verspreiding van het foraminiferengeslacht *Lepidocyclina* en haar beteekenis voor de palaeogeographie. Handelingen 3e Nederlandsch-Indisch Natuurwetenschappelijk Congres, Buitenzorg 1924, p. 371-380.
('The distribution of the foraminiferal genus Lepidocyclina and its significance for paleogeography')

Van der Vlerk, I.M. (1925)- Het foraminiferen genus *Spiroclypeus* en zijn beteekenis voor de stratigraphie van het Tertiair van den Indo-Australischen Archipel. Verhandelingen Geologisch-Mijnbouwkundig Genootschap Nederland Kolonien, Geologische Serie 8 (Gedenkboek Verbeek, memorial volume), p. 561-568.
(online at: https://books.google.com/books?id=Yy0RAAAAIAAJ&pg)
('The larger foram genus Spiroclypeus and its significance for the stratigraphy of the Tertiary of the Indo-Australian Archipelago')

Van der Vlerk, I.M. (1928)- Het genus *Lepidocyclina* in het Indo-Pacifische gebied. Wetenschappelijke Mededeelingen Dienst Mijnbouw Nederlandsch-Indie, 8, p. 7-86.
(Classification of Indo-Pacific Lepidocyclina, primarily based on characteristics of embryon: Eulepidina, Trybliolepidina, Nephrolepidina, Isolepidina and Pliolepidina. Incl. Lepidocyclina boetonensis n.sp. from Eo-Oligocene of Buton)

Van der Vlerk, I.M. (1928)- The genus *Lepidocyclina* in the Far East. Eclogae Geologicae Helvetiae 21, 1, p. 182-211.
(online at: http://retro.seals.ch/cntmng?type=pdf&rid=egh-001:1928:21::604&subp=hires)
(Early review of Oligo-Miocene larger foram genus Lepidocyclina. With species determination table)

Van der Vlerk, I.M. (1931)- Cenozoic Amphineura, Gastropoda, lamellibranchiata, Scaphopoda. In: B.G. Escher et al. (eds.) De palaeontologie en stratigraphie van Nederlandsch Oost-Indie, Leidsche Geologische Mededelingen 5 (K. Martin memorial volume), p. 206-296.
(online at: https://repository.naturalis.nl/pub/505678)
(Listing of all (100s) Cenozoic mollusc species described from Indonesia before 1931, mainly by K. Martin)

Van der Vlerk, I.M. (1950)- Stratigraphy of the Caenozoic of the East Indies based on foraminifera. Report 18th International Geological Congress, Great Britain 1948, 15, p. 61-63.
(Summary of Tertiary larger foram Ta-Tg 'Letter zonation' used in shallow marine carbonates of Indonesia)

Van der Vlerk, I.M. (1951)- Tabulation of determinations of larger foraminifera. In: M. Reinhard & E. Wenk (eds.) Geology of the Colony of North Borneo, Bull. Geological Survey Department, British Territories in Borneo 1, p. 137-145.
(Incl. samples from Banggi and Kudat areas of E Miocene limestones with reworked Eocene Pellatispira, Discocyclina, etc.)

Van der Vlerk, I.M. (1955)- Correlation of the Tertiary of the Far East and Europe. Micropaleontology 1, p. 72-75.
(One of the first and basically correct, correlations of the 'Far East Letter Classification' (Cenozoic larger foraminifera zonation) with successions in Pakistan, Middle East and France)

Van der Vlerk, I.M. (1959)- Problems and principles of Tertiary and Quaternary stratigraphy. Quarterly J. Geological Society, London, 115, p. 49-63.

Van der Vlerk, I.M. (1961)- *Lepidocyclina radiata* (K. Martin), 1880. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B64, 5, p. 620-626.
(Description of type specimen of M-L Miocene Lepidocyclina radiata from south coast of W Java)

Van der Vlerk, I.M. (1963)- Biometric research on *Lepidocyclina*. Micropaleontology 9, p. 425-426.
(On evolutionary tendency in iso-, nephro- and trybliolepidine embryons in Lepidocyclina embryon of deuteroconch to enclose protoconch to an increasing degree. Measured as 'grade of enclosure, maximum values increase from 25-30% in Late Eocene, 40-45% in Late Oligocene, 45-50% at Te-Tf boundary and 50-55% in E Miocene/ Burdigalian)

Van der Vlerk, I.M. (1966)- Stratigraphie du Tertiare des domaines Indo-Pacifiques et Mesogeen (essai de correlation). Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B69, 3, p. 336-344.
('Stratigraphy of the Tertiary of the Indo-Pacific and Mesogean domains (attempt of correlation)'. Stratigraphic correlations between Far East and Europe using evolutionary stages of Lepidocyclina)

Van der Vlerk, I.M. (1968)- Evolutie van een embryo. Geologie en Mijnbouw 47, 2, p. 121-125.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0TklUS0F0Zkt5T0k/view)
('Evolution of an embryon'. Evolution of Late Oligocene- Miocene Lepidocyclina embryon can be measured as 'degree of curvature', which can then be used for age determination and correlations. With example of values in East Java- Madura stratigraphy)

Van der Vlerk, I.M. (1968)- Two methods of worldwide correlation. Micropaleontology 14, 3, p. 334-338.
('Degree of curvature' preferred over 'Grade of enclosure' method to characterize evolutionary stage of Lepidocyclina)

Van der Vlerk, I.M. (1973)- Nomenclature and numerical taxonomy (name and number). Verhandelingen Naturforschenden Gesellschaft Basel 84, 1 (Kugler memorial Volume), p. 245-255.
(On the different ways of classifying Lepidocyclina)

Van der Vlerk, I.M. (1973)- An improved method of biometric research. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B76, 4, p. 245-259.

Van der Vlerk, I.M. & D.D. Bannink (1969)- Biometrical investigations on *Operculina*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 72, 3, p. 169-174.
(Includes measurements on Operculina foram material from SE Asia. Observed gradual decrease in grade of enclosure of second chamber by third in specimens from Eocene- Recent)

Van der Vlerk, I.M. & R.E. Dickerson (1927)- Distinctions among certain genera of larger foraminifera for the field geologist of the East Indies. Journal of Paleontology 1, 3, p. 185-192.
(Brief paper on how larger foraminifera genera can be used by field geologists to determine broad ages of Tertiary limestones)

Van der Vlerk, I.M. & H. Gloor (1968)- Evolution of an embryo. Genetica 39, p. 45-63.
(Measurements on embryonic shambers of Lepidocyclina from Java and Madura. Extent to which deuteroconch encloses protoconch or extent to which dividing-wall between them is curved ("degree of curvature") increases from 10% in M Oligocene to 67% in M Miocene. Rate of evolution changes from very slow to very fast to slow again and to fast again. First rapid change in M Oligocene when genus migrated from America to Europe and to Far East- Australia. Second phase of rapid evolution from earliest Miocene up to extinction of genus)

Van der Vlerk, I.M. & J.H.F. Umbgrove (1927)- Tertiaire gidsforaminiferen van Nederlandsch Oost Indie. Wetenschappelijke Mededeelingen Dienst Mijnbouw Nederlandsch Oost-Indie 6, p. 1-45.
(online at: https://resolver.kb.nl/resolve?urn=MMKB24:067734000:pdf)
('Tertiary index fossils from the Netherlands East Indies'. Small guidebook describing and illustrating the main index larger foram genera in the Tertiary of Indonesia. The first paper to define a 'Letter Table' for subdivision of the Tertiary of the Netherlands Indies, with subdivisions a-f, based on vertical distribution of larger foraminifera genera. This would become known as the 'East Indies Letter Classification' (Adams, 1970))

Van Eijden, A.J.M. & J. Smit (1991)- Eastern Indian Ocean Cretaceous and Paleogene quantitative biostratigraphy. In: J. Weissel et al. (eds.) Proc. Ocean Drilling Program (ODP), Scientific Results, 121, p. 77-123.
(online at: www-odp.tamu.edu/publications/121_SR/VOLUME/CHAPTERS/sr121_03.pdf)
(Upper Cretaceous- Oligocene planktonic foram biostratigraphy of Indian Ocean ODP sites along Ninety-East Ridge, W and SW of Sumatra. Most intervals reflect temperate- subtropical climate)

Van Eijden, A.J.M. & G.M. Ganssen (1995)- An Oligocene multi-species foraminiferal oxygen and carbon isotope record from ODP Hole 758A (Indian Ocean): paleoceanographic and paleo-ecologic implications. Marine Micropaleontology 25, p. 47-65.
(Oligocene- basal Miocene (Zones P19-P22/N4) O and C isotope stratigraphy of benthic and plankonic forams from E Indian Ocean ODP Hole 758A. Lack of covariance in planktic and benthic δ18O ratios indicates that many Oligocene sea level fluctuations, including major fall at 30 Ma, not of glacio-eustatic origin)

Van Gorsel, J.T. (1981)- Paleoenvironmental distribution of Mid-Cretaceous to Recent larger foraminifera. Esso Production Research-European, Bordeaux, Exploration Memo EPR-E.1EX.81, p. 1-12. *(Unpublished)*

Van Gorsel, J.T. (1987)- Depositional sequence stratigraphy. Short Course, Indonesian Association of Geologists (IAGI), Jakarta, p. 1-180.
(Course notes for first 2-day sequence stratigraphy course in Indonesia)

Van Gorsel, J.T. (1988)- Biostratigraphy in Indonesia: methods, pitfalls and new directions. Proc. 17th Annual Conv. Indonesian Petroleum Association (IPA), Jakarta, 1, p. 275-300.
(online at: https://www.researchgate.net/publication/323775420_Biostratigraphy_in_Indonesia_methods_pitfalls_and_new_directions)
(Overview of methods, zonations, facies interpretations, etc., in Indonesian Tertiary biostratigraphy)

Van Gorsel, J.T. (2014)- An introduction to Cenozoic macrofossils of Indonesia. Berita Sedimentologi 30, p. 63-76.
(online at: https://journal.iagi.or.id/index.php/FOSI/article/view/137)
(Brief overview of principal groups and literature of Cenozoic macrofossils known from outcrops in Indonesia. Prior to 1930's macrofossils (mainly molluscs), were principal objects of paleontological and biostratigraphic studies in Indonesia. Since then focus shifted to microfossils. Cenozoic marine macrofossil assemblages from Indonesia all represent tropical faunas of Indo-Pacific province affinity)

Van Gorsel, J.T., P. Lunt & R. Morley (2014)- Introduction to Cenozoic biostratigraphy of Indonesia- SE Asia. Berita Sedimentologi 29, p. 6-40.
(online at: https://journal.iagi.or.id/index.php/FOSI/article/download/140/110)
Overview of Cenozoic biostratigraphy and biofacies interpretation of SE Asia. Principal microfossil groups used in region are foraminifera, calcareous nannofossils and palynology. Also brief reviews of work on ostracodes, diatoms and radiolaria)

Van Konijnenburg-van Cittert, J.H.A., I.M. van Waveren & J. Jonkers (2004)- Catalogue of the Mesozoic and Cenozoic holotypes in the collection of plant fossils in the Nationaal Natuurhistorisch Museum, Leiden. Nationaal Natuurhistorisch Museum (NNM) Technical Bulletin 7, p. 1-27.
(online at: www.repository.naturalis.nl/document/43244)
(Inventory of holotypes of fossil plants in Leiden Naturalis museum collections, mainly from Tertiary in Java, in 'Martin Collection'. Incl. Jurassic Novoguineoxylon lacunosum from Kamundan (W Papua), species from Java Mio-Pliocene described by Goeppert (1854), Crie (1888), Krausel) (1926, etc.)

Van Regteren Altena, C.O. (1938)- Renamed Mollusca from the Dutch East Indian Tertiary. Natuurkundig Tijdschrift voor Nederlandsch-Indie 98, p. 209-212.
(Listing of proposed nomenclatural changes for mollusc names described from Indonesia by K. Martin)

Van Straelen, V. (1931)- Arthropoda. In: B.G. Escher et al. (eds.) Onze palaeontologische kennis van Nederlandsch Oost Indie, Leidsche Geologische Mededelingen 5 (Feestbundel K. Martin), p. 156-163.
(online at: www.repository.naturalis.nl/document/549583)
(Listings of arthropod fossils from Indonesia, as known in 1931: (1) Permian trilobites (Phillipsia, Neoproetus indicus, Griffithides sumatrensis) from Timor, Leti, Sumatra, W Papua; (2) Neogene ostracodes, (3) Cenozoic crab fossils from Java, Kalimantan, Sulawesi, W Papua and M Jurassic Eryma boehmi n.sp. from of Jefbie, Misool)

Van Straelen, V. (1938)- Crustaces decapodes Cenozoiques des Indes Orientales neerlandaises. Leidsche Geologische Mededelingen 10, p. 90-103.
(online at: www.repository.naturalis.nl/document/549700)
('Cenozoic decapod crustaceans from the Netherlands East Indies'. Review of Neogene crab fossils from Neogene of Java (Calappa sangiranensis n.sp., Philyra, Ixoides, Cancer javanicus n.sp, etc.), Madura (Calappa madoerensis n. sp.), Eocene of Sumba (Ranina (Lophoranina) soembaensis n.sp.), etc.)

Van Vessem, E.J. (1977)- The internal structure of *Miogypsina polymorpha* and *Miogypsina bifida*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen B80, 5, p. 421-428.
(Study of two Miogypina (Lepidosemicyclina) species, originally described by Rutten (1911, 1912) from E Kalimantan, based on material from E Kalimantan and E Java. M. (L) polymorpha may be ancestor of M (L.) bifida))

Van Vessem, E.J. (1978)- Study of Lepidocyclinidae from Southeast Asia, particularly from Java and Borneo. Utrecht Micropaleontological Bulletin 19, p. 1-163.
(online at: https://dspace.library.uu.nl/handle/1874/205830)
(Quantitative study of lepidocyclinids of Nephrolepidina group from 42 samples from N Borneo (Klias Peninsula, Kinabatangan River), E Kalimantan (Kutei, Sangkulirang), C and E Java (Lodan, Tremboel, Boegoel wells) and Madura. Assemblages subdivided into five successive biometric units, called species: L. isolepidinoides, L. sumatrensis, L. angulosa, L. martini and L. rutteni)

Varol, O. (1983)- Proposed calcareous nannofossil zonation scheme for the Miocene to Holocene of Southeast Asia. Bull. Geological Society Malaysia 16, p. 37-46.
(online at: https://gsmpubl.files.wordpress.com/2014/09/bgsm1983004.pdf)
(Miocene- Recent SE Asia calcareous nannofossil zonation, modified from schemes of Martini (1971) and Okada & Bukry (1980))

Vermeij, G.J. & H. Raven (2009)- Southeast Asia as the birthplace of unusual traits: the Melongenidae (Gastropoda) of northwest Borneo. Contributions to Zoology 78, 3, p. 113-127.
(online at: http://dpc.uba.uva.nl/cgi/t/text/get-pdf?idno=m7803a03;c=ctz)
(On characteristics of large Melongena-type gastropods from Miocene-Recent of Sarawak- Brunei)

Vincent, E. (1977)- Indian Ocean Neogene planktonic foraminiferal biostratigraphy and its paleoceanographic implications. In: J. Heirtzler (ed.) Indian Ocean geology and biostratigraphy, American Geophysical Union (AGU) Memoir, Chapter 20, p. 469-584.
(Classic documentation of northern Indian Ocean Neogene planktonic foraminifera)

Wade, B.S., P.N. Pearson, W.A. Berggren & H. Palike (2011)- Review and revision of Cenozoic tropical planktonic foraminiferal biostratigraphy and calibration to the geomagnetic polarity and astronomical time scale. Earth-Science Reviews 104, p. 111-142.
(Modern review of calibration of planktonic foraminiferal datum levels to the geomagnetic polarity and astronomical time scale)

Wagner, C.W. (1964)- Manual of larger foraminifera. Bataafsche Internationale Petroleum Mij (BPM), The Hague, p. 1-307.
(Well-illustrated Shell manual of worldwide Mesozoic-Cenozoic and Paleozoiclarger foraminifera. Unpublished, but distributed among various academic institutions (although not always specifically mentioned, several of the illustrated genera are from the Indonesian region; JTvG)

Whipple, G.L. (1934)- Larger foraminifera from Vitulevu, Fiji. In: H.S. Ladd, Geology of Vitulevu, Fiji, Bernice P. Bishop Museum Bull. 119, p. 141-154.
(3 larger foram horizons on Fiji, all Te, Early Miocene)

Wright, J.D. & R.C. Thunell (1988)- Neogene planktonic foraminiferal biogeography and paleoceanography of the Indian Ocean. Micropaleontology 34, 3, p. 193-216.
(Indian Ocean planktonic foraminiferal biogeography examined in DSDP sites for 5 Neogene time-slices and used to reconstruct surface circulation patterns. Closure of Indo-Pacific passage at end of E Miocene strengthened low-latitude circulation in Indian Ocean and led to development of distinct tropical and subtropical faunal provinces. Global cooling during M-L Miocene resulted in steeper latitudinal temperature gradients and enhanced faunal provincialization)

Yabe, H. (1921)- Notes on some Eocene foraminifera, III. Notes on *Pellatispira* Boussac. Science Reports Tohoku Imperial University, 2nd ser. (Geol.), 5, p. 106-108.
(online at: https://tohoku.repo.nii.ac.jp/record/10618/files/KJ00004176256.pdf)
(General discussion of Eocene larger foram Pellatispira, partly based on material from Borneo and Japan)

Young, J.R. (1998)- Neogene. In: P.R. Bown (ed.) Calcareous nannofossil biostratigraphy, Chapter 8, British Micropalaeontological Society Publications Series, Chapman & Hall, p. 225-265.
(Thorough review of Miocene-Pliocene calcareous nannofossils and biozonations)

Zachariasse, W.J., W.A. Berggren, W. Si & B.T. Huber (2017)- On the taxonomic identity of the planktonic foraminiferal species *Globorotalia barisanensis* Le Roy, 1939. J. Foraminiferal Research 47, 4, p. 389-398.
(online at: https://pdfs.semanticscholar.org/fc9d/05e55c75424d1fd9590ca8ec7cbfe39d4a51.pdf)
(Holotype of Globorotalia barisanensis Le Roy 1939 from M Miocene Lower Palembang Fm in upper Kassikan section, Tapung Kiri River, C Sumatra, is non-keeled species, transitional between Gr. peripheroacuta and Gr praefohsi (zones N10-N11). Holotype should be viewed as senior synonym of Gr. peripheroacuta, but more practical to conserve latter name)

Zachariasse, W.J. & Sudijono (2012)- New data on the morphology and classification of the Oligocene-Miocene planktonic foraminifer *Paragloborotalia siakensis* (LeRoy, 1939). J. Foraminiferal Research 42, 2, p. 156-168.
(SEM images of specimens of 'Globigerina siakensis' from near its type locality in C Sumatra show spinose test and straight intercameral sutures on spiral side of test. Globorotalia mayeri has curved spiral-side intercameral sutures and lacks spinosity)

X.3. Jurassic- Cretaceous

Numerous additional papers with Jurassic-Cretaceous biostratigraphy content but are case studies specific to certain areas, are listed under those areas, and are not repeated here.

This chapter does include several papers on Jurassic-Cretaceous faunas-floras outside Indonesia that may help in the interpretation of related occurrences in Indonesia.

Arkell, W.J. (1954)- Indo-China, Hongkong and Indonesia (Ch. 17) and New Guinea, New Caledonia, New Zealand and Australia. In: Jurassic geology of the world, Oliver and Boyd, Edinburgh, Chapter 18, p. 431-462.
(Old review of knowledge of Jurassic stratigraphies, incl. the Indonesian region (Sumatrra, Borneo, eastern Indonesia, New Guinea). Late Jurassic deposition in eastern Indonesia (Misool, Sula islands, Rotti, Timor, etc.), and New Guinea is in direct communicaton with that of the Himalayas, as suggested by macrofossil assemblages (Tethyan N Gondwana))

Baumgartner, P.O., A. Bartolini, E.S. Carter, M. Conti, G. Cortese, T. Danelian, P. De Wever et al. (1995)- Middle Jurassic to Early Cretaceous radiolarian biochronology of Tethys based on Unitary Associations. In: P.O. Baumgartner et al. (eds.) Middle Jurassic to Lower Cretaceous Radiolaria of Tethys: occurrences, systematics, biochronology, Memoires Geologie, Lausanne, p.

Baumgartner, P.O., L. O'Dogherty, S. Gorican, E. Urquhart, A. Pillevuit & P. de Wever (eds.) (1995)- Middle Jurassic to Lower Cretaceous radiolaria of Tethys: occurrences, systematics, biochronology. InterRad Jurassic-Cretaceous Working Group, Memoires Geologie (Universite de Lausanne) 23, p. 1-1172.
(Key publication on M Jurassic- E Cretaceous radiolaria from Tethys region, mainly western Tethys and Japan)

Baumgartner, P.O., L. O'Dogherty, S. Gorican, R. Dumitrica-Jud, P. Dumitrica, A. Pillevuit, E. Urquhart, A. Matsuoka et al. (1995)- Radiolarian catalogue and systematics of Middle Jurassic to early Cretaceous Tethyan genera and species. In: P.O. Baumgartner et al. (eds.) Middle Jurassic to Lower Cretaceous Radiolaria of Tethys: occurrence, systematics, biochronology. Memoires Geologie (Universite de Lausanne) 23, p. 37-685.

Beauvais L. (1989)- Jurassic corals of the Circum-Pacific area. Memoir Assoc. Australasian Palaeontologists (AAP) 8, p. 291−302.

Beauvais, L., M.C. Bernet-Rollande & A. Maurin (1985)- Reinterpretation of Pretertiary classical reefs from Indo-Pacific Jurassic examples. In: C. Gabrie & M. Harmelin (eds.) Proc. Fifth International Coral Reef Congress, Tahiti 1985, 6, Misc. Paper (B), p. 581-586.
(Jurassic carbonate mounds in W Thailand (M-U Jurassic, Mae Sot basin), C Sumatra (U Jurassic, Padang-Tembesi River) and Philippines (M Jurassic, Mindoro, U Jurassic Calamian Isl.) not 'reefs' like present day reefs. Corals typically float in lime mud matrix and are mainly digitate or lamellar, to cope with muddy conditions. Calcareous sponges also common. Main rock-building organisms are Bacinellid- Lithocodium-stromatolite assemblage, as encrusters over exotic grains or as single builder. Jurassic corals, sponges etc., have no major rock building potential)

Beauvais, L., H. Fontaine & A. Maurin (1987)- A review of recent data on mud-mounds discoveries in Asia. Oil and Gas Geology 1987, 12, p. 373-376.
(Many of Mesozoic carbonates in SE Asia are probably microbial mud mounds, not reefs: Jurassic of Sumatra, Thailand, Burma, Philippines, Sarawak-Kalimantan)

Belford D.J. & V. Scheibnerova (1971)- Turonian foraminifera from the Carnarvon Basin, Western Australia, and their palaeogeographical significance. Micropaleontology 17, p. 331-344.
(Presence of E-M Turonian planktonic foraminifera Praeglobotruncan stephani, P. hagni, P. imbricata, P. helvetica and Hedbergella spp from wells in Carnarvon Basin show Tethyan character)

Bolli, H.M. (1974)- Jurassic and Cretaceous calcisphaerulidae from DSDP Leg 27, Eastern Indian Ocean. In: J.J. Veevers et al. (eds.) Initial Reports Deep Sea Drilling Project (DSDP) 27, p. 843-907.

(online at: www.deepseadrilling.org/27/volume/dsdp27_39.pdf)
(19 new species of Pithonella, family Calcisphaerulidae, from U Jurassic and Cretaceous sediments from Sites 259, 260, 261, and 263 of Leg 27 in E Indian Ocean. Previously, calcisphaerulidae (av. size 40-120μ) described only from thin sections, like Stomiosphaeridae and Cadosinidae from Late Jurassic- earliest Cretaceous pelagic limestones of Seram, Timor, Roti, Buton and Misool by Wanner 1940 and Vogler 1941)

BouDagher-Fadel, M.K. (2008)- The Mesozoic larger benthic foraminifera: the Jurassic. In: Evolution and geological significance of larger benthic foraminifera, Chapter 4, Developments in Palaeontology and Stratigraphy, Elsevier, 21, p. 157-213.
(General review of Jurassic larger foraminifera)

Boudagher-Fadel, M.K. & G.D. Price (2019)- Global evolution and paleogeographic distribution of mid-Cretaceous orbitolinids. UCL Press, University College London, p. 1-21.
(online at: https://www.scienceopen.com/document?vid=c36a0eb8-aaf1-4f37-87e7-009f3718bea3)
(Review of Aptian- Cenomanian larger foraminifera Orbitolina. Includes new material from W flank of Meratus Mountains, S Kalimantan, identified here as Late Aptian- E Albian Mesorbitolina texana, Palorbitolinoides orbiculata, Mesorbitolina subconcava and Conicorbitolina sp.)

Brunnschweiler, R.O. (1963)- A review of the sequence of *Buchia* species in the Jurassic of Australasia. Proc. Royal Society of Victoria 76, p. 163-168.
(On ranges of Buchia (= Aucella) bivalve mollusc species in Oxfordian-Tithonian. Now assigned to genus Malayomaorica)

Challinor, A.B. (1989)- The succession of *Belemnopsis* in the Late Jurassic of Eastern Indonesia. Palaeontology 32, 3, p. 571-596.
(online at: https://www.palass.org/publications/palaeontology-journal/archive/32/3/article_pp571-596)
(Belemnopsis from Misool and Sula all part of B. moluccana lineage. Misool Late Jurassic stratigraphy condensed rel. to Sula. Misool: 85m of Oxfordian Demu Fm carbonate/ shale overlain by ~100m of Kimmeridgean-Tithonian Lelinta shale with minor sandstone)

Challinor, A.B. (1991)- Belemnite successions and faunal provinces in the Southwest Pacific, and the belemnites of Gondwana. BMR J. Australian Geology Geophysics 12, 4, p. 301-325.
(online at: www.ga.gov.au/corporate_data/81298/Jou1991_v12_n4_p301.pdf)
(M Jurassic- E Cretaceous belemnites from SW Pacific, New Guinea, Misool, Sula, tied to dinoflagellate zonations. Two belemnite provinces in SW Pacific region from M Jurassic- E Cretaceous: (1) Tethyan: E Indonesia, W Papua, PNG, parts of N and W Australia; (2) S Pacific: New Zealand, most of Australia)

Challinor, A.B. (1992)- Belemnites of the Southwest Pacific. In: G.E.G. Westermann (ed.) The Jurassic of the Circum-Pacific, World and Regional Geology 3, Cambridge University Press, p. 308-309.
(Brief summary of Jurassic belemnites in E Indonesia, PNG, New Zealand. Three main assemblages Dicoelites-Conodicoelites (Late Bajocian- E Oxfordian), Hibolithes (late Callovian- Oxfordian) and Belemnopsis (basal Oxfordian- latest Tithonian))

Cloos, H. (1916)- Doggerammoniten aus den Molukken. I. Text. Habilitationsschrift, Hohen Phil. Fakultat Konigliche Universitat Marburg, Schweizerbart, Stuttgart, p. 1-50.
('Dogger (= M Jurassic) ammonites from the Moluccas'. 'Habilitation' thesis on Hammoceratid ammonites from Misool and Mangoli, and Stephanoceratids from Mangoli and Taliabu. Text volume only; part 2 never published due to WWI)

Colom, G. (1965)- Essais sur la biologie, la distribution geographique et stratigraphique des Tintinnoidens fossiles. Eclogae Geologicae Helvetiae 58, 1, p. 319-334.
(online at: http://dx.doi.org/10.5169/seals-163267)

('Review of the biology, geographic and stratigraphic distribution of the fossil Tintinnids'. Classic paper on Jurassic-Cretaceous tintinnid nannofossils, incl. records of Late Jurassic Calpionella alpina group in PNG (Maril Shale; Rickwood 1955) N Australia (Brunnschweiler, 1960))

Cookson, I.C. & A. Eisenack (1958)- Microplankton from Australia and New Guinea Upper Mesozoic sediments. Proc. Royal Society of Victoria 70, p. 19-79.
(online at: http://takata.slv.vic.gov.au/...)
(Early paper of Late Jurassic- Early Cretaceous dinoflagellates. Mainly taxonomic descriptions of 75 species from Australian NW shelf, some from PNG (Omati River, Era River))

Cookson, I.C. & A. Eisenack (1960)- Upper Mesozoic microplankton from Australia and New Guinea. Palaeontology 2, 2, p. 243-261.
(online at: http://cdn.palass.org/publications/palaeontology/volume_2/pdf/vol2_part2_pp243-261.pdf)
(Upper Jurassic- basal Cretaceous dinoflagellates and hystrichospheres from Canning basin, W Australia and other localities in Australia and New Guinea)

Cookson, I.C. & A. Eisenack (1974)- Mikroplankton aus Australischen Mesozoischen und Tertiaeren Sedimenten. Palaeontographica, B, Palaeophytologie, 148, p. 44-93.
('Microplankton from Australian Mesozoic and Tertiary sediments')

Crame, J.A. (1983)- The occurrence of the Upper Jurassic bivalve *Malayomaorica malayomaorica* (Krumbeck) on the Orville Coast, Antarctica. J. Molluscan Studies 49, p. 61-76.
(online at: http://mollus.oxfordjournals.org/content/49/1/61.full.pdf
(First record of Malayomaorica malayomaorica in Antarctica. Late Jurassic bivalve species (originally assigned to Aucella, then Buchia) mainly limited to Kimmeridgean. Appears to be typical of Late Jurassic margins of Gondwanaland, including NW Australia, New Zealand, New Caledonia and E Indonesia (Misool, Timor, New Guinea, Sula, E Sulawesi, Ceram, Buru; JTvG)

Dhondt, A.V. (1992)- Cretaceous inoceramid biogeography: a review. Palaeogeogr. Palaeoclim. Palaeoecology 92, p. 217-232.
(Cretaceous inoceramid bivalves did not thrive in shallow or warm seas, and therefore rare in Tethyan shallow deposits. Occur mainly in temperate seas, and distribution often bipolar. Not much on SE Asia)

Fontaine, H. & L. Beauvais (1986)- Distribution of the Jurassic corals in Southeast Asia. In: P.Q. Tuong (ed.) Proc. First Conf. Geology of Indochina, Ho Chi Minh City 1986, Gen. Dept. of Geology Vietnam, 1, p. 137-145.
(On Jurassic corals from S Vietnam, Cambodia, S Laos, Philippines (Mindoro, Calamian Islands, NE Palawan), Borneo (W Sarawak, W Kalimantan), Sumatra, Thailand (Mae Sot))

Hallam, A. (1977)- Jurassic bivalve biogeography. Paleobiology 3, p. 58-73.
(No maps. Incl. Late Jurassic SW Pacific Province: with Malayaomaorica as only endemic bivalve genus. Same or closely re lated species in Spiti Shales of Himalayas. Buchia and Retroceramus abundant in the Arctic/ Antarctic, but also common in lower latitudes)

Hayami, I. (1984)- Jurassic marine bivalve faunas and biogeography in Southeast Asia. In: T. Kobayashi et al. (eds.) Geology and Palaeontology of Southeast Asia 25, University of Tokyo Press, p. 229-237.
(Unique E Jurassic (Pliensbachian?) heavy bivalve assemblage from Timor with Lithiotis, Pachymegalodon, Gervilleioperna, etc. described from Fatu Lst of Timor by Krumbeck (1923). Upper Jurassic bivalves in W Borneo part of East Asian Province with Philippines and Japan. Timor-Roti, Seram, Misool, etc., are part of Maorian Province with Malayomaorica and Retroceramus haasti)

Hofker, J., Jr. (1963)- Studies on the genus *Orbitolina* (Foraminiferida). Leidsche Geologische Mededelingen 29, p. 181-253.
(online at: https://repository.naturalis.nl/pub/505874)

(Study of mid-Cretaceous (Barremian-Cenomanian) 'Tethyan' larger foram genus Orbitolina. Includes material from W Kalimantan Seberuang area (collected by Van Schelle and Wing Easton), and classified as relatively primitive 'Group I- Lower Aptian' species (N.B. Schlagintweit and Simmons (2023) assigned 'Orbitolina' from Borneo to subgenus Palorbitolina lenticularis, signifying late Barremian- E Aptian age; JTvG)

Iba, Y. & S. Sano (2006)- *Mesorbitolina* (Cretaceous larger foraminifera) from the Yezo Group in Hokkaido, Japan and its stratigraphic and paleobiogeographic significance. Proc. of the Japan Academy, B, 82, 7, p. 216-223.
(online at: https://www.jstage.jst.go.jp/article/pjab/82/7/82_7_216/_pdf)
('Tethyan' low-latitude Aptian Orbitolina (Mesorbitolina) parva from limestone olistoliths from olistostrome horizon in lower Yezo Group, Hokkaido, represent first report of this species from Circum-North Pacific)

Iba, Y., S. Sano & T. Miura (2011)- Orbitolinid foraminifers in the Northwest Pacific: their taxonomy and stratigraphy. Micropaleontology 57, 2, p. 163-171.

(Four 'Tethyan' orbitolinid larger foram species ('Palorbitolina lenticularis', Praeorbitolina cf. wienandsi, Mesorbitolina parva, M. texana) recognized in Late Hauterivian- Early Albian of Japan and South Sakhalin)

Jaworski, E. (1931)- Ueber *Arnioceras geometricum* Oppel 1856 und verwandte Spezies; nebst einem Anhang uber *Arnioceras natrix* v. Schlotheim 1820. Neues Jahrbuch Mineralogie Geologie Palaontologie, Beilage Band 65, B, p. 83-140.
('Arnioceras geometricum Oppel (1856) and related species; with Appendix on Arnioceras natrix v. Schloth. 1820'. Preliminary results of study on Early Jurassic ammonites from Netherlands Indies (Babar island, etc.))

Jaworski, E. (1933)- Revision der Arieten, Echioceraten und Dactylioceraten des Lias von Niederlandisch-Indien. Neues Jahrbuch Mineralogie Geologie Palaontologie, Abhandlungen, Beilage Band 70, B, p. 251-333.
(online at: http://mmtk.ginras.ru/pdf/Jaworski_1933_J_Arieten_Indonesia_HI.pdf)
('Revision of the Arietidae, Echioceratidae and Dactylioceratidae from the Liassic of Netherlands Indies'. Mainly taxonomic descriptions of Early Jurassic ammonites from Roti, Babar, Timor and Sula Islands from collections in Amsterdam, Leiden, Utrecht, Delft, Bonn, Berlin and Basel: Arnioceras spp., Arieticeras, Dactylioceras spp., Coeloceras moermanni, Euechioceras wichmanni, etc. Little on stratigraphy, no locality maps)

Jeletzky, J.A. (1963)- *Malayomaorica* gen. nov. (Family Aviculopectinidae) from the Indo-Pacific Upper Jurassic, with comments on related forms. Palaeontology 6, p. 148-160.
(online at: http://palaeontology.palass-pubs.org/pdf/Vol%206/Pages%20148-160.pdf)
(S Hemisphere Late Jurassic bivalves described as Buchia and Aucella differ from N Hemisphere-Boreal Buchia, therefore assigned to new genus Malayomaorica. Typical of Kimmeridgean of Gondwana margin, including NW Australia, New Zealand New Guinea, Misool, Sula, E Sulawesi, Timor, Ceram, Buru; JTvG)

Kobayashi, T. (1978)- The Jurassic palaeogeography of Japan and Southeast Asia. Proc. of the Japan Academy, Series B, 54, 10, p. 583-588.
(online at: https://www.jstage.jst.go.jp/article/pjab1977/54/10/54_10_583/_pdf/-char/en)
(Review of Jurassic facies in Japan and across SE Asia, Shallow marine E Jurassic facies inSumatra and NW Kalimantan. Also Jurassic--Cretaceous marine sequence in Sarawak. With one small E Jurassic facies map (no mention of widespread marine Jurassic in East Indonesia- New Guinea; JTvG))

Kobayashi, T. (1979)- The *Trigonioides* basins and the Cretaceous palaeogeography of East and Southeast Asia. Proc. of the Japan Academy, Series B, 55, B 1, p. 1-5.
(online at: https://www.jstage.jst.go.jp/article/pjab1977/55/1/55_1_1/_pdf)
(On distribution of Early-Middle Cretaceous non-marine bivalve mollusc Trigonioides in SE Asia, including in continental facies of Rantaulajung Fm near Martapura, SE Kalimantan with Upper Cretaceous conchostracans)

Kruizinga, P. (1931)- Cephalopoda. In: B.G. Escher et al. (eds.) De palaeontologie en stratigraphie van Nederlandsch Oost-Indie, Leidsche Geologische Mededelingen 5 (K. Martin memorial volume), p. 297-389.
(online at: www.repository.naturalis.nl/document/549628)
(Summary of state of knowledge of ammonites and belemnites in Indonesia in 1931. Highest number of species known is from Timor. Other significant cephalopd faunas on Roti, Sula Islands, Buru and Misool. Most known species of Triassic age, other rich faunas of Jurassic and Permian, but relatively few species from Cretaceous)

Leinfelder, R.R., D.U. Schmid, M. Nose & W. Werner (2002)- Jurassic reef patterns- the expression of a changing globe. In: W. Kiessling et al. (eds.) Phanerozoic Reef Patterns, Society for Sedimentary Geology (SEPM) Special Publ. 72, p. 465-520.
(online at: http://userpage.fu-berlin.de/leinfelder/palaeo_de/leinfelder/rrlprints/Leinfelder_etal_2002_lr.pdf)
(Includes brief discussions of Jurassic carbonates of W Thailand, Sumatra and Philippines. Early- Middle Jurassic reefs absent in SE Asia, except small Lithiotis bivalve mounds on Timor, due to end-Triassic extinction event, etc. Minor Late Jurassic reefs in Sumatra and Bau Limestone of Sarawak- NW Kalimantan border area)

Leupold, W. & W. Maync (1935)- Das Auftreten von *Choffatella*, *Pseudocyclammina*, *Lovcenipora* (*Cladocoropsis*) und *Clypeina* im alpinen Faziesgebiet. Eclogae Geologicae Helvetiae 28, p. 129-139.
(online at: http://retro.seals.ch/digbib/view?pid=egh-001:1935:28::191)
('The occurrences of Choffatella, Pseudocyclammina, Lovcenipora (Cladocoropsis) and Clypeina in the Alpine facies region'. First record of Late Jurassic Pseudocyclammina associated with Cladocoropsis mirabilis Felix from Switzerland. Assemblage formerly known from Japan and Barisan Mts of Sumatra (see also Maync 1952))

Livsey, A.R., D.J. Mantle, C.D. Bates & A.J. Charles (2024)- The use of quantitative palynology in Pre-Tertiary hydrocarbon exploration and development in Eastern Indonesia. Proc. 48th Annual Conv. Indonesian Petroleum Association (IPA), Jakarta, IPA24-SS-03, p. 2006-2015.
(Review of Jurassic- E Cretaceous palynology method and zonation, as used in NW Australian shelf and in Bintuni Basin of W Papua. With biozonation charts. (thin on stratigraphic results in E Indonesia; JTvG))

Mamgain, V.D. & B.R.J. Rao (1965)- Orbitolines from the limestone intercalations of Dras Volcanics, Jammu and Kashmir State. J. Geological Society India 6, p. 122-129.
(Rich and well-preserved E Cretaceous Orbitolina (incl. Orbiqia drasensis n.sp.) in limestone intercalations in volcanic rocks of Dras, in Jammu and Kashmir state (Ladakh, N India). Also Orbitolina trochus. Barremian-Aptian age (Orbiqia now accepted as Mesorbitolina; = part of the intra-Neo?Tethyan Dras-Kohistan-Ladakh volcanic arc? = related to Woyla terrane of Sumatra?; JTvG))

Matsumaru, K., M. Aizawa, K. Mukai & A. Furusawa (2007)- Note of orbitolinid foraminifera from the Lower Aptian (Cretaceous) Shimanoshita Mudstone, Lower Yezo Group, Hokkaido, Japan. J. Saitama University Fac. Educ. 56, l, p. 367-372.
(Mid-Cretaceous 'Tethyan' larger foram genus Orbitolina at several localities across accretionary prism of SE Japan and also Taiwan Shimanoshita location E Aptian with Palorbitolina lenticularis, Mesorbitolina parva, Praeorbitolina)

Matsumaru, K., M. Ehiro & S. Kojima (2006)- On *Orbitolina* (Foraminiferida) from the Shyok suture zone, Ladakh, NW India. J. Palaeontological Society India 51, 2, p. 43-49.
(online at: http://palaeontologicalsociety.in/vol51_2/v4.pdf)
(E-M Albian Orbitolina from Lower Shyok Fm in Shyok suture zone, Ladakh, NW India, in sediments overlying ultramafic unit. Incl. Mesorbitolina texana, M. minuta, Simplorbitolina cf. conulus)

Matsumaru, K. & A. Furusawa (2007)- On orbitolinid foraminifera from the lower Aptian (Cretaceous) of Hokkaido, Japan. J. Palaeontological Society India 52, 1, p. 39-44.
(online at: http://palaeontologicalsociety.in/vol52_1/v4.pdf)
(Five E Aptian orbitolinid species in Takisato Orbitolina Lst of Hokkaido: Palorbitolina lenticularis, Mesorbitolina parva, M. minuta, M. libanica, Paleodictyoconus conica (compares to Paleo-Pacific active margin of Borneo?; JTvG))

Matsuoka, A. (1995)- Jurassic and Lower Cretaceous radiolarian zonation in Japan and in the western Pacific. The Island Arc 4, p. 140-153.
(Radiolarian zonation for Jurassic- Lower Cretaceous from Japan outcrop sections and W Pacific seafloor; applicable to low and middle paleolatitude portions of Paleo-Pacific ocean. 11 zones proposed)

Maync, W. (1952)- Critical taxonomic study and nomenclatural revision of the Lituolidae based upon the prototype of the Family, *Lituola nautiloidea* Lamarck, 1804. Contr. Cushman Foundation Foraminiferal Research 3, 2, p. 35-56.
(online at: https://cushmanfoundation.allenpress.com/portals/_default/files/pubarchive/CCFFR/03ccffr2.pdf)
(Includes discussion of Lacazina lamellifera Silvestri 1925 and Loftusia bemmeleni Silvestri 1932 from Saling Lst of S Sumatra (not species of Pseudocyclammina). Confirms presence of true Late Jurassic- E Cretaceous Loftusia and Pseudocyclammina in samples from Gumai Mts of S Sumatra (= Woyla Terrane; JTvG))

Munasri (2000)- Microfossils radiolaria in Indonesia: introducing the technique of preparation. Proc. 29th Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 4, p. 259-264.

Munasri (2001)- Radiolarian research in some Mesozoic provinces in Indonesia. Berita Sedimentologi 16, p. 26-30.
(online at: https://journal.iagi.or.id/index.php/FOSI/article/view/242)
(Brief review of radiolaria studies from Cretaceous of C Java (Karangsambung), M Cretaceous from S Sulawesi (Bantimala), Jurassic-Cretaceous from SE Kalimantan (Meratus Range), M-L Triassic and E Cretaceous from W Timor (Nefokoko, Kefamenanu, Kolbano) and M Jurassic from Roti island)

Munasri (2012)- Penggunaan fosil radiolaria dalam sintesis geologi. Pros. Pemaparan Hasil Penelitian Pusat Penelitian Geoteknologi LIPI, Bandung 2012, p. 337-350.
('The use of fossil radiolaria in geological synthesis'. Review of Mesozoic radiolarian research in Indonesia. Radiolarian fossils found in deep-sea environments and are used for biostratigraphy, tectonic reconstructions, paleo-oceanography and paleogeography. With summary of 15 areas in Indonesia with radiolarian sediments)

Munasri, K. Wakita & K. Sashida (1999)- Fosil radiolaria sebagai alat biostratigrafi yang baru di Indonesia. Proc. 27th Annual Conv. Indonesian Association Geologists (IAGI), 2 (Sedimentology Paleontology Stratigraphy), Yogyakarta, p. 48-52.
(https://www.iagi.or.id/web/digital/56/31.pdf)
('Fossil radiolaria as a new biostratigraphic tool in Indonesia'. Review of radiolaria studies in Cretaceous subduction complexes in Java (Karangsambung), S Sulawesi (Bantimala, Barru), SE Kalimantan (Pulau Laut), Timor (Kolbano))

O'Dogherty, L. (2009)- Inventory of Mesozoic radiolarian species (1867-2008). Geodiversitas (Museum National d'Histoire Naturelle, Paris) 31, 2, p. 371-481.
(online at: https://sciencepress.mnhn.fr/sites/default/files/articles/pdf/g2009n2a6.pdf)
(Comprehensive listing 6296 Mesozoic radiolarian species names, in combinations as originally described, with approximate ages. E.g. helpful in confirming correctness of Hojnos (1934) Late Jurassic- E Cretaceous age assignments of E Sulawesi radiolarian rocks)

O'Dogherty L., E.S. Carter, P. Dumitrica, S. Gorican, P. De Wever, A.N. Bandini, P.O. Baumgartner & A. Matsuoka (2009)- Catalogue of Mesozoic radiolarian genera. Part 2: Jurassic-Cretaceous. Geodiversitas 31, 2, p. 271-356.
(online at: https://www.researchgate.net/publication/232690604_Catalogue_of_Mesozoic_Radiolarian_Genera_Part_2_Jurassic-Cretaceous
(Key work. Illustrated catalogue of type species of 581 genera of Jurassic- Cretaceous radiolaria. No range charts)

Oloriz, F. & G.E.G. Westermann (1998)- The perisphinctid ammonite *Sulaites* n. gen. from the Upper Jurassic of the Indo-Southwest Pacific. Alcheringa 22, 3-4, p. 231-240.
(New genus Sulaites comprises Oxfordian group of 'Perisphinctes' sularus and moluccanus, described from Sula Islands, and Late Oxfordian-?E Kimmeridgian 'Pseudoparaboliceras aramaraii' group described from W Papua. Genus Sulaites also known from W Papua, PNG and probably New Zealand and Nepal)

Oosting, A.M. (2004)- Palaeoenvironmental and climatic changes in Australia during the Early Cretaceous. Thesis University Utrecht, p. 1-203.
(online at: http://dspace.library.uu.nl/handle/1874/1578)
(Biochronostratigraphy for Tethyan and Boreal M Cretaceous traditionally based on ammonites, but because of lack of useful ammonites in Australian M Cretaceous, and faunal and floral latitudinal contrasts, correlations between different realms is not straightforward. Dinoflagellate cyst events combined with changes in carbon-isotope stratigraphy used here to assess Barremian- Albian stage boundaries in Australia)

Pannekoek, A.J. (1931)- Brachiopoda. In: B.G. Escher et al. (eds.) De palaeontologie en stratigraphie van Nederlandsch Oost-Indie, Leidsche Geologische Mededelingen 5, 1 (K. Martin memorial volume), chapter 12, p. 396-435.
(online at: https://repository.naturalis.nl/pub/505703)
(Summary of state of knowledge in 1931 of Late Paleozoic- Mesozoic- Tertiary brachiopod species and distribution in Indonesia)

Philippe, M., A. Boura, C. Oh & D. Pons (2014)- *Shimakuroxylon* a new homoxylous Mesozoic wood genus from Asia, with palaeogeographical and palaeoecological implications. Review Palaeobotany Palynology 204, p. 18-26.
(New type of (Late?) Jurassic-E Cretaceous fossil wood with radial pitting of 'japonicum type', named here Shimakuroxylon. Geographic distribution limited to terranes which lined S-most E Asia during Jurassic (Lhasa, Indochina, Semitau, etc.). W Kalimantan specimen in British Museum collected from Buduk (Boedak) 100km N of Pontianak, associated with bivalves identified as M Jurassic by Newton (1903) (probably Late Jurassic?). Also present in Outer Zone of SW Japan. Probably indicator for warm and wet climates)

Philippe, M., H.E. Jiang, K. Kim, C. Oh, D. Gromyko, M. Harland, I.S. Paik & F. Thevenard (2009)- Structure and diversity of the Mesozoic wood genus *Xenoxylon* in Far East Asia: implications for terrestrial palaeoclimates. Lethaia 42, p. 393-406.
(Mesozoic fossil wood Xenoxylon is indicator of wet temperate biotopes and is common in Far East Asia in Carnian-Maastrichtian. It is part of 'Northern-type' leaf-flora (also call Tetori-type or Siberian-Canadian), although still present in Vietnam. Diversity peak of Xenoxylon spp. centered on NE China, where wet-temperate climate probably prevailed through Late Triassic- Cretacous (genus not known from Indonesia or Gondwana))

Remane, J. (1985)- Calpionellids. In: H.M. Bolli, J.B. Saunders & K. Perch-Nielsen (eds.) Plankton Stratigraphy, Cambridge University Press, p. 555-572.
(Review of Late Tithonian- E Valanginian planktonic protozoans of unknown affinities. Includes reported, but not illustrated, presence of calpionellids in PNG by Rickwood (1955))

Renz, G.W. (1974)- Radiolaria from Leg 27 of the Deep Sea Drilling Project. In: J.J. Veevers et al. (eds.) Initial Reports Deep Sea Drilling Project (DSDP) 27, p. 769-841.
(online at: www.deepseadrilling.org/27/volume/dsdp27_38.pdf)
(Study of Cretaceous radiolarians from Leg 17 sites, mainly Site 261 in eastern Indian Ocean and Site 262 in Timor Trough of SW Timor. Succession of three assemblages, from young to old: (1) Bathropyramis timorensis, (2) Eucyrtis columbarius and (3) Spongocyclia lanigera Assemblage, of uncertain ages (but placed in ~Berriasian-Aptian range by Sanfilippo and Riedel 1985, Fig. 2; JTvG). Includes brief discussion of Tan Sin Hok (1927) Roti sample 149 assemblage, which is believed to be younger than this DSDP material)

Riding, J.B. (2012)- A compilation and review of the literature on Triassic, Jurassic, and earliest Cretaceous dinoflagellate cysts. American Assoc. Stratigraphic Palynologists (AASP), Contributions Series 46, Dallas, p. 1-126. *(also 2020 edition, 325 p.)*
(online at: https://nora.nerc.ac.uk/id/eprint/19423/1/Jurassic_dinocyst_reference_list_-_revised_manuscript_March_2012_word_2003.pdf)
(Parts of latest documentation of the highly successful Mesozoic dinolagellate biostratigraphy, applicable of Australia NW Shelf and New Guinea margins)

Riding, J.B. (2013)- The literature on Triassic, Jurassic and earliest Cretaceous dinoflagellate cysts: supplement 1. Palynology 37, 2, p. 345-354.

Riding, J.B., E. Mariani & R.A. Fensome (2022)- A review of the Jurassic dinoflagellate cyst genus *Gonyaulacysta* Deflandre 1964 emend. nov. Review Palaeobotany Palynology 299, 104605, p. 1-37.
(Review of M-L Jurassic gonyaulacacean dinoflagellate cyst genus Gonyaulacysta Deflandre, including material from Australia NW Shelf)

Sahni, M.R. & V.V. Sastri (1957)- A monograph of the orbitolines found in the Indian continent (Chitral, Gilgit, Kashmir), Tibet and Burma, with observations on the age of the associated volcanic series. Memoirs Geological Survey India, Palaeontologia Indica 33, 3, p. 1-50.
(online at: http://histdept.crma.ac.th:8011/ebooks/20100511-the-geological-survey-of-india.pdf)
(Mid-Cretaceous larger foraminifera Orbitolina from localities in collision zones around N and NE margin of Indian subcontinent. Including multiple Myanmar localities (Yanbo, Ku Taung, Mesan, etc.), initially sampled by Clegg. Two Orbitolina limestone horizons in Burma: (1) Barremian- Aptian with Orbitolina bulgarica, O wadiai n.sp., O. raoi n.sp. and O. hukawngensis n.sp.; (2) Cenomanian with Orbitolina birmanica Sahni 1937 (species with great affinity with Orbitolina concava)

Sano, S.I. & P.W. Skelton (2010)- *Epidiceras* (Bivalvia, Hippuritoidea) from the Tithonian-Berriasian Torinosu-type Limestones of the Sakawa Area, Southwest Japan. Turkish J. Earth Sciences 19, p. 733-743.
(online at: http://journals.tubitak.gov.tr/earth/issues/yer-10-19-6/yer-19-6-5-0905-2.pdf)
(Primitive rudists Epidiceras speciosum and E. guirandi from Tithonian-Berriasian Torinosu limestones in SW Japan. Epidiceras speciosum also present in Kimmeridgean-Tithonian Bau Limestone of SW Sarawak. Tethyan rudists proven here to extend into W Pacific province)

Sarjeant, W.A.S., W. Volkheimer & W.P. Zhang (1992)- Jurassic palynomorphs of the Circum-Pacific region. In: G.E.G. Westermann (ed.) The Jurassic of the Circum-Pacific, World and Regional Geology 3, p. 273-292.
(Includes discussion of Jurassic palynomorphs/ dinoflagellates of Papua New Guinea, Australia. The only Jurassic palynoflora from SE Asia is partly published work from Misool by Helby and Hasibuan: Yefbie Shale has characteristic Toarcian Susadinium assemblage and Bajocian- E Bathonian Dissioliodinium association;base Demu Fm has Late Callovian- E Oxfordian Rigaudella aemula zone, Late Oxfordian- E Kimmeridgean Wanaea spectabilis zone in rest of Demu Fm and basal Lelinta shale; later assemblages suggest age as young as Kalyptea wisemaniae zone, E Berriasian)

Sato, T. (1956)- Correlation du Jurassique inferieur japonais en basant sur les ammonites fossiles. J. Geological Society Japan 62, 732, p. 490-503. *(Japanese with French abstract)*
(online at: www.jstage.jst.go.jp/article/geosoc1893/62/732/62_732_490/_pdf)
(Incl. circum-Pacific 'Aalenian' (Late Toarcian?) ammonite distribution map, showing distribution of Hammatoceras in E Indonesia)

Sato, T. (1975)- Marine Jurassic formations and faunas in Southeast Asia and New Guinea. In: T. Kobayashi & R. Toriyama (eds.) Geology and palaeontology of Southeast Asia, University of Tokyo Press, 15, p. 151-189.
(Review of Jurassic macrofossils and stratigraphy in SE Asia, including Sumatra, Timor,New Guinea, E Sulawesi, Sula, Buru, Ceram and Misool)

Sato, T. & T. Ishibashi (1984)- Ammonoids of Southeast Asia. In: T. Kobayashi, R. Toriyama & W. Hashimoto (eds.) Geology and palaeontology of Southeast Asia, University of Tokyo Press, 25, p. 267-272.
(Brief review of Triassc- Cretaceous ammonoids in Indonesia and surrounding SE Asia. No figures)

Sato, T. & G.E.G. Westermann (1991)- 4. Japan and Southeast Asia. In: G.E.G. Westermann & A.C. Ricardi (eds.) Jurassic taxa ranges and correlation charts fror the Circum-Pacific. Newsletters on Stratigraphy 24, 1-2, p. 81-108.
(Useful compilation of distribution and ranges of Jurassic macrofossils in Indonesia)

Scrivenor, J.B. (1912)- Radiolaria-bearing rocks in the East Indies. Geological Magazine (V), 9, 6, p. 241-248.
(Review of mainly Mesozoic radiolarian-bearing rocks on the Malay Peninsula and Indonesia. Radiolarites of Danau Fm of C Borneo (probable Jurassic age) do look like oceanic deposits, but other radiolaria-bearing rocks often associated with clastic material and may be of shallower marine origin)

Sha, J. (2007)- Cretaceous trigonioidid (non-marine Bivalvia) assemblages and biostratigraphy in Asia with special remarks on the classification of Trigonioidacea. J. Asian Earth Sciences 29, 1, p. 62-83.
(Seven zones distinguished in distribution of Cretaceous Trigonioides-group fresh water molluscs. Mainly on mainland Asia (China, Korea, some Thailand) and Japan; nothing on Indonesia although Cretaceous non-marine bivalves are present in SE Kalimantan)

Sha, J. (2010)- Historical distribution patterns of trigonioidids (non-marine Cretaceous bivalves) in Asia and their palaeogeographic significance. Proc. Royal Society (London), B (Biological Science), 277, p. 277-283.
(online at: http://rspb.royalsocietypublishing.org/content/early/2009/07/15/rspb.2009.0936.full.pdf)
(Non-marine trigonioidid bivalves five phases of radiation in Cretaceous of Pal-Asia: pre-Aptian (?Valanginian/ Hauterivian- Barremian), Aptian, Albian, Cenomanian and Turonian-Maastrichtian. Distribution patterns show two distinct paleo-river systems feeding trigonioidids. Trigonioidid distribution pattern suggests Japan was attached to part of E China and/or Korea in Valanginian-Cenomanian (no discussion of records from Indonesia))

Sha, J., A. Meesook & X.K. Nguyen (2012)- Non-marine Cretaceous bivalve biostratigraphy of Thailand, Southern Lao PDR and Central Vietnam. J. of Stratigraphy 36, 2, p. 382-399.
(online at: https://dcxz.cbpt.cnki.net/portal/journal/portal/client/paper/448cc2031c16a43c1e8e5005490a83d6)
(Bivalve molluscs from non-marine Cretaceous basin of Thailand, Laos and Vietnam (Khorat Group). Two trigoniodid bivalve assemblages: Aptian Trigonioides kobayashii- Plicatounio assemblages and mainly Albian Trigonioides diversicostatus- Pseudohyria assemblage)

Skelton, P.W. (1985)- Preadaptation and evolutionary innovation in rudist bivalves. Special Papers in Palaeontology 33, p. 159-173.
(Includes mention of early rudists Epidiceras speciosum (Goldfuss) and Valletia sp. from Bau Limestone in SW Sarawak, in collections of British Museum of Natural History, London)

Skelton, P.W., S.I. Sano & J.P. Masse (2013)- Rudist bivalves and the Pacific in the Late Jurassic and Early Cretaceous. J. Geological Society, London, 170, p. 513-526.
(Rudist distribution in Pacific region. (very little on SE Asia region rudists (Borneo, Misool))

Skwarko, S.K. & F. Hasibuan (1989)- A brief review of literature on the larger marine invertebrates in the Cretaceous of Indonesia with list of fossils hitherto identified. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 6, p. 44-52.
(Listing of 32 papers with descriptions of Cretaceous marine macrofossils from Indonesia since 1883. 22 faunas documented, 8 from Kalimantan, 3 each from Sumatra, Sula Islands, Misool and W Papua, one from Sulawesi. Bivalve molluscs (80 species) and cephalopods (70 species) are most diverse groups. Most determinations in need of revision)

Stevens, G.R. (1963)- Designation of a lectotype for the nominal species *Belemnopsis alfurica* (Boehm) 1907 (Cephalopoda, Dibranchiata). Trans. Royal Society of New Zealand, Geology, 2/6, p. 101-104.
(Designation of a lectotype for U Jurassic Belemnopsis alfurica (Boehm) from Upper Wai Lagoi River, Taliabu, Sula Islands. Also known as Belemnites alfurica from N India Himalayas, N Australia and New Zealand)

Stevens, G.R. (1964)- The belemnite genera *Dicoelites* Boehm and *Prodicoelites* Stolley. Paleontology 7, 4, 9, 606-620.
(online at: http://palaeontology.palass-pubs.org/pdf/Vol%207/Pages%20606-620.pdf)
(Belemnite genus name Dicoelites first used by Boehm (1906) for D. dicoelus Rothpletz from Callovian-Lower Oxfordian of Roti Island, then in 1912 for material from Callovian-Kimmeridgean? of Wai Miha, Taliabu, Sula islands (Dicoelites keeuwensis). Both have two grooves, but latter re-assigned to new genus Conodicoelites)

Stevens, G.R. (1965)- The Jurassic and Cretaceous belemnites of New Zealand and a review of the Jurassic and Cretaceous belemnites of the Indo-Pacific region. New Zealand Geological Survey, Paleontological. Bulletin 36, p. 1-283.
(Major review of J-K belemnites. In Oxfordian development of Indo-Pacific province. Kimmeridgean abundant Belemnopsis developed in Indo-Pacific, incl. Belemnopsis uhligi complex (Belemnopsis spp., incl. uhligi, indica, moluccana, etc.; formerly comonly called B. gerardi). Usually associated with Buchia and Inoceramus. In Tithonian uhligi complex persists, but progressively replaced by Hibolithes and Duvalia assemblage. Etc.)

Stolley, E. (1929)- Uber Ostindische Jura-Belemniten. Palaontologie von Timor, Schweizerbart, Stuttgart, 16, 29, p. 91-213.
('On East Indies Jurassic belemnites'. Belemnites from Molengraaff, Jonker and Weber collections from Timor, Roti, Misool, Sula islands, Seram, E Sulawesi and Yamdena/Tanimbar. Includes reports of Belemnopsis aucklandica from Timor (Ofu) and Roti (re-assigned to Belemnopsis uhligi-jonkeri group by Stevens 1964. B. aucklandica from Yamdena, re-described as Belemnopsis stolleyi by Stevens 1964; JTvG)

Street, C. & P.R. Bown (2000)- Palaeobiogeography of Early Cretaceous (Berriasian-Barremian) calcareous nannoplankton. Marine Micropaleontology 39, p. 265-291.
(Early Cretaceous nannoplankton biogeography Watznaueria spp. dominant in all settings. Assemblage composition relatively uniform between ~50° N and S. High-paleolatitude assemblages less rich, lower diversity and with more Crucibiscutum salebrosum, Stradnerlithus silveradius, Broinsonia matalosa, etc. Argo Abyssal Plain, NW of Australia, carries southern high-latitude nannofossil assemblages)

Thierry, J. (1976)- Paleobiogeographie de quelques Stephanocerataceae (Ammonitina) du Jurassique moyen et superieur; une confrontation avec la theorie mobiliste. Geobios 9, 3, p. 291-331.
(Paleogeography of Middle-Late Jurassic ammonites, showing distribution of 'Tethyan' Macrocephalites-Mayaites group (also known from E Indonesia and New Guinea; JTvG)

Van Gorsel, J.T. (1978)- Late Cretaceous orbitoidal foraminifera. In: R.H. Hedley and C.G. Adams (eds.) Foraminifera 3, Academic Press, London, p. 1-120.
(General review of Campanian-Maastrichtian orbitoidal larger foraminifera. In SE Asia members of 'Caribbean-Tropical Pacific' assemblage with Pseudorbitoides, Orbitoides, Asterorbis, etc. present in PNG, W Papua and E Philippines. Occurrence of Maastrichtian Omphalocyclus in Kalimantan never described elsewhere)

Van Gorsel, J.T. (2014)- An introduction to Mesozoic faunas and floras of Indonesia. Berita Sedimentologi 31, p. 27-56.
(online at: https://journal.iagi.or.id/index.php/FOSI/article/view/123/93)
(Review of Mesozoic fossils of Indonesia. Mesozoic-age faunas relatively widespread in Indonesia, from Sumatra, Java and Kalimantan in W to Sulawesi, Outer Banda Arc (Sumba, Timor, Tanimbar, Seram), Sula Islands and New Guinea in East. Triassic low-latitude Tethyan faunas present in both W and E Indonesia. Late Jurassic ammonites and bivalves from E Indonesia- New Guinea of more temperature character)

Vitacca, J.J., N. Sinclair, D.J. Mantle, N. Marshall & D. Peyrot (2022)- The taxonomy of selected marine microplankton from the Middle and Upper Jurassic (Callovian-Kimmeridgian) of the North West Shelf, Australia. Review Palaeobotany Palynology 304, 104668, p. 1-31.
(11 new species of M-U Jurassic dinoflagellates from Australia NW Shelf)

Von Hillebrandt, A., P. Smith, G.E.G. Westermann & J.H. Callomon (1992)- Ammonite zones of the Circum-Pacific region. In: G.E.G. Westermann (ed.) The Jurassic of the Circum-Pacific, Cambridge University Press, p. 247-272.
(Jurassic ammonite zonations of Circum-Pacific region. Lower Jurassic (Toarcian) ammonites known from Kalimantan, Sula Islands, Misool and Timor and Roti. M Jurassic ammonites mainly on Sula Islands and New Guinea (Bajocian-Bathonian Sulaites, Lower Callovian Macrocephalites keeuwensis assemblage. U Jurassic (Oxfordian Mayaites) also rich on Sula Islands; Kimmeridgean ammonites virtually unknown)

Von Hillebrandt, A., G.E.G. Westermann, J.H. Callomon & R.L. Detterman (1992)- Ammonites of the Circum-Pacific region. In: G.E.G. Westermann (ed.) The Jurassic of the Circum-Pacific, Cambridge University Press, New York, p. 342-359.
(Review of Jurassic ammonite stratigraphic and geographic distributions in Pacific region. Sula- New Guinea sub-province of Indo-Pacific Province in Bajocian- Bathonian, characterized by endemic Irianites, Satoceras and Praetulites. In Oxfordian more extended Indo-SE Pacific realm, with mayaitids also in New Zealand)

Wan, X., Y. Wu & G. Li (2003)- Distribution of Mid-Cretaceous Orbitolinids in Xizang (Tibet) and its paleobiogeographic implications. Acta Geologica Sinica 77, 1, p. 1-8. *(Chinese with English Abstract)*
(online at: www.jourlib.org/paper/4876549#.V9XKwvkrKUk)
(Mid-Cretaceous Orbitolinid larger foraminifera common in Barremian- Cenomanian of shallow marine Neotethys margins. In Tibet limited to Lhasa Block (= SE Eurasia margin during M Cretaceous) and parts of Qiantang, extending E to Myanmar and W to Ladakh. Not present on India Plate (also known from Sumatra, Java, Borneo; JTvG))

Wandel, G. (1936)- Beitrage zur Kenntnis der Jurassischen Molluskenfauna von Misol, Ost Celebes, Buton, Seran und Jamdena. In: J. Wanner (ed.) Beitrage zur Palaeontologie des Ostindischen Archipels 13, Neues Jahrbuch Mineralogie Geologie Palaeontologie, Beilage Band 75B, p. 447-526.
('Contributions to the knowledge of Jurassic molluscs from Misool, East Sulawesi, Buton, Seram and Yamdena'. Description of Mollusca, mainly collected by F. Weber. Misool faunas include upper Liassic Harpoceraten beds, lower Dogger Hammoceraten beds, Oxfordian Aucella malayomaorica marls (also in E Sulawesi). 'Lithacocera' from E Sulawesi is a Tithonian form (Arkell, 1954). Etc.)

Yabe, H. & S. Hanzawa (1926)- *Choffatella* Schlumberger and *Pseudocyclammina*- a new genus of arenaceous foraminifera. Science Reports Tohoku Imperial University, 2nd series, Geology, 9, p. 9-13.
(online at: http://ir.library.tohoku.ac.jp/re/bitstream/10097/30196/1/KJ00004178170.pdf)
(New genus name Pseudocyclammina for Late Jurassic foram 'Cyclammina' lituus from Torinosu Limestone of Japan, which species is very similar to Choffatella cyclamminoides n. sp. described by Silvestri (1925) from Sungi Tuni in Korinci, Jambi Province, Sumatra)

Yabe, H. & S. Toyama (1927)- *Cladocoropsis mirabilis* Felix from the Torinosu Limestone of Japan. Japanese J. Geology and Geography 5, p. 107-110.
(online at: https://babel.hathitrust.org/cgi/pt?id=nyp.33433110166588&seq=351)
(On (Late) Jurassic Cladocoropsis, a reefal sponge or branching stromatoporoid, in Pacific accreted terranes belt of SW Japan (frequently erroneously assigned to (Triassic) genus Lovcenipora). Widespread in Tethyan reefal limestone facies (Leinfelder et al. 2005) (also known from Sumatra (Gumai Mts= Woyla Terrane), Borneo, Timor, possibly same as Lovcenipora vinassai from Triassic of Buru- Seram?; JTvG))

Yin, J. & R. Enay (2004)- Tithonian ammonoid biostratigraphy in eastern Himalayan Tibet. Geobios 37, 5, p. 667-686.

(Rich ammonoid faunas in Tithonian- Lw Berriasian in E Himalayas of Tibet, with Tithonian Virgatosphinctes-Aulacosphinctoides and Uhligites-Aulacosphinctes; U Tithonian dominated by Blanfordiceras wallichi, etc. Also presence of Kimmeridgean Sulaites and Belemnopsis galoi. (NB: strong affinities between M-L Jurassic ammonite faunas of N India- Tibetan Himalayas area and E Indonesia (Sula islands, New Guinea)- SW Pacific ammonoid faunas, typical of the Jurassic North Gondwana margin- southern Meso- or Neo-Tethys; JTvG)

X.4. Triassic

1. *As mentioned for the other sub-chapters on Biostratigraphy, many papers documenting occurrences of Triassic fossils from Indonesia as well as surrounding areas are grouped with their respective areas, and are not duplicated here;*
2. *This listing below captures more general papers on Triassic fossils, including papers from outside Indonesia, that may be relevant for determining paleobiogeographic affinities of Triassic faunas and floras of Indonesia*
3. *Of particular interest are Late Triassic limestones, with Tethyan faunas are particularly widespread, from mainland SE Asia (Thailand, Laos, Vietnam, Indonesia (Sumatra, Timor, Seram), NW Australia (Wombat Plateau) to 'Panthalassan' seamount carbonates in accretionary complexes in Japan*

Ager, D.V. (1968)- The supposedly ubiquitous Tethyan brachiopod *Halorella*. J. Paleontological Society of India 5-9, p. 54-70.
(online at: http://palaeontologicalsociety.in/vol5/v11.pdf)
(Late Triassic rhynchonellid brachiopod Halorella common in Europe. Halorella nimassica described from Timor by Krumbeck (1921-1924) not true Halorella, but assigned to Timorhynchia n. gen., Halorella spp. described from Seram by Wanner (1907) look more like true Halorella, but here placed in Halorelloidea)

Al-Shaibani, S., D. Altiner, P. Bronnimann, D.J. Carter & L. Zaninetti (1982)- *Triasina hantkeni* Majzon, 1954 (Foraminifere), dans le Trias superieur de la Tethys (Europe et Asie). Archives Sciences Geneve 35, 2, p. 137-142.
('Triasina hantkeni Majzon 1954 (foraminifer), in the Upper Triassic of the Tethys (Europe and Asia)'. Norian-Rhaetian small benthic foram Triasina, known in Tethys from Europe to E Indonesia (Seram))

Ando, H. (1987)- Paleobiological study of the Late Triassic bivalve *Monotis* from Japan. The University of Tokyo Museum Bull. 30, p. 1-110.
(online at: http://umdb.um.u-tokyo.ac.jp/DKankoub/Bulletin/no30/no30000.html)
(Monograph of Late Triassic 'flat clam' Monotis in Japan, with discusssion of global distribution)

Carter, E.S. (2007)- Global distribution of Rhaetian radiolarian faunas and their contribution to the definition of the Triassic-Jurassic boundary. In: S.G. Lucas & J.A. Spielmann (eds.) The global Triassic, New Mexico Museum of Natural History and Science Bulletin 41, p. 27-31.
(online at: http://paleo.cortland.edu/globaltriassic/Bull41/09-Carter%20(radiolarians).pdf)
(Includes comparison with Rhaetian radiolarian faunas from Meto River, West Timor as described by Rose (1994). Assemblages not typically Tethyan, but mix of cosmopolitan taxa and species with stronger affinities to Japan and Philippines)

Chablais, J. (2010)- Sedimentology and biostratigraphy of the Upper Triassic atoll-type carbonates of the Sambosan Accretionary Complex (Panthalassan domain; Japan). Doct. Thesis Universite de Geneve, Sc.4212, p. 1-204.
(online at: https://archive-ouverte.unige.ch/unige:8438)
(Part of series of papers on U Triassic (Norian-Rhaetian) carbonates from limestone-capped seamount in Sambosan Accretionary Complex. With typical 'Tethyan' Late Triassic benthic foram assemblages, incl. Aulotortus friedli, etc. (related assemblages also in Timor; JTvG))

Chablais, J., R. Martini, F. Kobayashi, G.M. Stampfli & T. Onoue (2011)- Upper Triassic foraminifers from Panthalassan carbonate buildups of Southwestern Japan and their paleobiogeographic implications. Micropaleontology 57, 2, p. 93-124.
(online at: https://www.researchgate.net/publication/230776641_Upper_Triassic_foraminifers_from_Panthalassan_carbonate_buildups_of_Southwestern_Japan_and_their_paleobiogeographic_implications)
(60 species of foraminifera in U Triassic 'Panthalassan' seamount atoll-type carbonates in Late Jurassic- E Cretaceous Sambosan Accretionary Complex of SW Japan. Four associations: (1) lagoonal with common Aulotortidae, Nodosariidae and Endotebidae; (2) Back- and fore-reef: mainly Duostomina and Variostoma; (3) reefal: Galeanella, Hoyenella, Ophthalmidium and Cucurbita; (4) Shoal facies rich in ooids and near-

monospecific Pilammina sulawesiana association (as also known from E Sulawesi). Carnian-Norian age similar to Tethyan carbonate platform/ reefs in European Alps, Asinepe Limestone in Seram, etc.. Foraminifera of Tethyan tropical affinity, suggesting paleoposition at low-middle latitude in S Hemisphere, in agreement with presence of endemic foraminifers and corals of Timor and Sulawesi. Sambosan AC seamounts moved >15000 km to be accreted against Asian blocks, which fits with velocity/direction of plates during Neotethys ridge opening)

Chablais, J., R. Martini, E. Samankassou, T. Onoue & H. Sano (2010)- Microfacies and depositional setting of the Upper Triassic mid-oceanic atoll-type carbonates of the Sambosan Accretionary Complex (southern Kyushu, Japan). Facies 56, p. 249-278.
(Of interest for comparison with Triassic 'Fatu Limestones' of Timor;also Seram?; JTvG)

Chonglakmani, C. & J.A. Grant-Mackie (1984)- Handbook of Triassic index fossils (preliminary). Dept. Mineral Resources (DMR) , Bangkok, p. 1-21.
(online at: http://library.dmr.go.th/Document/DMR_Technical_Reports/1984/24271.pdf)
(Selection of age-significant Triassic microfossils from Thailand and adjacent regions. Ammonites and thin-shelled molluscs (Daonella, Halobia, Monotis) most important. With 5 plates and range chart)

Cirilli, S. (2010)- Upper Triassic- lowermost Jurassic palynology and palynostratigraphy: a review. In: S.G. Lucas (ed.) The Triassic timescale, Geological Society, London, Special Publ. 334, p. 285-314.
(Late Triassic- E Jurassic palynostratigraphy and latitudinal control on distribution of 'Onslow' (rel. warm, southern Tethys margin) vs. 'Ipswich' (cool, high-latitude Gondwanan) microfloral provinces in S Hemisphere)

De Franceschi, D. & C. Vozenin-Serra (1997)- The Upper Vietnamese Triassic flora: palaeogeographical significance. Comptes Rendus Academie Sciences, Paris, ser. 2, 324, 4, p. 333-340.
(Vietnamese Triassic flora belongs to coastal floristic assemblage of SW Pacific. Affinities of plant-fossil assemblage with 'Krusin flora' of NW Borneo)

Diener, C. (1918)- Nachtrage zur Dibranchiatenfauna der Halstatter Kalke. Jahrbuch Geologischen Reichsanstalt, Wien, 1918, 68, 3, p. 475-492
(online at: https://opac.geologie.ac.at/ais312/dokumente/JB0683_475_A.pdf)
('Additions to the Dibranchiate fauna of the Halstatter Limestone'. Mainly comparison of Late Triassic belemnoid faunas from Timor and North Calcareous Alps in Austria. In both areas very similar thin-ribbed Aulacoceras sulcatum group. Prefers to maintain Aulacoceras timorense Wanner as separate species)

Dobruskina, I.A. (1994)- Triassic floras of Eurasia. Osterrreichische Akademie der Wissenschaften, Erdwissenschaftlichen Kommissionen, Band 10, Springer Verlag, Vienna, p. 1-422.
(online at: http://jurassic.ru/pdf/Dobruskina1994.pdf)
(Extensive review of Triassic plant fossils from Europe and Asia. Brief discussions on Far East Asia, incl. Triassic floras from Thailand (Norian- Rhaetian, near base of Khorat Group), Vietnam (Norian-Rhaetian 'Tonkin flora') and Sarawak (Carnian 'Krusin flora', near basal conglomerates of Halobia clastics series))

Dolby, J.H. & B.E. Balme (1976)- Triassic palynology of the Carnarvon Basin, Western Australia. Review Palaeobotany Palynology 22, p. 105-168.
(Five Triassic palynological assemblage zones in wells from Carnarvon Basin: I. Kraeuselisporites saeptatus (Griesbachian-Smithian, II. Tigrisporites playfordii (Smithian-Anisian), III. Staurosaccites quadrifidus (Anisian-Carnian), IV. Samaropollenites speciosus (Carnian) and V. Minutosaccus crenulatus (Carnian-?Norian). Provincialism in M-L Triassic floras:(1) Onslow microflora on NW Shelf, with mixed Gondwanan-European elements; (2) Ipswich microflora: less diverse Falcisporites-dominated assemblages in E and S Australia; European elements not present)

Foster, C.B., B.E. Balme & R. Helby (1994)- First record of Tethyan palynomorphs from the Late Triassic of East Antarctica. BMR. J. Australian Geology Geophysics 15, p. 239-246.
(online at: www.ga.gov.au/corporate_data/49409/Jou1994_v15_n2.pdf)

(Remarkable 'Onslow-type' ('Tethyan') Norian microflora from Prince Charles Mts, E Antarctica, with Minutosaccus crenulatus, Ovalipollis ovalis, Samaropollenites speciosus, Falcisporites australis, etc. (also known from Australian NW Shelf, Timor, etc; JTvG.))

Gardin, S., L. Krystyn, S. Richoz, A. Bartolini & B. Galbrun (2012)- Where and when the earliest coccolithophores? Lethaia, 10.1111, p. 1-17.
(First coccoliths appear in Late Triassic, with oldest species Crucirhabdus minutus and Prinsiosphaera triassica appearing in latest Norian. Across Norian-Rhaetian boundary increase in abundance of Prinsiosphaera triassica and appearance of Euconusphaera zlambachensis (two most important Rhaetian pelagic carbonate producers). Both present on Timor and the Wombat Plateau (off NW Australia) (also on Seram?; JTvG))

Goudemand, N., M.J. Orchard, H. Bucher & J. Jenks (2012)- The elusive origin of *Chiosella timorensis* (conodonts, Triassic). Geobios 45, 2, p. 199-207.
(online at: https://www.zora.uzh.ch/id/eprint/57300/1/1-s2.0-S0016699511000969-main.pdf)
(First occurrence of conodont Chiosella timorensis is good marker for Anisian age (species originally described as Gondolella timorensis from Lacan, Manatuto district, Timor, by Nogami (1968))

Grant-Mackie, G.A. (1975)- The stratigraphy and taxonomy of the Upper Triassic bivalve *Monotis* in New Zealand. Ph.D. Thesis University of Auckland, p. 1-380.
(online at: https://researchspace.auckland.ac.nz/handle/2292/2580)
(20 taxa recognized in New Zealand Late Triassic bivalve genus Monotis (of interest for comparisons with M-L Triassic Monotis species from Borneo, Buton, Seram, etc.; JTvG))

Grant-Mackie, G.A. (1978)- Subgenera of the Upper Triassic bivalve *Monotis*. New Zealand J. Geology Geophysics 21, 1, p. 97-111.
(online at: www.tandfonline.com/doi/abs/10.1080/00288306.1978.10420726)
(Proposal of five subgenera of Monotis. Monotis s.s. resembles salinaria group, M. (Entomonotis) includes M. ochotica, subcircularis, and zabaikalica groups; M. (Eomonotis) typha group and M. (Maorimonotis))

Hasibuan, F. (2010)- The Triassic marine biota of Eastern Indonesia and its interregional and global correlation: a review. Jurnal Geologi Indonesia (IJOG) 5, 1, p. 31-47.
(online at: https://ijog.geologi.esdm.go.id/index.php/IJOG/article/view/91/91)
(Marine macrofossil biostratigraphy of Triassic in Indonesia. Mainly on ammonoids from Triassic of Timor (E-M and Late Triassic) and Misool (M-L Triassic), and correlations with regions outside Indonesia)

Hasibuan, F. & Purnamaningsih (1998)- Pre-Tertiary biostratigraphy of Indonesia. In: J.L. Rau (ed.) Proc. 34th Session Coord. Comm. Coastal Offshore Geoscience Programmes E and SE Asia (CCOP), Taejon, Korea 1997, 2, Technical Reports, p. 40-54.
(Review of Paleozoic- Mesozoic marine macrofossil biostratigraphy of Indonesia, particularly in Misool island)

Hautmann, M. (2001)- Die Muschelfauna der Nayband-Formation (Obertrias, Nor- Rhat) des ostlichen Zentraliran. Beringeria 29, p. 1-181.
(online at: http://opus.bibliothek.uni-wuerzburg.de/frontdoor/index/index/docId/1817)
('The bivalve fauna of the Nayband-Formation (U Triassic, Norian-Rhaetian) of east-central Iran'. Well-preserved bivalve fauna with >100 species. Paleogeographic analysis of Tethyan bivalve faunas suggests E Tethys province, with northern (Iran, Yunnan, Vietnam, Burma, W Sumatra) and southern subprovinces (Lhasa-block, E Indonesia terranes; attached to Gondwana until end Triassic and separated later by opening of 'Ceno-Tethys' Ocean (= Mesotethys?; JTvG))

Hautmann, M., M.J. Benton & A. Tomasovych (2008)- Catastrophic ocean acidification at the Triassic-Jurassic boundary. Neues Jahrbuch Geologie Palaontologie, Abhandlungen 249, p. 119-127.
(On end-Triassic extinction of reefal organisms and end of carbonate deposition at Triassic- Jurassic boundary in many parts of world, caused by ocean acidification, tied to volcanic degassing (this may be a potential

explanation why Late Triassic carbonate platform limestones are rel. widespread in E Indonesia, but Early Jurassic carbonates are rare to absent; JTvG))

Hautmann, M. (2012)- Extinction: end-Triassic mass extinction. In: eLS online, John Wiley & Sons, p. 1-10.
(On mass extinction at end of Triassic, at ~200 Ma. This event eliminated conodonts and nearly annihilated corals, sphinctozoan sponges and ammonoids. Probably caused by volcanic activity of Central Atlantic Magmatic Province. Lead to virtual absence of reef systems for nearly 10 Myrs in E Jurassic).

Helby, R., V.D. Wiggins & G.J. Wilson (1987)- The circum-Pacific occurrence of the Late Triassic dinoflagellate *Sverdrupiella*. Australian J. Earth Sciences 34, p. 151-152.
(Late Triassic dinoflagellate genus Sverdrupiella Bujak and Fisher widespread, abundant and relatively diverse in Norian strata. Reported from Seram, E Indonesia, in Norian Kanikeh Fm, associated with undescribed dinoflagellate suite including suessioids and Heibergella (=Hebecysta balmei zone?; JTvG). Circum-Pacific distribution of Sverdrupiella in Late Triassic similar to distribution of Late Triassic bivalve Monotis)

Hinde, G.J. (1908)- Radiolaria from Triassic and other rocks of the Dutch East Indian Archipelago. In: R.D.M. Verbeek, Molukkenverslag. Geologische verkenningstochten in het oostelijke gedeelte van den Nederlandsch Oostindische Archipel. Jaarboek Mijnwezen Nederlandsch Oost-Indie 37 (1908), Wetenschappelijk Gedeelte, p. 694-736.
(Radiolaria from Timor, Savu, Ceram, Sulawesi, Buru and Mangoli collected by Verbeek, probably mainly of Late Triassic-Jurassic age. 83 species identified, 74 new. Richest assemblages found in Halobia-Daonella-bearing cherty limestones from Rote and Savu and Halobia limestone from Timor. Fewer, but similar species in loose chert pebbles collected at Seram and E Sulawesi)

Ichikawa, K. (1958)- Zur Taxonomie und Phylogenie der triadischen „Pteriidae" (Lamellibranchiata) mit besonder Berucksichtigung der Gattungen *Claraia, Eumorphotis, Oxytoma* und *Monotis*. Palaeontographica, A111, 5-6, p. 131-212.
('On the taxonomy and phylogeny of the Triassic 'Pteriidae' (Lamellibranchiata), with particular emphasis on the genera Claraia, Eumorphotis, Oxytoma und Monotis'. Includes description of new species Monotis (Entomonotis) timorica from Timor (which is same as Monotis subcircularis Gabb; Westermann 1962))

Ishibashi, T. (1975)- Some Triassic ammonites from Indonesia and Malaysia. In: T. Kobayashi & R. Toriyama (eds.) Geology and Palaeontology of Southeast Asia, University of Tokyo Press, 16, p. 45-56

Jasin, Basir (2018)- Radiolarian biostratigraphy of Malaysia. Bull. Geological Society Malaysia 65, p. 45-58.
(online at: https://gsmpubl.files.wordpress.com/2018/08/bgsm201805.pdf)
(Two types of radiolarian cherts in Malaysia:bedded chert and chert blocks. Bedded cherts in Peninsular Malaysia mainly in Western Belt (Kubang Pasu, Kenny Hill and Semanggol Fms). In Sabah in Sabah Ophiolite Complex. In Sarawak bedded cherts in Serian Volcanic Fm and basal part of Pedawan Fm. Chert blocks mainly in Bentong Raub Suture Zone of Peninsular Malaysia and melanges of Sabah and Sarawak. Radiolarians from Peninsular Malaysia Late Devonian-Triassic, with 16 biozones identified. Radiolarians from Sabah and Sarawak E Jurassic- Cretaceous ages. Five hypersiliceous periods, related to volcanism: Late Devonian-E Carboniferous, Permian, Triassic, E Jurassic and Late Jurassic-Cretaceous)

Kanmera, K. (1964)- Triassic coral faunas from the Konose Group in Kyushu. Memoirs Faculty of Science, Kyushu University, Ser. D, Geology, 15, 1, p. 117-147.
(online at: https://catalog.lib.kyushu-u.ac.jp/opac_download_md/1543615/15_01_p117.pdf)
(Corals from U Triassic U Konose Gp in the southern accretionary belt of Japan (Pacific seamount limestones). associated with common basic volcanoclastics. Close affinities to (Tethyan) Norian-Rhaetian corals of Austrian Alps and from Timor (author mentions species previously known only from Timor;like Montlivaltia timorica, M. norica, JTvG))

Kobayashi, F., R. Martini & L. Zaninetti (2005)- Anisian foraminifers from allochthonous limestones of the Tanoura formation (Kurosegawa Terrane, West Kyushu, Japan). Geobios 38, 6, p. 751-763.

(35 species of M Triassic foraminifera in allochthonous oolitic limestone blocks in Carnian Tanoura Fm in Kurosegawa Terrane of W Kyushu. Blocks of Anisian in age, based on Pilammina densa and Meandrospira dinarica, associated with Triadodiscus, Aulotortus, etc. Terrane was part of N Gondwana margin, then isolated eastwards before E Cretaceous amalgamation with S China in E Cretaceous)

Kobayashi, T. & T. Kimura (1944)- A study on the radiolarian rocks. J. Faculty of Science, Imperial University of Tokyo, Section 2, 7, 2, p. 75-187.
(Includes review of Paleozoic- Mesozoic radiolaria data from Sumatra, Kalimantan, etc. Kalimantan Danau Fm radiolaria transitional between Jurassic and Carboniferous faunas, therefore possibly more likely of Triassic age. Cherts in Tuhul Fm of S Sumatra are M-L Triassic or Permo-Triassic in age)

Kobayashi, T. & M. Tamura (1983)- On the Oriental Province of the Tethyan Realm in the Triassic Period. Proc. of the Japan Academy, Series B, 59, 7, p. 203-206.
(online at: https://www.jstage.jst.go.jp/article/pjab1977/59/7/59_7_203/_pdf/-char/en)
(Short paper on provinciality in Triassic bivalves. In addition to cosmopolitans, many Triassic bivalve genera are distributed from Alps to China, Indochina and USSR. Oriental Province of Tethys with species indigenous to E and SE Asia. Stretches from Kashmir, Burma, S China, Malay Peninsula, to E Indonesia. Krumbeckia Diener 1908 is known from Upper Triassic in Sumatra and Yunnan. No maps)

Kobayashi, T. & M. Tamura (1983)- The Arcto-Pacific Realm and the Trigoniidae in the Triassic Period. Proc. of the Japan Academy, Series B, 59, 7, p. 207-210.
(online at: https://www.jstage.jst.go.jp/article/pjab1977/59/7/59_7_207/_pdf/-char/en)
(Family Trigoniidae evolved from Myophoriidae polyphyletically in M and Late Triassic. Triassic bivalves of New Zealand, New Caledonia and New Guinea constitute distinct fauna. U Triassic cosmopolitan genus Monotis with paleogeographically restricted species; Borneo is junction of three Monotis seas: (1) M. salinaria group, distributed from Europe-Himalaya to Borneo; (2) M. ochotica group from Siberia to Borneo through Japan and (3) M. subcircularis group from Canada to New Zealand and Borneo through E Pacific coast. No figures)

Kobayashi, T. & M. Tamura (1984)- The Triassic Bivalvia of Malaysia, Thailand and adjacent areas. In: T. Kobayashi et al. (eds.) Geology and Palaeontology of Southeast Asia 25, University of Tokyo Press, p. 201-227.
(Review of Triassic bivalves Claraia, Daonella-Halobia, Monotis, Trigoniaceae (incl. Myophoria, Costatoria), megalodontids in Malay Peninsula, Thailand and Sumatra)

Koike, T. (1984)- Summary of Triassic conodonts of Southeast Asia. In: T. Kobayashi, R. Toriyama & W. Hashimoto (eds.) Geology and Palaeontology of Southeast Asia, University of Tokyo Press, 25, p. 295-302.
(Brief review of Triassic conodont faunas in Malay Peninsula, Thailand and Indonesia (Carnian of Lake Toba, Sumatra; Anisian-Ladinian and Carnian and W Timor; Norian of Timor Leste))

Kristan-Tollmann, E. (1988)- A comparison of Late Triassic agglutinated foraminifera of Western and Eastern Tethys. Abhandlungen Geologischen Bundesanstalt, Vienna, 41, p. 245-253.
(online at: www.landesmuseum.at/pdf_frei_remote/AbhGeolBA_41_0245-0253.pdf)
(Examples of common species of Triassic arenaceous and calcareous agglutinated foraminifera across Tethys realm (mainly from Austrian Calcareous Alps, PNG Kuta Limestone and Timor). Timor fauna includes Verneulinoides mauritii and Variostoma helictum)

Kristan-Tollmann, E. (1991)- Triassic Tethyan microfauna in Dachstein limestone blocks in Japan. In: T. Kotaka, J.M. Dickins et al. (eds.) Proc. International Symposium in Shallow Tethys 3, Sendai 1990, Saito Hoon Kai Special Publ. 3, p. 35-49.
(First description of U Triassic foraminifera from Panthalassan seamount reefal limestone in Sambosan accretionary complex in Japan; see also Chablais et al. 2008)

Kristan-Tollmann, E. (1995)- Weitere Beobachtungen an Rhatischen nannofossilien der Tethys. Geologisch-Palaontologische Mitteilungen Innsbruck 20, p. 1-11.

(online at: www2.uibk.ac.at/downloads/c715/gpm_20/20_001-011.pdf)
('Additional observations on Rhaetian nannofossils of the Tethys'. Eoconusphaera zlambachensis and Prinsiosphaera triassica are most common species in uppermost Triassic of Calcareous Alps; also present off Wombat Plateau, NW Australia, confirming Tethys-wide distribution)

Kummel, B. (1969)- Ammonoids of the Late Scythian (Lower Triassic). Bull. Museum Comparative Zoology, Harvard University, 137, p. 311-701.
(online at: www.archive.org/details/bulletinofmuseum13719681969harv)
(Treatise on Early Triassic/ Scythian ammonoids, including material from Timor (p. 349-351), mainly from 'Block E near Nifoekoko', described previously by Welter (1922). Contains Hungarites, Pronorites spp, etc.)

Lehrmann, D.J., J.Y. Wei & P. Enos (1998)- Controls on facies architecture of a large Triassic carbonate platform: the Great Bank of Guizhou, Nanpanjiang Basin, south China. Journal of Sedimentary Research 68, p. 311-326.
(Model of E-M Triassic carbonate platform deposition on S China block. E Triassic low relief bank with cyanobacterial limestones in platform interior and oolites along margins. M Triassic higher relief platform margins composed of Tubiphytes boundstone, with cyclic tidal flats formed ine platform interior)

Mantle, D.J., J.B. Riding & C. Hannaford (2020)- Late Triassic dinoflagellate cysts from the Northern Carnarvon Basin, Western Australia. Review Palaeobotany Palynology 281, 104254, p. 1-53.

McRoberts, C.A. (2008)- Rhaetian (Late Triassic) *Monotis* (Bivalvia: Pectinoida) from the eastern Northern Calcareous Alps (Austria) and the end-Norian crisis in pelagic faunas. Palaeontology 51, 3, p. 721-735.
(Species of marine bivalves of pectinoid genus Monotis provide useful biochronologic indices for Late Triassic (M Norian-earliest Rhaetian). Profound extinction event in pelagic realm at Norian-Rhaetian boundary where ~15 monotids became extinct. Surviving Monotis dwarfed compared to Norian predecessors)

McRoberts, C.A. (2010)- Biochronology of Triassic bivalves. In: S.G. Lucas (ed.) The Triassic Timescale, Geological Society, London, Special Publ. 334, p. 201-219.
(General zonation scheme for Triassic based on bivalves Claraia, Peribositria, Enteropleura, Daonella (M Triassic), Halobia (Carnian-M Norian), Eomonotis and Monotis (Late Norian). Widely distributed across Tethys, Panthalassa and Boreal regions)

Nicoll, R.S. & C.B. Foster (1998)- Triassic biozonation and stratigraphy, 1998 Chart 20. Australian Geological Survey Organisation (AGSO), 1p. *(chart)*
(online at: www.ga.gov.au/corporate_data/76687/Chart_20_NWS_Triassic.pdf)

O'Dogherty L., E.S. Carter, P. Dumitrica, S. Gorican, P. De Wever, A. Hungerbuhler, A.N. Bandini & A. Takemura (2009)- Catalogue of Mesozoic radiolarian genera. Part 1: Triassic. Geodiversitas 31, 2, p. 213-270.
(online at: https://www.researchgate.net/publication/232694009_Catalogue_of_Mesozoic_radiolarian_genera_Part_1_Triassic)
(Systematic revision of Triassic radiolarian genera, validating 282 genera, of which only 30 genera crossed Triassic-Jurassic boundary

O'Dogherty L., E.S. Carter, S. Gorican & P. Dumitrica (2010)- Triassic radiolarian biostratigraphy. In: S.G. Lucas (ed.) The Triassic Timescale, Geological Society, London, Special Publ. 334, p. 163-200.
(General discussion of Triassic radiolarian biostratigraphy. Work on Triassic in Indonesia limited to Sashida et al. (1999) on Timor. Many studies on Thailand Triassic)

Payne, J.L. & B. van de Schootbrugge (2007)- Life in Triassic oceans: links between planktonic and benthic recovery and radiation. In: P. Falkowski & A.H. Knoll (eds.) Evolution of primary producers in the sea, Academic Press, Amsterdam, p. 165-189.
(Review of faunal trends through Triassic. E Triassic global reef gap after end-Permian extinctions, commonly associated with black shale. Increase in coral and algal diversity through M Triassic, but reefs dominated by

Tubiphytes. E-M Carnian reefs dominated by Porifera, Norian-Rhaetian reefs dominated by corals. E Jurassic is another reef gap, again with common black shale)

Peybernes, C. (2016)- Upper Triassic mid-oceanic shallow water ecosystems of the Panthalassa Ocean: insights from the Sambosan Accretionary Complex, Southwest Japan. Doct. Thesis Universite de Geneve, Sc 4914, p. 1-229.
(online at: https://archive-ouverte.unige.ch/unige:84250)
(U Triassic carbonates of Sambosan Accretionary Complex in SW Japan: (1) limestone clasts embedded in volcaniclastic matrix with microbialite-rich Ladinian?- Lower Carnian reef biota; (2) sponge-dominated Late Carnian- Norian (Rhaetian?) reefs (built on top of basalts of low-latitude Panthalassic seamount). Paleogeographic affinities with S. Tethys. Limestone commonly as clasts in volcanoclastic breccias, probably mass-movement deposits from seamount collapse in mid-oceanic realm (very few ooids?))

Peybernes, C., J. Chablais & R. Martini (2015)- Upper Triassic (Ladinian?-Carnian) reef biota from the Sambosan Accretionary Complex, Shikoku, Japan. Facies 61, 4, p. 1-27.
(M-L Triassic (Ladinian?-Carnian) reef limestone from Sambosan Accretionary Complex, Shikoku Island, SW Japan, with scleractinian corals, calcified sponges, calcareous algae, foraminifera and microproblematica (older than previously identified reef limestones in Sambosan Complex))

Peybernes, C., J. Chablais, T. Onoue, G. Escarguel & R. Martini (2016)- Paleoecology, biogeography, and evolution of reef ecosystems in the Panthalassa Ocean during the Late Triassic: Insights from reef limestone of the Sambosan Accretionary Complex, Shikoku, Japan. Palaeogeogr. Palaeoclim. Palaeoecology 457, p. 31-51.
(U Triassic sponge-coral- Tubiphytes reef limestone from Sambosan Accretionary Complex at Shikoku Island, Japan, with two types of reefs, Ladinian?- E Carnian and Late Carnian-Rhaetian? Strong paleobiogeographic affinity of Late Triassic W Panthalassa reef biota with those of S Tethys Ocean)

Rigaud, S., R. Martini, R. Rettori & G.D. Stanley (2010)- Stratigraphic potential of the Upper Triassic benthic foraminifers. Albertiana 38, p. 34-39.
(online at: http://archive-ouverte.unige.ch/unige:6737)
(Summary of rich Carnian-Norian benthic foraminiferal assemblages in isolated limestone remnant of Panthalassa Ocean in N Wallowa Mts, Oregon. Foraminifera ~75% well-known Tethyan species, with comparable stratigraphic distribution, incl. Triasina oberhauseri, Aulotortus tumidus, A. communis, A. tenuis, etc. Also with Norian Heterastridium conglobatum)

Rigaud, S., R. Martini & R. Rettori (2013)- A new genus of Norian involutinid foraminifers: its morphological, biostratigraphic, and evolutionary significance. Acta Palaeontologica Polonica 58, 2, p. 391-405.
(online at: www.app.pan.pl/archive/published/app58/app20110072.pdf)
(New genus name Aulosina for Late Triassic (Norian- E Rhaetian) involutinid foram Triasina oberhauseri, morphologically transitional between Aulatortus and (Rhaetian) Triasina hantkeni)

Schafer, P. & J.A. Grant Mackie (1998)- Revised systematics and palaeobiogeography of some Late Triassic colonial invertebrates from the Pacific region. Alcheringa 22, 1-2, p. 87-122.
(Revision of U Triassic colonial organisms from New Zealand, New Caledonia, Timor, etc. Heterastridium conglobatum, a hydrozoan of uncertain affinity and possible pelagic lifestyle, is known from Norian of Tethys (Hallstatt Lst in Alps, Middle East, etc.), Timor, New Caledonia, New Zealand and W North America. 'Monotrypella timorica' is calcareous demosponge)

Stanley, G.D. & T. Onoue (2015)- Upper Triassic reef corals from the Sambosan Accretionary Complex, Kyushu, Japan. Facies 61, 1, p. 1-27.
(Ten U Triassic coral taxa, incl. Retiophyllia, from limestones of Sambosan accretionary complex of Japan, with remains of reefs and carbonate sediment deposited on Pacific volcanic atolls. High degree of endemism; some paleogeographic connection with W Tethys, Pamir Mts and Timor (Craspedophyllia ramosa Roniewicz))

Schatz, W. (2004)- Revision of the subgenus *Daonella* (Arzelella) (Halobiidae; Middle Triassic). Journal of Paleontology 78, 2, p. 300-316.
(Mainly taxonomic revision of M Triassic Daonella (thin-shelled 'paper mollusc'). Includes Late Ladinian- E Carnian species Daonella indica Bittner 1899, known from SE Europe, through Turkey, Pamir, Himalayas (Spiti, etc.), SW China, Malay Peninsula (Pahang), Timor, Japan (in N Thailand and Malay Peninsula in area of Paleotethys suture; JTvG))

Thenius, E. (1980)- Zum Problem der „zirkumpazifischen" und der Tethys-Verbreitung mariner Evertebraten in der Trias. Annalen Naturhistorischen Museum Wien 83, p. 285-301
(online at: www.landesmuseum.at/pdf_frei_remote/ANNA_83_0285-0301.pdf)
('On the problem of 'Circum-Pacific' and the Tethys distribution of marine invertebrates in the Triassic'. Figure 2 shows distribution of Norian bivalve Monotis, modified from Westermann (1973), showing Indonesia assemblages with 'Arctic- W Pacific' Monotic ochotica and M. subcircularis in NW Kalimantan, while 'Tethyan' Monotis salinaria and M. subcircularis are present on Timor)

Vachard, D. & H. Fontaine (1988)- Biostratigraphic importance of Triassic foraminifera and algae from South-East Asia. Revue de Paleobiologie, Geneve, 7, 1, p. 87-98.
(Triassic forams from Thailand, NW Malay Peninsula, Indonesia (Sibaganding limestone of Sumatra, Seram), Philippines (Malajon Island), Myanmar, Vietnam)

Van Voorthuysen, J.H. (1940)- Beitrag zur Kenntnis des inneren Baus von Schale und Sipho bei triadischen Ammoniten. Doct. Thesis, University of Amsterdam, p. 1-143.
(Study of internal structure of U Triassic ammonites from Timor (Tropitida, Placites spp., Cladiscitidae, Phylloceratidae and Arcestidae))

Vozenin-Serra, C. (1971)- Notes sur des bois Mesozoiques et Cenozoique du Viet-nam et du Cambodge. Archives Geologiques Vietnam 14, p. 1-47.
('Notes on the Mesozoic and Cenozoic fossil wood from Vietnam and Cambodia')

Vozenin-Serra, C. (1977)- Contribution a l'etude de la paleoflore du Sud-est Asiatique (Cambodge, Laos, Vietnam). These Doctoral Etat., Universite de Paris, p. 1-310. *(Unpublished)*
('Contribution to the study of the paleoflora of SE Asia (Cambodia, Laos, Vietnam)')

Vozenin-Serra, C. (1983)- Etat de nos connaissances sur les flores mesozoiques du Sud-Est Asiatique. Comptes Rendus 108e Congres National des Societes Savantes, Grenoble 1983, 1, Sciences de la Terre, 2, p. 101-116.
('Status of our knowledge of the Mesozoic floras of SE Asia'. Review of Triassic-Cretaceous floras, mainly from Vietnam, Thailand, etc. For Indonesian region only mentions W Sarawak Krusin flora of Kon'no 1972, which is part of Late Triassic 'Dictyophyllum- Clathropteris floral province' of SW Pacific at end-Triassic (also known from Vietnam, Thailand, W Japan))

Vozenin-Serra, C. (1984)- Etat de nos connaissances sur les flores du Paleozoique superieur et du Mesozoique du Sud-Est Asiatique. Interpretations paleogeographiques. Memoires Societe Geologique France 147, p. 169-181.
('Status of our knowledge of the Upper Paleozoic and Mesozoic floras of SE Asia and paleogeographic interpretations')

Westermann, G.E.G. (1973)- Species distribution of the world-wide Triassic pelecypod *Monotis* Bronn. Proc. 22nd International Geological Congress, India 1964, Section 8, p. 374-389.

Zammit, M. (2010)- A review of Australasian ichthyosaurs. Alcheringa 34, 3, p. 281-292.
(Ichthyosaur fossils recorded from M Triassic of Timor (Mixosaurus sp.), from U Triassic of New Caledonia (Shonisauru) and Lower Cretaceous of Australia and New Zealand (Platypterygius))

Zaninetti, L. (1976)- Les foraminiferes du Trias, essai de synthese et correlation entre les domaines mesogeens europeen et asiatique. Rivista Italiana Paleontologia Stratigrafia 82, p. 1-258.
(Synthesis of Triassic foraminifera and correlation between European and Asian domains)

X.5. Paleozoic

Numerous additional papers with Paleozoic biostratigraphy content that are case studies specific to certain areas, are listed under those areas, and are not repeated here.

This chapter does include several papers on Paleozoic faunas-floras outside Indonesia that may help in the interpretation of related occurrences in Indonesia.

Permian stratigraphies and fossils are particulary useful for distinguishing higher latitude Gondwanan terranes (with Early Permian glacial deposits, Glossopteris flora, Eurydesma mollusc fauna,etc.), separated by the Paleotethys Ocean from low latitude Asian (Cathaysian) terranes during Permian time, as well as the 'Cimmerian ribbon terranes' (Sibumasu- Baoshan- Qiangtang) that rifted off Gondwana during the Early Permian opening of the Mesotethys Ocean.

Remarkable occurrences of 'East Paleotethys- Panthalassan' low-latitude Middle? Permian low-latitude fusulinid limestones deposited on basic seamount volcanics, now found from West Sumatra to Timor(?), Japan, New Zealand and NW North America

Aitchison, J.C., N. Suzuki, M. Caridroit, T. Danelian & P. Noble (2017)- Paleozoic radiolarian biostratigraphy. Geodiversitas 39, 3, p. 501-532.
(online at: https://sciencepress.mnhn.fr/sites/default/files/articles/pdf/geodiversitas-2017-n3a5-biostratigraphy-of-paleozoic-radiolarian-genera.pdf)
(Paleozoic (Cambrian-Permian) radiolarian biostratigraphy, with references to papers from Thailand, etc. (see also companion papers in same Geodiversitas volume 39, by Noble, Caridroit))

Archbold, N.W. (1983)- West Australian Permian brachiopoda: their taxonomy, biostratigraphy and provincialism; with an Appendix of published articles including a description of Permian brachiopoda from Irian Jaya, Indonesia. Ph.D. Dissertation, University of Melbourne, 2 vols.

Archbold, N.W. (2001)- Permian Productida of Australasia: palaeobiogeographical and palaeoclimatological implications. In: S.L. Long et al. (eds.) Brachiopods, Chapter 37, CRC Press, p. 363-371.
(Permian Productid brachiopods few genera in common between Westralian (rel. warm, S Tethys margin) and Austrazean (colder water, mainly endemics, strong links with New Zealand) provinces)

Archbold, N.W. & T. Hogeboom (2000)- Subsurface brachiopoda from borehole cores through the Early Permian sequence of the Carnarvon Basin, Western Australia: correlations with palynological biostratigraphy. Proc. Royal Society of Victoria 112, 1, p. 93-109.
(Early Permian brachiopods from five wells in onshore Carnarvon Basin, tied to spore-pollen zonation. Four earliest Permian brachiopod zones of W Australia (Lyonia lyoni, Trigonotreta occidentalis, Strophalosia irwinensis and Strophalosia jimbaensis zones, in ascending order) correlated with palynological zones (Granulatisporites confluens, Pseudoreticulatispora pseudoreticulata, Striatopodocarpites fusus, Didecitriletes byroensis and Microbaculispora trisina zones, in ascending order))

Asama, K., A. Hongnusonthi, J. Iwai, E. Kon'no, S.S. Rajah & M. Veeraburas (1975)- Summary of the Carboniferous and Permian plants from Thailand, Malaysia and adjacent areas. In: T. Kobayashi & R. Toriyama (eds.) Geology and Palaeontology of Southeast Asia, University of Tokyo Press, 15, p. 77-101.
(online at: http://library.dmr.go.th/library/DMR_Technical_Reports/1975/324.pdf)
(Oldest plants in SE Asia from Lower Carboniferous: known from Malaysia only (Kuantan flora, with Lepidodendron, on E Malaya Block). Five known Permian floras. Permian Jambi flora of Sumatra greatest similarities with Artinskian-Kungarian Shansi Fm of N China. Four other U Permian Cathaysian floras with Gigantopteris: Linggiu and Jengka floras from E Malay Peninsula; Phetchabun and Loei floras of N Thailand. Permian plants from W New Guinea are Gondwana-type flora, but some apparent Cathaysian elements)

Belasky, P. (1994)- Biogeography of Permian corals and the determination of longitude in tectonic reconstructions of the Paleopacific region. Canadian Society of Petroleum Geologists, Special Publ., p. 621-646.

(Mainly focused on American terranes. South China was center of diversity of Permian Tethyan coral province and was located near Permian Equator and W margin of Paleopacific Ocean)

BouDagher-Fadel, M.K. (2008)- The Palaeozoic larger benthic foraminifera: the Carboniferous and Permian. In: Evolution and geological significance of larger benthic foraminifera, Chapter 2, Developments in Palaeontology and Stratigraphy, Elsevier, 21, p. 39-118.
(General review of Paleozoic larger foraminifera, mainly Carboniferous-Permian fusulinids. End of Permian is major extinction event)

Bronnimann, P., J.E. Whittaker & L. Zaninetti (1978)- *Shanita*, a new pillared Miliocean foraminifera from the Late Permian of Burma and Thailand. Rivista Italiana Paleontologia 84, p. 63-92.
(New late Middle- Late Permian pillared miliolid Shanita amosi from W margin of Shan Plateau, E Myanmar. Species believed to be mainly characteristic of M Permian of Sibumasu/ Cimmerian terranes)

Broutin, J., J. Yu, X. Shi, W. Shu & X. Qing (2020)- Terrestrial palaeofloral succession across the Permian-Triassic boundary in the North and South China blocks: a brief review. Palaontologische Zeitschrift 94, p. 633-644.
(From Late Cisuralian (E Permian) to M Lopingian, N and S China blocks located in the equatorial domain. Both with homogeneous 'Cathaysian Palaeoflora', resulting from Cisuralian southern extension and diversification from northern cradle of Cathaysian plants to South. Progressive change in floral successions due to N-ward drift of N China block, while S China block remained in sub-equatorial domain)

Cantrill, R.C. (2003)- Aspects of Ordovician conodonts and the stratigraphy of Thailand, Malaysia, and Tasmania. Ph.D. Thesis, University of Tasmania, p. 1-158.
(online at: http://eprints.utas.edu.au/19158/1/whole_CantrillRobinCrawford2003_thesis.pdf)
(Incl. Ordovician conodonts from shallow tropical Lower Setul Lst from Langkawi Island (Tremadoc- Ashgill), Thung Song Lst of mainland Thailand and Ko Tarutao, Karmberg Lst of central S Tasmania)

Cantrill, R.C. & C. Burrett (2003)- The Greater Gondwana distribution of the Ordovician conodont *Panderodus nogamii* (Lee) 1975. Courier Forschungsinstitut Senckenberg, Frankfurt, 245, 1, p. 407-421.
(Panderodus nogamii (formerly Scolopodus nogamii), first described from N Korea, also in Lower Ordovician of Thailand, Malaysia, N and S China, Australia and Argentina. It ranged through Middle- early Late Ordovician and was restricted to shallow water carbonates in tropical- subtropical paleolatitudes of Greater Gondwana)

Caridroit, M., T. Danelian, L. O'Dogherty, J. Cuvelier, J.C. Aitchison, L. Pouille, P. Noble et al. (2017)- An illustrated catalogue and revised classification of Paleozoic radiolarian genera. Geodiversitas 39, 3, p. 363-417.
(online at: https://sciencepress.mnhn.fr/sites/default/files/articles/pdf/geodiversitas-2017-n3a3-catalogue-of-paleozoic-radiolarians_0.pdf)

Chapman, F. & W.J. Parr (1937)- On the discovery of fusulinid foraminifera in the Upper Palaeozoic of Northwest Australia. Victorian Naturalist 53, p. 175-179.
(Describe presence of fusulinid genera Verbeekina and Neoschwagerina in NW Australia. Identifications were later shown to be erroneous, they are fish remains (Crespin 1958, Quilty 1975) (still no fusulinid forams recorded yet from Australia, but are present on Timorisland and also in Birds Head of New Guinea; JTvG)

Cleal, C.J. (2017)- A global review of Permian macrofloral biostratigraphical schemes. In: S.G. Lucas & S.Z. Shen (eds.) The Permian Timescale, Geological Society, London, Special Publ. 450, p. 349-363.
(Incl. mention of Djambi flora of Sumatra, which may be regarded as transitional flora with taxa characteristic of both Euramerica (e.g. medullosaleans and marattialean ferns) and Cathaysia (e.g. Tingia, Cathaysiodendron) plant Paleokingdoms)

De D. Hornibrook, N. (1951)- Permian fusulinid foraminifera from the North Auckland Peninsula, New Zealand. Trans. Royal Society of New Zealand 79, 2, p. 319-321.

(online at: http://rsnz.natlib.govt.nz/volume/rsnz_79/rsnz_79_02_004230.pdf)
(Permian fusulinid foraminifera from limestone blocks associated with spilitic pillow lavas from Whangaroa Harbour, North Island, are first record of fusulines from New Zealand. Verbeekina, Neoschwagerina margaritae, Yabeina multiseptata probably of M Permian (Wordian- Capitanian) age (and signify non-Gondwana low latitude; JTvG))

De Neve, G.A. (1961)- Correlation of fusulinid rocks from southern Sumatra, Bangka, and Borneo, with similar rocks from Malaya, Thailand and Burma. Proc. 9th Pacific Science Congress, Bangkok 1957, Geology and Geophysics 12, p. 249. *(Abstract only)*
(Four occurrences of U. Paleozoic rocks with fusulinids in Indonesia, incl. Sumatra localities of limestone with Neoschwagerina and Fusulina spp. in Palembang area, S Sumatra, (3a) E of Bukit Pendopo, discovered by Keil and (3b) 18 km W of Palembang, in Sekaju area pebbles with fusulinids in Old Neogene conglomerate by Van Tuijn (1931))

Dickins, J.M. (1956)- Permian pelecypods from the Carnarvon Basin, Western Australia. Bureau Mineral Resources Geology Geophysics, Bull. 29, p. 1-42.
(online at: https://s3-ap-southeast-2.amazonaws.com/corpdata/224/Bull_029.pdf)
(Documentation of Permian bivalve molluscs from northern Gondwana margin in NW Australia)

Dickins, J.M. (1957)- Permian pelecypods and gastropods from the Carnarvon Basin, Western Australia. Bureau Mineral Resources Geology Geophysics, Bull. 41, p. 1-75.
(online at: https://d28rz98at9flks.cloudfront.net/173/Bull_063.pdf)
(More documentation of Permian bivalve molluscs from northern Gondwana margin in NW and W Australia. Including the supposedly cool-water genus Atomodesma spp.))

Dickins, J.M. (1963)- Permian pelecypods and gastropods from Western Australia. Bureau Mineral Resources Geology Geophysics, Bull. 63, p. 1-203.
(online at: https://d28rz98at9flks.cloudfront.net/173/Bull_063.pdf)
(>150 species of pelecypods and 60 gastropods from 'Gondwanan' Permian of Carnarvon Basin and Canning/ Fitzroy Basins. Lower Permian four stages, lower stage characterized by 'Eurydesma fauna')

Edgell, H.S. (2004)- Upper Devonian and Lower Carboniferous Foraminifera from the Canning Basin, Western Australia. Micropaleontology 50, p. 1-26.
(Foraminifera from U Devonian reef complex and overlying Lower Carboniferous in N Canning Basin of NW Australia 20 species of tournayellids and endothyrids. Striking resemblance to microfaunas of Russia, Kazakhstan and South China)

Fortey, R.A. & L.R.M. Cocks (1986)- Marginal faunal belts and their structural implications, with examples from the Lower Palaeozoic. J. Geological Society, London, 143, p. 151-160.
(Record of Ordovician (Llanvirn) graptolites in Heluk River, E Irian Jaya (4°25'S, 139°17'E). Assigned to isograptid biofacies and taken as evidence of Ordovician ocean margin here. Oldest fossils in Indonesia ?)

Foster, C.B. (1982)- Spore-pollen assemblages of the Bowen Basin, Queensland (Australia): their relationship to the Permian/Triassic boundary. Review Palaeobotany Palynology 36, p. 165-183.
(Five palynofloral zones in M Permian- E Triassic of Bowen Basin, from old to young: 'Upper Stage 5' (= Dulhuntyipora zone of authors; commonly associated with Glossopteris); Playfordiaspora crenulata Zone (M Permian; = youngest coal seams and with last known Glossopteris), Protohaploxypinus microcorpus Zone (also with P. samoilovichii), Lunatisporites pellucidus Zone (also contains Falcisporites= Alisporites?; latest Permian or E Triassic?) and Protohaploxypinus samoilovichii Zone (= E Triassic). No range charts (Morante 1996 suggests Base P. microcorpus is ~Base E Triassic = base of Mt Goodwin shale in Bonaparte basin)

Foster, C.B., G.A. Logan, R.E. Summons, J.D. Gorter & D.S. Edwards (1997)- Carbon isotopes, kerogen types and the Permian-Triassic boundary in Australia: implications for exploration. The APPEA Journal 37, p. 472-489.

(Permian- Triassic boundary characterized by massive extinction of marine fauna. In non-marine sections in E Australia, top of coal measures used as top Permian. Carbon isotopic (δ13C) shift of either organic matter or carbonates may be used to delimit P-T boundary)

Furnish, W.M. & B.F. Glenister (1970)- Permian ammonoid *Cyclolobus* from the Salt Range, West Pakistan. In: Stratigraphic boundary problems: Permian and Triassic of West Pakistan, p. 153-175.
(On M Permian ammonite Cyclolobus, incl. mentions of occurrences in Basleo and Ruasnain, W Timor)

Gerth, H. (1931)- Porifera. In:B.G. Escher et al. (eds.) Onze palaeontologische kennis van Nederlandsch Oost Indie, Leidsche Geologische Mededelingen 5 (Feestbundel K. Martin), p. 115-119.
(online at: www.repository.naturalis.nl/document/549435)
(Porifera (sponges) chapter in 'Our paleontological knowledge of the Netherlands East Indies'. Isolated sponge needles known from various formations in Indonesia, but whole sponges mainly known from Permian and Triassic of Timor. Permian of Timor 25 species, mainly siliceous sponges. Triassic of Timor mainly calcareous sponges. No figures)

Gobbett, D.J. (1967)- Palaeozoogeography of the Verbeekinidae (Permian foraminifera). In: C.G. Adams & D.V. Ager (eds.) Aspects of Tethyan biogeography: A Symposium, The Systematics Association, London, Special Publication 7, p. 77-91.
(Fusulinid formifera of Verbeekinidae family largely restricted to M Permian of Tethys seaway. In Malaya and Hapan (and ASumatra; JTvG) verbeekinid limestones are associated with volcanics. With M Permian distribution map from Mediterranean area, through the Asia Minor-C Asian Tethyan belt to SE Asia and Japan. In Indonesia limited to Neoschwagerina fauna in Sumatra and Timor)

Gould, R.E. (1975)- The succession of Australian Pre-Tertiary megafossil floras. The Botanical Review 41, 4, p. 453-483.
(Review of Devonian- Cretaceous floras of Australia, incl. Devonian Baragwanathia flora, Permian Glossopteris flora, Triassic Dicroidium flora, etc.)

Hahn, G. & C. Brauckmann (1975)- Revision zweier Trilobiten Arten aus dem Perm Asiens. Geologica et Palaeontologica 9, p. 117-124.
('Revision of two trilobite species from the Permian of Asia'. With distinction of new subgenus of Pseudophillipsia (Carniphillipsia))

Hayasaka, I. (1917)- On the brachiopod genus *Lyttonia* with several Japanese and Chinese examples. J. Geological Society Japan, Tokyo, 24, 288, p. 43-53.
(online at: https://www.jstage.jst.go.jp/article/geosoc1893/24/288/24_288_43/_pdf/-char/en)
(Discussion of Permian brachiopods of Lyttonia/ Leptodus type in Japan and China and Indochina (also known from Sumatra, Malay Peninsula, Timor, etc.; JTvG)

Heritsch, F. (1937)- Die rugosen Korallen und die Stratigraphie der Permformation. In: F.E. Suess Festschrift, Mitteilungen der Geologischen Gesellschaft in Wien, 29, p. 307-328.
(online at: https://www.zobodat.at/pdf/MittGeolGes_29_0307-0328.pdf)
('The rugose corals and the stratigraphy of the Permian'. Review of Permian coral zonation, including discussion of M Permian Basleo coral fauna of Timor, as first described by Gerth (1921))

Heritsch, F. (1937)- Rugose Korallen aus dem Salt Range, aus Timor und aus Djoulfa, mit Bemerkungen uber die Stratigraphie des Perms. Sitzungsberichte Akademie Wissenschaften, Wien, Math.- Naturwissenschaftliche Klasse, Abt. 1, 146, p. 1-16.
('Rugose corals from the Salt Range (Himalaya), from Timor and from Djoulfa, with remarks on the stratigraphy of the Permian'. Brief descriptions of some Permian rugose corals)

Hess, H. (1999)- Permian. In: H. Hess et al. (eds.) Fossil crinoids, Cambridge University Press, Chapter 20, p. 160-165.

(Timor Permian crinoid faunas most diverse and abundant in world, with 320 species described by Wanner, most new and unique to Timor. Permian crinoids from Australia cooler water faunas, with much lower diversity than Timor faunas)

Hill, D. (1957)- The sequence and distribution of Upper Palaeozoic coral faunas. Australian J. Science 19, p. 42-61.
(Review of Permian corals, including Timor material)

Igo, H. (1984)- Summary of the Palaeozoic conodonts from Malaysia and Thailand. In: T. Kobayashi, R. Toriyama & W. Hashimoto (eds.) Geology and Palaeontology of Southeast Asia, University of Tokyo Press, 25, p. 289-293.
(Brief review of Paleozoic conodonts in SE Asia. Oldest conodonts are of Ordovician age, from S Thailand)

Ishii, K.I. (1975)- On the genus *Colaniella* and its biostratigraphic significance. J. Geosciences, Osaka City University 19, 6, p. 107-138.
(online at: http://dlisv03.media.osaka-cu.ac.jp/infolib/user_contents/kiyo/DBe0190006.pdf)
Small latest Permian benthic foram genus Colaniella, generally associated with fusulinids (Palaeofusulina, Reichelina, Staffella) and small foraminifera (Pachyphloia, Globivalvulina, Agathammina, Endothyra, etc.). Common in Japan, S China, Himalayas, Mediterranean, also in Thailand, Malay Peninsula (Kelantan))

Ishii, K., Y. Okimura & K. Ichikawa (1985)- Notes on Tethys biogeography with reference to Middle Permian fusulinaceans. In: K. Nakazawa & J.M. Dickins (eds.) The Tethys: her paleogeography and paleobiogeography from Paleozoic to Mesozoic, Tokai University Press, Tokyo, p. 139-155.

Isozaki, Y. (2006)- Guadalupian (Middle Permian) giant bivalve Alatoconchidae from a mid-Panthalssan paleo-atoll complex in Kyushu, Japan: a unique community associated with Tethyan fusulines and corals. Proc. of the Japan Academy, B82, 1, p. 25-32.
(online at: http://ea.c.u-tokyo.ac.jp/earth/Members/Isozaki/06Alatoconchidae.pdf)
(Large Alatoconchidae bivalves in M Permian, with typical Tethyan fusulinids of Neoschwagerina and Lepidolina zones. Range up to end-Guadalupian extinction level)

Isozaki, Y. & D. Aljinovic (2009)- End-Guadalupian extinction of the Permian gigantic bivalve Alatoconchidae: end of gigantism in tropical seas by cooling. Palaeogeogr. Palaeoclim. Palaeoecology 284, p. 11-21.
(Large, thick Permian Alatoconchidae bivalves from E-M Permian shallow-marine carbonates in 9 areas in low-latitudes of Tethyan and Panthalassan domains, incl. Thailand, Malaysia, Philippines, and Japan. Always in association with large fusulines (Verbeekinidae) and/or rugose corals (Waagenophyllidae). This 'tropical trio' became extinct near Guadalupian-Lopingian boundary, probably due to temperature drop (Kamura cooling))

Kanmera, K., K. Ishii & R. Toriyama (1976)- The evolution and extinction patterns of Permian Fusulinaceans. In: T. Kobayashi & W. Hashimoto (eds.) Geology and Palaeontology of Southeast Asia, University of Tokyo Press, 17, p. 129-154.
(Review of parallel evolution of several lineages of fusulinid foraminifera, each with short periods of rapid diversification, interspersed with longer periods of relative stability. No sudden and catastrophic extinctions of fusulinaceans, but gradual waning of group through later Permian)

Kawamura, T. & H. Machiyama (1995)- A Late Permian coral reef complex, South Kitakami Terrane, Japan. Sedimentary Geology 99, p. 135-150.
(M Permian Iwaizaki Lst of S Kitakami Terrane in NE Japan, represents coral reef complex. With rugose and tabulate corals, solenoporacean algae, calcisponges, Tubiphytes, Archaeolithoporella, etc. High diversity unique among Permian reefs, most of which are mainly formed by calcisponges and calcareous algae. Coral reefs developed mainly around S China and Indochina continents, in tropics. Calcisponge reefs and Tubiphytes-algal crust reefs common in tropical- subtropical regions; stromatolite-bryozoan reefs in arid areas)

Kemp, E.M., B.E. Balme, R.J. Helby, R.A. Kyle, G. Playford & P.L. Price (1977)- Carboniferous and Permian palynostratigraphy in Australia and Antarctica: a review. BMR J. Australian Geology Geophysics 2, p. 177-208.
(online at: www.ga.gov.au/corporate_data/80927/Jou1977_v2_n3_p177.pdf)

Kobayashi, F. (2005)- Permian foraminifers from the Itsukaichi-Ome area, West Tokyo, Japan. Journal of Paleontology 79, 3, p. 413-432.
(Continuous carbonate deposition of >70 My on Panthalassan buildup in Carboniferous-Permian limestone blocks in Itsukaichi- Ome area. Blocks and breccias resulted from collapse of seamount and tectonic mixing with trench-fill deposits in Jurassic accretionary complexes of Chichibu Terrane. 105 species of Permian and 65 species of Carboniferous forms in 24 fusulinacean zones)

Kobayashi, F. (2011)- Permian fusuline faunas and biostratigraphy of the Akasaka Limestone (Japan). Revue de Paleobiologie, Geneve, 30, 2, p. 431-574.
(online at: www.ville-ge.ch/mhng/paleo/paleo-pdf/30-2/pal_30_2_01a.pdf)
(Mainly M Permian Akasaka Lst in C Japan with 63 species of fusulinids, divided into 7 zones: Parafusulina nakamigawai (U Kungurian), Cancellina nipponica (Lw Roadian), Neoschwagerina simplex (U Roadian), N. craticulifera (Lw Wordian), N. colaniae (U Wordian) Yabeina globosa (Capitanian) and Nanlingella suzukii Zone (Wuchiapingian))

Kobayashi, F. (2012)- Permian non-fusuline foraminifers of the Akasaka Limestone (Japan). Revue de Paleobiologie, Geneve, 31, 2, p. 313-335.
(online at: www.ville-ge.ch/mhng/paleo/paleo-pdf/31-2/pal_31_2_03.pdf)
(Description of diverse Permian smaller foram assemblages with Tethyan affinities, from 250m thick Akasaka Lst, deposited on Permian Panthalassan seamount and emplaced as exotic block in Jurassic-E Cretaceous accretionary complexes of Mino Terrane in C Japan. Associated with rich fusulinid assemblages, with 7 zones.)

Kobayashi, F. (2012)- Middle and Late Permian foraminifers from the Chichibu Belt, Takachiho Area, Kyushu, Japan: implications for faunal events. Journal of Paleontology 86, p. 669-687.
(M-L Permian- Triassic carbonates (<100m thick?) with fusulinids in S Chichibu Belt of C Kyushu, originated as Panthalassan seamount, now in ?Jurassic accretionary complex. Permian (Guadalupian-Lopingian) and Triassic ages)

Kobayashi, F., C.A. Ross & J.R.P. Ross (2010)- *Thailandina* and *Neothailandina*, and their subfamily Thailandininae: an example of an invalid taxonomic group of Permian fusuline foraminifera. Journal of Paleontology 84, 2, p. 360-361.
(Thailandina and Neothailandina genera of Permian fusulinids from Rat Buri Limestone in Thailand first proposed by Toriyama and Kanmera (1968), based on unique wall structure. However, thailandina-type wall is secondary mineralization of calcite as seen in specimens of Staffellidae, Misellina, etc.)

Kobayashi, T. & T. Hamada (1978)- Three suites of Carboniferous trilobites in Southeast Asia. Proc. of the Japan Academy, Series B, 54, 3, p. 92-95.
(online at: https://www.jstage.jst.go.jp/article/pjab1977/54/3/54_3_92/_pdf)
(18 species of trilobites known from Carboniferous of SE Asia. Oldest assemblage in Langgon Red Beds in NW Malay Peninsula (Langgonbole, Waribole, Macrobole, etc.). Younger assemblage (Dinantian) with Linguphillipsia (= Phillipsia of older authors?))

Krassilov, V.A. (2000)- Permian phytogeographic zonality and its implications for continental position and climates. Paleontological Journal (Moscow) 34, Supplement 1, p. 587-598.
(online at: http://paleobotany.ru/pdf/Krassilov%202000%20-%20Permian%20Phytogeographic%20Zonality%20and%20Its.pdf)
(Revised scheme of Permian plant geography. Contrary to prevalent opinion views Sumatra (Jambi) and West Irian Jaya floras as 'mixed Eurogondwana' floras, and mapped in 'subtropical humothermic belt' with 'Gondwanan' India- Australia floras)

Krijnen, W.F. (1931)- Palaeozoic and Mesozoic Gastropoda, Lamellibranchiata and Scaphopoda. In: B.G. Escher et al. (eds.) De palaeontologie en stratigraphie van Nederlandsch Oost-Indie, Leidsche Geologische Mededelingen 5 (K. Martin Memorial Volume), p. 164-205.
(online at: www.repository.naturalis.nl/document/549430)
(Extensive listings of Paleozoic- Mesozoic molluscs (gastropods, bivalves) known from Indonesia in 1931. No figures)

Laveine, J.P., S. Zhang & Y. Lemoigne (2003)- Additional documentation to the knowledge of the Late Palaeozoic floras of East and Southeast Asia: general conclusions and references. Revue de Paleobiologie, Geneve, 22, p. 831-849.
(Floras suggest that Indochina, E Malaysia, and S and N China were closely connected during Carboniferous)

Leven, E.Y. (1997)- Permian stratigraphy and fusulinida of Afghanistan with their paleotectonic implications. Geological Society of America (GSA), Special Paper 316, p. 1-134.
(Complete Permian section in Afghanistan, with rich fusulinid faunas: 58 genera, 282 species, incl. 41 new. Asselian- Sakmarian in N Afghanistan and N Pamir tropical shelf carbonates and fusulinid assemblages. Age-equivalent rocks of S Afghanistan- S Pamir all siliciclastic, lacking fusulinids, with cold-water macrofaunas typical of Peri-Gondwanan seas. In S Afghanistan first fusulinids in Sakmarian, low diversity, mainly species of Pseudofusulina. By late Yahtashian- Bolorian (late E Permian= ~late Artinskian-Kungurian; JTvG), due to milder climate and widespread transgression, distinction between S and N fusulinid assemblages starts to disappear. In C-S Afghanistan occurrences of Monodiexodina in Bolorian (= Kungurian; late E Permian), Neoschwagerina- Verbeekina- Sumatrina in Murgabian-Midian (M Permian) and miliolid Shanita amosi-Hemigordiopsis renz in Midian-Dzhulfian (M-L Permian))

Leven, E.Y. (2003)- The Permian stratigraphy and Fusulinids of the Tethys. Rivista Italiana Paleontologia Stratigrafia 109, 2, p. 267-280.
(online at: https://riviste.unimi.it/index.php/RIPS/article/view/5507/5553)

Leven, E.J. & H.J. Campbell (1998)- Middle Permian (Murgabian) fusuline faunas, Torlesse Terrane, New Zealand. New Zealand J. Geology Geophysics 41, p. 149-156.
(online at: www.tandfonline.com/doi/pdf/10.1080/00288306.1998.9514799)
(Two fusuline foram limestone localities in Torlesse Gp of Canterbury, S Island, New Zealand. Limestones associated with volcanics and hemipelagic sediments which appear 'allochthonous' (fortuitous accumulations associated with oceanic seamounts). Low diversity fauna dominated by Parafusulina (Skinnerella) japonica, also Parafusulina (S.) cuniculata. Fauna correlates best with late M Permian (E Murgabian) sequences of Tethyan affinity in Eurasia)

Leven, E.J. & J.A. Grant-Mackie (1997)- Permian fusulinid foraminifera from Wherowhero Point, Orua Bay, Northland, New Zealand. New Zealand J. Geology Geophysics 40, p. 473-486.
(online at: www.tandfonline.com/doi/pdf/10.1080/00288306.1997.9514777)
(Abundant, diverse E-M Midian fusulinids (24 species) from Waipapa Terrane, N Island. Probably East Paleotethys-Panthalassa seamount fauna. Key species Neoschwagerina margaritae, Yabeina spp, Reichelina sp, Lepidolina shiraiwensis, etc. Much more diverse than Murgabian faunas from Torlesse, S Island)

Long, J.A. (1990)- Late Devonian Chondrichthyans and other microvertebrate remains from Northern Thailand. J. Vertebrate Paleontology 10, 1, p. 59-71.
(Microvertebrates from limestones near Burmese border town of Mae Sam Lap, N Thailand, dated as Late Famennian by conodonts. Fauna contains chondrichthyans, several types of chondrichthyan scales, actinopterygian scales and teeth, and rare acanthodian scales. Taxa suggest close affinity between Shan-Thai, East Gondwana, and S China Terranes at end of Devonian)

Macurda, D.B. (1983)- Systematics of the fissiculate Blastoidea. University of Michigan Papers on Paleontology, Ann Arbor, 22, p. 1-291.

(Strictly taxonomic review of Permian blastoids, much of which based on Timor material. No location info, biostratigraphic succession, etc.)

Mamet, B.L. & E. Saurin (1970)- Sur la microfaune des foraminiferes carboniferes du Sud-est asiatique. Bull. Societe Geologique France (7) 12, 2, p. 356-363.
('On the Carboniferous foraminiferal microfauna of Southeast Asia'. Eight Carboniferous foram assemblages, similar to Europe/ Asia, described from Laos, Vietnam, Malaysia)

Martodjojo, Sujono (1964)- A study of the superfamily Adrianitaceae. Contributions Department of Geology, Institute of Technology Bandung (ITB) 58, p. 75-81.
(Two groups recognized in ?Permian ammonoid family, from Timor?)

McLoughlin, S. (1993)- Glossopterid megafossils in Permian Gondwanic non-marine biostratigraphy. In: R.H. Findlay et al. (eds.) Proc. Gondwana Eight- Assembly, evolution and dispersal, Balkema, Rotterdam, p. 253-264.
(Review of Permian Glossopteris floras, characteristic of Gondwana. NE Australian Strong provincialism: Bowen Basin Glossopteris present, but no species in common with Indian or South African successions))

McLoughlin, S. (1995)- New records of *Bergiopteris* and glossopterid fructifications from the Permian of Western Australia and Queensland. Alcheringa 19, 3, p. 175-192.

Metcalfe, I. & Y. Isozaki (2009)- Current perspectives on the Permian-Triassic boundary and end-Permian mass extinction: Preface. J. Asian Earth Sciences 36, p. 407-412.
(End-Permian mass extinction nowwell dated at 252.6 ± 0.2 Ma (U-Pb) and Permian-Triassic GSSP level is dated by interpolation at 252.5 Ma. Conodonts evolved rapidly in first 1 million years following mass extinction leading to high-resolution conodont zones. Nature of double-phased Late Permian extinction (at Guadalupian-Lopingian boundary and P-T boundary, linked to large igneous provinces, suggests superplume activity that involved geomagnetic polarity change and massive volcanism)

Morante, R. (1996)- Permian and early Triassic isotopic records of carbon and strontium in Australia and a scenario of events about the Permian-Triassic boundary. Historical Biology 11, p. 289-310.
(Organic carbon in wells in Australian basins showed negative d13C excursions of 6-10%, believed to correlate with negative d13C excursion in marine carbonates at Permian-Triassic boundary around world. Shift consistently near base of miospore Protohaploxypinus microcorpus Zone of Helby et al. (1987))

Nestell, M.K. & G.P. Pronina (1997)- The distribution and age of the genus *Hemigordiopsis*. In: C.A. Ross et al. (eds.) Late Paleozoic foraminifera; their biostratigraphy, evolution, and paleoecology; and the mid-Carboniferous boundary, Cushman Foundation Foraminiferal Research, Special Publ. 36, 3, p. 105-110.
(online at: https://www.academia.edu/62505006/The_Distribution_and_age_of_the_genus_Hemigordiopsis)
(M-L Permian (Late Midian) foraminifer Hemigordiopsis appears to be characteristic of 'Cimmerian' terranes (including Sibumasu) which rifted off N Gondwana margin in Permian, now as remnants in Tethyan belt from Mediterranean to Peninsular Thailand and W Malay Peninsula. Hemigordiopsis evolved into Shanita in Late Midian; both go extinct at end of Late Midian))

Newell Arber, E.A. (2005)- Catalogue of the fossil plants of the *Glossopteris* flora in the Department of Geology, British Museum (Natural History); being a monograph of the Permo-Carboniferous flora of India and the Southern Hemisphere. British Museum (Natural History). Dept. of Geology, London, p. 1-255.
(online at: www.books.google.com/...)
(Old monograph on Carboniferous-Permian plant fossils of Gondwanaland (mainly India))

Nicoll, R.S. & I. Metcalfe (2011)- The Permian conodont biostratigraphy of Australia and New Zealand. In. XVII Int. Congress Carboniferous and Permian, Perth 2011, Geological Survey West Australia, Record 2011/20, p. 96. *(Abstract only)*

(Summary of Permian conodont studies. High paleolatitude Permian conodont faunas of W Australia dominated by Vjalovognathus with occasional Hindeodus, Mesogondella and Sweetognathodus. Lower latitude faunas from Timor, Pakistan, Nepal/Tibet higher diversity, attributed to warmer temperatures)

Okimura, Y., K. Ishii & C.A. Ross (1985)- Biostratigraphical significance and faunal provinces of Tethyan Late Permian smaller foraminifera. In: K. Nakazawa & J.M. Dickins (eds.) The Tethys: her paleogeography and paleobiogeography from Paleozoic to Mesozoic, Tokai University Press, Tokyo, p. 115-138.

Ota, A. & Y. Isozaki (2006)- Fusuline biotic turnover across the Guadalupian-Lopingian (Middle-Upper Permian) boundary in mid-oceanic carbonate buildups: biostratigraphy of accreted limestone in Japan. J. Asian Earth Sciences 26, p. 353-368.
(Fusulinid biostratigraphy of upper M- lower U Permian shallow-water limestones at Kamura and Akasaka in SW Japan. Both sections represent seamount carbonate buildups developed on basaltic basement in mid-oceanic environment. Sections contain abundant Tethyan fusulines and record extinction of M Permian large-sized fusuline family Verbeekinidae at Guadalupian- Lopingian boundary in mid-Panthalassa Ocean, similar to shallow-water Tethyan shelf areas)

Ozawa, T. (1970)- Notes on the phylogeny and classification of the Superfamily Verbeekinoidea: Studies of the Permian Verbeekinoidean Foraminifera-I. Memoirs Faculty Science, Kyushu University, Ser. D, Geology, 20, 1, p. 17-58.
(online at: https://catalog.lib.kyushu-u.ac.jp/opac_download_md/1543907/20_01_p017.pdf)
(Review of evolution and paleogeography of Tethyan M-L Permian Verbeekina group fusulinid foraminifera)

Ozawa, T. (1975)- Evolution of *Lepidolina multiseptata* (Permian foraminifer) in East Asia. Memoirs Faculty of Science, Kyushu University, Ser. D, Geology, 23, 2, p. 117-264.
(online at: https://catalog.lib.kyushu-u.ac.jp/opac_download_md/1544173/23_02_p117.pdf)
(Review of evolution in M-U Permian verbeekinoid foraminifer Lepidolina multiseptata (Deprat), which occurs in SE Asia (Sumatra, Malay Peninsula (C Pahang), W Cambodia,), S China, Japan and New Zealand (in E Asia Lepidolina now viewed as typicalof the E Tethyan- Cimmerian Province (Sibumasu/ Lhasa Block, etc.;Huang et al. 2022) and late M Permian?))

Palmieri, V. (1994)- Permian foraminifera in the Bowen Basin, Queensland. Queensland Geology 6, p. 1-125.

Palmieri, V., C.B. Foster & E.V. Bondareva (1994)- First record of shared species of Late Permian small foraminiferids in Australia and Russia: time correlations and plate reconstructions. AGSO J. Australian Geology Geophysics 15, p. 359-365.
(online at: https://d28rz98at9flks.cloudfront.net/81403/Jou1994_v15_n3_p359.pdf)
(At least 12 species of small calcareous foraminiferids from early Late Permian assemblages from Arctic Russia also occur in assemblages from E Australian Ingelara Fm of Bowen Basin)

Pia, J. (1937)- Die wichtigsten Kalkalgen des Jungpalaeozoicums und ihre geologische Bedeutung. Comptes Rendus 2nd Congres Avancement Etudes de Stratigraphie du Carbonifere, Heerlen 1935, 2, p. 765-856.
('The most important calcareous algae from the Late Paleozoic and their stratigraphic significance'. Incl. description of M Permian algae assemblages with Vermiporella sumatrana n.sp. from Sumatra (Fontaine 1989))

Pickett, J.W. (2011)- Fossil corals of Australia, New Zealand, New Guinea and Antarctica: bibliography and index. Memoir Assoc. Australasian Palaeontologists (AAP) 40, p. 1-189.
(Bibliography and index of published fossil coral research from Antarctica, Australia, New Guinea and New Zealand, covering 1343 species names, 607 genus names and 639 references)

Posthumus, O. (1931)- Plantae. In: B.G. Escher et al. (eds.) Onze palaeontologische kennis van Nederlandsch Oost Indie, Leidsche Geologische Mededelingen 5 (Feestbundel K. Martin), p. 485-508.
(online at: www.repository.naturalis.nl/document/549382)

(Listings of Permian- Pleistocene fossil plant and calcareous algae species from Indonesia, as known in 1931: mainly from Permian of Sumatra and Tertiary of Java, Sumatra, Kalimantan)

Ross, J. (1978)- Biogeography of Permian ectoproct bryozoa. Palaeontology 21, 2, p. 341-356.
(online at: http://palaeontology.palass-pubs.org/pdf/Vol%2021/Pages%20341-356.pdf)
(Overview of Permian bryozoa distribution, incl. comments on Timor assemblages)

Segroves, K.L. (1969)- Saccate plant microfossils from the Permian of Western Australia. Grana Palynologica 9, p. 174-227.
(online at: www.tandfonline.com/doi/pdf/10.1080/00173136909436435)
(29 species of saccate spores from Permian of N Perth Basin)

Shen, S.Z. (2018)- Global Permian brachiopod biostratigraphy: an overview. In: S.G. Lucas & S.Z. Shen (eds.) The Permian timescale, Geological Society, London, Special Publ. 450, p. 289-320.
(Permian brachiopod successions in five major paleobiogeographical realms. For Gondwanaland and peri-Gondwanan regions including Cimmerian blocks, Bandoproductus, Cimmeriella characteristic of Cisuralian (E Permian). Lower Permian brachiopods from Mengkareng Fm in Sumatra (Crippa et al. 2014) viewed as Sakmarian in age and grouped with S Thailand- Malaysia Cimmerian/ Sibumasu faunas (despite association with 'Jambi Flora, which is generally viewed as 'Cathaysian'?; JTvG)))

Shen, S.Z. & G.R. Shi (2002)- Paleobiogeographical extinction patterns of Permian brachiopods in the Asian-western Pacific region. Paleobiology 28, p. 449-463.
(End-Permian extinction eliminated ~90% of genera and 95% of species of Brachiopoda. End-Guadalupian extinction less profound)

Shen, S.Z., H. Zhang, W.Z. Li, L. Mu & J.F. Xie (2006)- Brachiopod diversity patterns from Carboniferous to Triassic in South China. Geological Journal 41, p. 345-361.
(Carboniferous to Triassic includes (1) 100 My-long stable biodiversity stage from Late Carboniferous- late Middle Permian, with highly diverse brachiopod faunas; (2) end-Permian most severe mass extinction in Phanerozoic; (3) bleak stage in E Triassic and (4) rapid recovery stage in M Triassic)

Smith, T.E., T. Bernecker, S. Bodorkos, J. Gorter, L. Hall, T. Hill, E. Holmes, A. Kelman, K. Khider, J. Laurie et al. (2017)- The impact of recalibrating palynological zones to the chronometric timescale: revised stratigraphic relationships in Australian Permian and Triassic hydrocarbon-bearing basins. AAPG/SEG 2017 Int. Conf. Exhibition, London, Search and Discovery Article 51443, p. 1-9. *(Poster Presentation)*
(online at: http://www.searchanddiscovery.com/documents/2017/51443smith/ndx_smith.pdf)
(Recalibration of Permian and Triassic spore-pollen palynozones and numerical ages from high-precision radiometric dating of tuffs)

Smith, T.E. & D. Mantle (2013)- Late Permian palynozones and associated CA-IDTIMS dated tuffs from the Bowen Basin, Australia. Geoscience Australia Record 2013/46, p. 1-39.
(onine at: https://d28rz98at9flks.cloudfront.net/72990/Rec2013_046.pdf)
(Calibration of Late Permian palynozones with radiometric ages of associated tuffs. Dulhuntyispora parvithola zone APP5 spans >5 Myrs (~254.4- 263 Ma). Latest Permian Playfordiaspora crenulata and Protohaploxypinus microcorpus palynozones APP6 between ~252-254.4 Ma) (see also Laurie et al. 2016))

Stehli, F.G. (1961)- New genera of Upper Paleozoic Terebratuloids. Journal of Paleontology 35, 3, p. 457-466.
(New genus and species name Timorina broili for small terebratulid brachiopod, originally described as Notothyris minuta by Broili (1916) from M Permian of Basleo, Timor)

Stephenson, M.H. (2008)- A review of the palynostratigraphy of Gondwanan Late Carboniferous to Early Permian glacigene successions. In: C.R. Fielding et al. (eds.) Resolving the Late Paleozoic ice age in time and space, Geological Society of America (GSA), Special Paper 441, p. 317-330.

(E Permian W Australian, Arabian and S African sequences can be correlated using taxa like Converrucosisporites confluens and Pseudoreticulatispora pseudoreticulata. C. confluens and P. pseudoreticulata zones considered to be Sakmarian, and Striatopodocarpites fusus zone is Artinskianin age. Difficult to correlate Gondwana palynological assemblages precisely to Russian type areas because of scarcity of marine fauna in Gondwana and different paleolatitudes, so Carboniferous-Permian boundary cannot be precisely correlated in Gondwana by palynology)

Stephenson, M.H. (2017)- Permian palynostratigraphy: a global overview. In: S.G. Lucas & S.Z. Shen (eds.) The Permian timescale, Geological Society, London, Special Publ. 450, p. 321-347.
(online at: https://www.lyellcollection.org/doi/10.1144/SP450.2)

Sun, K. (1999)- Origin, evolution and extinction of Cathaysia flora. Chinese Science Bull. 44, 2, p. 100-108.
(Carboniferous- Permian Cathaysia flora mainly distributed in China, Korea, Japan, Laos, Thailand, Sumatra and Malaysia. Mixed Cathaysian and Gondwanan flora known in S Tibet, Kashmir, Turkey and New Guinea. Cathaysian flora developed from global uniform E Carboniferous Lepidodendropsis flora)

Taboada, A.C., A.J. Mory, G.R. Shi, D.W. Haig & M.K. Pinilla (2015)- An Early Permian brachiopod-gastropod fauna from the Calytrix Formation, Barbwire Terrace, Canning Basin, Western Australia. Alcheringa 39, 2, p. 207-223.
(Small brachiopod-gastropod fauna from core near base of Calytrix Fm in glacially-influenced Grant Group on Barbwire Terrace, Canning Basin. Age probably Sakmarian, but possibly as old as Asselian. With palynomorphs of Pseudoreticulatispora confluens Zone)

Teichert, C. & B.F. Glenister (1952)- Fossil nautiloid faunas from Australia. Journal of Paleontology 26, p. 730-752

Tien, Nguyen D. (1989)- *Sphairionia sikuoides* gen. et sp. nov., a Permian incertae sedis organism: *Sphairionia*. In: Comptes Rendus 11th Congress Carboniferous Stratigraphy and Geology, Beijing 1987, 3, p. 73-78.
(Description of 'small bubble', either lagenid foraminifer or charophyte algae from M Permian of W Thailand and Cambodia. Also already described in 1988 'Note on two 'Incertae Sedis' from the Permian of West Thailand', CCOP Technical Bulletin 20. (also reported from Thailand and other areas of Tethys; Pronina 1996))

Tong, J. & G.R. Shi (2000)- Evolution of Permian and Triassic foraminifera in South China. In: H. Yin, J.M. Dickins et al. (eds.) Permian-Triassic evolution of Tethys and Western Circum-Pacific, Developments in palaeontology and stratigraphy 18, Elsevier, Amsterdam, p. 291-307.
(Paper mainly on stratigraphic distribution of foram genera. No paleobiogeography, no fossil illustrations)

Tripathi, C. & G. Singh (1985)- Carboniferous flora of India and its contemporaneity in the world. Comptes Rendus 10me Congres International Stratigraphie et Geologie du Carbonifere, Madrid 1983, p. 295-306.

Tripathi, C. & G. Singh (1987)- Gondwana and associated rocks of the Himalaya and their significance. In: G.D. McKenzie (ed.) Gondwana Six: Stratigraphy, Sedimentology, and Paleontology, 6th Int Gondwana Symposium, Columbus, Ohio, 1985, American Geophysical Union (AGU), Geophysical Monograph 41, p. 195-205.
(Along N margin of India Plate (Permian rift margin) two Permian Gondwanan domains, both with widespread diamictite, Glossopteris flora and Eurydesma fauna: (1) (S) Lesser Himalaya belt with Permian Volcanics directly on Precambrian/ Cambrian basement and with Glossopteris flora; (2) (N) Tethyan Himalaya belt (Salt Range-Ladakh- Thakkola- Lingshi) with more complete Cambrian- Permian section, more marine influence (limestones) and mixed Gondwana-Cathaysia flora. In Kashmir Basin Artinskian? Mamal Fm above Panjal Volcanics with rich E Permian flora, incl. Gondwanan Glossopteris, endemic Gangamopteris kashmirensis, also common 'Cathaysian' elements like Taeniopteris, Rajahia, Lobatanularia, Gingkophyllum and Parasphenophyllum. Also endemic vertebrate fauna. In Karakorum Basin (= Lhasa or Qiantang Block?) Fenestella Shale, overlain by diamictite, Gangamopteris plant bed and fusulinid lst)

Ueno, K. (2022)- Carboniferous fusuline foraminifera: taxonomy, regional biostratigraphy, and palaeobiogeographic faunal development. In: S.G. Lucas et al. (eds.), The Carboniferous timescale, Geological Society, London, Special Publ. 512, p. 327- 496.
(online at: https://www.lyellcollection.org/doi/10.1144/SP512-2021-107)
(Major review of Carboniferous fusulinid foraminifera worldwide, incl. mainland SE Asia and Timor)

Vachard, D. (2018)- Permian smaller foraminifers: taxonomy, biostratigraphy and biogeography. In: S.G. Lucas & S.Z. Shen (eds.), The Permian Timescale, Geological Society, London, Special Publ. 450, p. 205-252.
(online at: www.academia.edu/40553701/Permian_smaller_foraminifers_taxonomy_biostratigraphy_and_biogeography)

Vachard, D. & J. Ferriere (1991)- Une association a *Yabeina* (foraminifere fusulinoide) dans le Midien (Permien superieur) de la region de Whangaroa (Baie d'Orua Nouvelle-Zelande). Revue de Micropaleontologie 34, p. 201-230.
('A Yabeina association (fusulinid foraminifera) in the Midian (U Permian) of the Whangaroa region (Orua Bay, New Zealand'. Microfauna/ microflora from exotic limestone blocks embedded in volcanic and volcano-sedimentary units in Whangaroa area, New Zealand. Algae dominated by Tubiphytes. Smaller foraminifera common Tethyan species. Fusulinids mainly Reichelina, Codonofusiella, Chusenella, also Yabeina parvula. Outcrop is terrane that belonged to Cathaysia and which had migrated 3000 km S across Paleopacific)

Vachard, D., L. Pille & J. Gaillot (2010)- Palaeozoic foraminifera: systematics, palaeoecology and responses to global changes. Revue de Micropaleontologie 53, p. 209-254.
(Review of biostratigraphy and facies models of Paleozoic forams)

Van Gorsel, J.T. (2014)- An introduction to Paleozoic faunas and floras of Indonesia. Berita Sedimentologi 31, p. 6-26.
(online at: https://journal.iagi.or.id/index.php/FOSI/article/view/122)
(Review of Ordovician-Permian fossils of Indonesia. E Paleozoic faunas mainly in W Papua. Late Paleozoic (mainly Permian) faunas and floras more widespread, mainly on Sumatra, Timor and NW Borneo. Paleozoic fossils from Indonesia are mainly marine organisms, but non-marine Permian plant fossils are also known from Jambi (Sumatra) and Bird Head (West Papua). Some assemblages or species signify 'low-latitude Tethyan' settings; others have 'anti-tropical/subtropical Tethyan' or 'Gondwanan' affinities)

Von Staff, H. (1909)- Beitrage zur Kenntnis der Fusuliniden. Neues Jahrbuch Mineralogie Geologie Palaontologie, Beilage Band 27, p. 461-508.
('Contribution to the knowledge of the fusulinids'. Permian larger foram Schwagerina verbeeki Geinitz from Padang Highlands, W Sumatra, should be classified in new genus Verbeekina (see also Thompson 1936; Genus name Verbeekina still used today, and is viewed as 'Tethyan' genus, also known from S China, Thailand, Tibet, Crimea, etc.; JTvG))

Wahlmann, G.P. (2002)- Upper Carboniferous- Lower Permian (Bashkirian- Kungurian) mounds and reefs. In: W. Kiessling et al. (eds.) Phanerozoic reef patterns, Soc. Sedimentary Geology (SEPM) Special Publ. 72, p. 271-338.
(Includes mention of Timor Permian (Sakmarian) Tubiphytes (= Shamovella) grainstones)

Waterhouse, J.B. (1973)- Permian brachiopod correlations for South-East Asia. In: B.K. Tan (ed.) Proc. Regional Conference on the Geology of SE Asia, Kuala Lumpur 1972, Bull. Geological Society Malaysia 6, p. 187-210.
(online at: https://gsm.org.my/wp-content/uploads/gsm_file_2/702001-101346-PDF.pdf)
(Correlations of Permian brachiopod assemblages in Indonesia (Timor), Malaysia, Thailand, Vietnam, Burma, etc., using fusulinids and ammonoids as main age framework)

Weidlich, O. (2002)- Middle and Late Permian reefs- distributional patterns and reservoir potential. In: W. Kiessling et al. (eds.) Phanerozoic reef patterns, Soc. Sedimentary Geology (SEPM) Special Publ. 72, p. 339-390.
(Includes SE Asia info: prolific Permian rugose coral faunas found in mainland SE Asia, Sumatra and Timor)

Xu, G. & R.E. Grant (1994)- Brachiopods near the Permian-Triassic Boundary in South China. Smithsonian Contr. Paleobiology 76, p. 1-68.
(online at: www.sil.si.edu/smithsoniancontributions/paleobiology/pdf_hi/SCtP-0076.pdf)
(Latest Permian (Changxingian) in S China with Cathaysian Tethyan brachiopod assemblage of 164 species, incl. Leptodus spp. After major extinction event 20 species in E Triassic (lower Griesbachian))

Yabe, H. & S. Hanzawa (1931)- Palaeozoic and Mesozoic foraminifera. In: B.G. Escher et al. (eds.) De palaeontologie en stratigraphie van Nederlandsch Oost-Indie (Feestbundel K. Martin), Leidsche Geologische Mededelingen 5, p. 23-34.
(online at: https://repository.naturalis.nl/pub/505689/LGM1931005001003.pdf)
(Listings of Paleozoic- Mesozoic foraminifera species reported from Indonesia: Carboniferous- Permian (Sumatra, Timor, Leti, Luang), Triassic- Jurassic (Sumatra) and Cretaceous (Sumatra, Java, Borneo, Timor, Roti, Ceram, etc.). Mid-Cretaceous Orbitolina known from Java, Sumatra and Borneo: typical Tethyan fauna, adapted to warm-water, carbonate bank environments, Borneo Orbitolina not O.concava, but O. scutum. probably U Aptian age)

Yancey, T.E. & D.W. Boyd (1983)- Revision of the Alatoconchidae: a remarkable family of Permian bivalves. Palaeontology 26, p. 497-520.
(online at: http://cdn.palass.org/publications/palaeontology/volume_26/pdf/vol26_part3_pp497-520.pdf)
(Review of M Permian large, thick-walled bivalves, including material from Kinta Valley, Malay Peninsula (H.S. Lee mine, etc.). Oldest known alatoconchids Shikamaia (Tanchintongia) perakensis and Saikraconcha (Dereconcha) kaparensis, of probable E Artinskian age)

Zaninetti, L., J. Whittaker & D. Altiner (1979)- The occurrence of *Shanita amosi* Bronnimann, Whittaker and Zaninetti (Foraminifera) in the Late Permian of the Tethyan region. Notes du Laboratoire Paleontologie Universite de Geneve 5, 1, p. 1-7.
(Late Permian small miliolid foraminifer Shanita appears to be marker for M-L Permian of N Gondwana margin; see also Jin & Yang, 2005)

Zhou, W., M. Wan, R.A. Koll & J. Wang (2018)- Occurrence of the earliest gigantopterid from the basal Permian of the North China Block and its bearing on evolution. Geological Journal 53, 2, p. 500-509.
(Gigantopterid plants characteristic floral element in Permian Cathaysian floras. However, in China oldest known occurrences later than in N America (Artinskian) and Sumatra/Indonesia (Asselian-Sakmarian). New gigantopterid Gigantonoclea cf. mira from basal Permian (Asselian) strata in N China Block represents oldest unequivocal evidence for gigantopterids)

X.6. Quaternary Hominids and Mammals, Archeology

N.B. See also references to related Quaternary mammals-hominids assemblages from Thailand, Myanmar, South China and Malay Peninsula .

Abdurahman, O., U.P. Wibowo, A. Zaennudin, M. Thamrin & J. Arif (2020)- M. Thamrin- Sang pemburu fosil dari Cirebon (dilengkapi dengan geologi dan koleksi fosil). Badan Geologi, Bandung, p. 1-119.
(online at: https://geologi.esdm.go.id/publikasi/laporan-dan-buku/m-thamrin-sang-pemburu-fosil-dari-cirebon-dilengkapi-dengan-geologi-dan-koleksi-fosil)
('M. Thamrin- The fossil hunter from Cirebon (with geology and fossil collections)'. Book about Javanese amateur fossil collector M. Thamrin and his Pleistocene mammal collections from 40 years of fossil hunting in the Cirebon area, W Java)

Adhityatama, S., R.R. Triwurjani, D. Yurnaldi, R. Janssen, M.D.K. Dhony, Suryatman et al. (2022)- Pulau Ampat site: a submerged 8th century iron production village in Matano Lake, South Sulawesi, Indonesia. Archaeological Research in Asia 19, 100335, p. 1-11.
(Population of Indonesia grew dramatically after discovery of iron sources and use of iron tools. Pamor Luwu iron used by Java's 13th-15th century Majapahit Empire, primary sourced from Matano Lake area in SE Sulawesi. Evidence for underwater site named Pulau Ampat)

Aiello, L.C. (2010)- Five years of *Homo floresiensis*. American J. Physical Anthropology142, 2, p. 167-179.
(online at: https://www.academia.edu/1353347/Five_years_of_Homo_floresiensis)
(Review of early debates on the nature of Homo floresiensis. Evidence supports the hypothesis that Homo floresiensis is late-surviving species of early Homo with its closest morphological affinities to early African pre-erectus/ergaster hominins. Controversy will probably continue)

Aimi, M. (1989)- A mandible of *Sus stremmi* Koenigswald 1933 from Cisaat, Central Java, Indonesia. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 6, p. 4-10.
(New mandible (jaw) of pig species from coarse sands of Kali Glagah Fm at Cisaat, Bumiayu region, C Java)

Aimi, M. & F. Aziz (1985)- Vertebrate fossils from the Sangiran dome, Mojokerto, Trinil and Sambungmacan, Indonesia. In: N. Watanabe & D. Kadar (eds.) Quaternary geology of the hominid bearing formations in Java. Geological Research Development Centre (GRDC), Bandung, Special Publ. 4, p. 155-198.

Aimi, M. & Sudijono (1979)- On the problematical species *Aceratherium boschi* von Koenigswald 1933. Bull. Geological Research Development Centre (GRDC) 1, p. 37-45.
(online at: http://www.rhinoresourcecenter.com/pdf_files/149/1498297867.pdf)
(On latest Miocene?-Pliocene? rhinoceros molar from calcarenite in upper Halang Beds in Cisande River, W Java. First described by Von Koenigswald in 1933. Here believed to be incomplete molar referable to Rhinoceros sondaicus. Foram age control deemed non-diagnostic)

Alink, G., S. Adithayama & T. Simanjuntak (2017)- Descriptive analysis of Palaeolithic stone tools from Sulawesi, collected by the Indonesian-Dutch expedition in 1970. AMERTA (J. Penelitian dan Pengembangan Arkeologi) 35, 2, p. 75-148. *(online at: www.researchgate.net/publication/323656343_DESCRIPTIVE_ANALYSIS_OF_PALAEOLITHIC_STONE_TOOLS_FROM_SULAWESI_COLLECTED_BY_THE_INDONESIAN-DUTCH_EXPEDITION_IN_1970)*
(1970 Indonesian-Dutch expedition, headed by Soejono, Van Heekeren and Hooyer collected 1100 lithic objects from sites Beru and Marale in Walanae Depression in S Sulawesi. Vertebrate fossils (mainly pigs, turtle) from Walanae valley from upper Late Pliocene and possible Lower Pleistocene Walanae Fm. 'Cabenge Industry' lithic artefacts in terrace deposits, may be of Late Pleistocene age)

Alink, G., W. Roebroeks & T. Simanjuntak (2016)- The *Homo erectus* site of Trinil: past, present and future of a historic place. AMERTA (J. Penelitian dan Pengembangan Arkeologi) 34, 2, p. 99-114.
(online at: https://jurnalarkeologi.kemdikbud.go.id/index.php/amerta/article/view/150)

(Review of excavation at the well-known Trinil Pithecanthropus erectus site along Solo River in Central Java, in 1891 (Dubois) and 1907 (Selenka Expedition))

Allen, H. (1991)- Stegodonts and the dating of stone tool assemblages in island S.E. Asia. Asian Perspectives 30, p. 243-266.
(online at: https://core.ac.uk/download/pdf/5105269.pdf)
(Stone tools found associated with U Pleistocene-Holocene Homo fossils on Java (Ngandong, Wajak, etc.), but no tools with any certainty associated with M Pleistocene Homo erectus (younger literature does suggest stone artifacts associated with H. erectus in Sangiran, etc.; JTvG))

Amano, N., T. Ingicco, A.M. Moigne, A.M. Semah, T. Simanjuntak & F. Semah (2022)- Monkey hunting in Early to Mid-Holocene Eastern Java (Indonesia). In: World Archaeoprimatology, Interconnections of humans and nonhuman primates in the past, Cambridge University Press, p. 474-496.
(online at: https://www.researchgate.net/publication/362430832_Monkey_Hunting_in_Early_to_Mid-Holocene_Eastern_Java_Indonesia)
(Excavations of E-M Holocene in Braholo Cave, Wonosari Plateau, western Gunung Sewu, East Java show high % of Javan langur (Trachypithecus auratus; ebony leaf monkey) bones, some with butchery marks, suggesting targeted hunting for food and as materials for tools and ornaments (see also Ingicco et al., 2020))

Amano, N., A.M. Moigne, T. Ingicco, F. Semah, R. DueAwe & T. Simanjuntak (2016)- Subsistence strategies and environment in Late Pleistocene- Early Holocene Eastern Java: evidence from Braholo Cave. Quaternary International 416, p. 46-63.
(Climatic shifts during Pleistocene- Holocene transition in Island SE Asia resulted in changes in landscapes, impacting vertebrate community composition and human subsistence economies. Braholo Cave in E Java with mainly arboreal fauna in Late Pleistocene - E-M Holocene, but older cave deposits dominated by animal taxa associated with open environments (bovids, cervids). Reflects forest expansion at onset of Holocene)

Amano, N., F. Rivals, A.M. Moigne, T. Ingicco, F. Semah & T. Simanjuntak (2016)- Paleoenvironment in East Java during the last 25,000 years as inferred from bovid and cervid dental wear analyses. J. Archaeological Science, Reports 10, p. 155-165.
(Dental wear analyses of bovid and cervid specimens from Late Pleistocene- M Holocene deposits of Braholo Cave and Song Terus in E Java. Pleistocene-Holocene transition accompanied by expansion of rainforests as evidenced by predominance of species that had browsing diet, but grazing signatures indicative of continued presence of patches of grasslands up to M Holocene. Dental wear from layers dating to Last Glacial Maximum (LGM) suggest a mosaic environment dominated by grasslands)

Ambrose, S.H. (1998)- Late Pleistocene human bottlenecks, volcanic winter, and differentiation of modern humans. J. Human Evolution 34, p. 623-651.
(Toba eruption on Sumatra may have caused human population bottleneck and modern human races may have differentiated abruptly only 70,000 years ago (see also commentary by Gathorne-Hardy 2003))

Ambrose, S.H. (2003)- Did the super-eruption of Toba cause a human population bottleneck? Reply to Gathorne-Hardy and Harcourt-Smith. J. Human Evolution 45, p. 231-237.
(~73 ka Toba eruption larger than previously estimated, and caused millennium of coldest temperatures of U Pleistocene. Genetic studies suggest real population bottleneck during first half of last glacial period, but no mass extinctions. We are descendants of few small groups of tropical Africans who united in face of adversity)

Anderson, C. (1937)- Palaeontological notes No. IV, Fossil marsupials from New Guinea. Records of the Australian Museum, Sydney, 20, 2, p. 73-76.
(online at: https://journals.australian.museum/media/Uploads/Journals/17261/250_complete.pdf)
(First record of Quaternary fossil marsupial from New Guinea (PNG), along the lower Bulolo and Watut Rivers, described as Nototherium watutense)

Ansyori, M.M. (2010)- Fauna from the oldest occupation layer in Song Terus cave, Eastern Java, Indonesia-biochronological significance of the Terus layer. Masters Thesis, Museum National Histoire Naturelle, Paris, p. 1-71.
(online at: www.ipt.pt/teses.digitais/ansyori.mirza/ansyori_mirza.pdf)
(Oldest archeological assemblage in Song Terus cave in Weru, Pacitan, S Mountains of S Java ranges in age from ~300-80 ka (M-L Pleistocene). Late Pleistocene faunas of upper Terus Layer (80-120 ka) resemble Punung fauna of Badoux. Big fauna dominated by Cervidae, Bovidae and Suidae. Tropical forest environment)

Ansyori, M. (2018)- Biostratigraphic significance of Sangiran Dome, Central Java, Indonesia: an insight on animal succession of Java during the Lower and Middle Pleistocene. Doctoral Thesis, International Doctorate in Quaternary and prehistory, Paris-Tarragone, p. 1-167.

Antoine, P.O. (2012)- Pleistocene and Holocene rhinocerotids (Mammalia, Perissodactyla) from the Indochinese Peninsula. Comptes Rendus Palevol 11, 2-3, p. 159-168.
(online at: https://www.sciencedirect.com/science/article/pii/S1631068311000492)
(Only living rhinoceros in Indochina Peninsula is Rhinoceros sondaicus (Vietnam). Extinct Pleistocene-Holocene Indochinese rhinocerotids include Dicerorhinus gwebinensis (E Pleistocene, Myanmar) and Recent Asian species Rhinoceros unicornis (M-L Pleistocene), R. sondaicus (M Pleistocene-Recent) and Dicerorhinus sumatrensis (M Pleistocene-Holocene). With SE Asia distribution maps)

Antoine, P.O., M.C. Reyes, N. Amano, A.P. Bautista, C.H. Chang, J. Claude, J. de Vos & T. Ingicco (2012)- A new rhinoceros clade from the Pleistocene of Asia sheds light on mammal dispersals to the Philippines. Zoological J. Linnean Society, London, 194, 2, p. 416-430.
(online at: https://academic.oup.com/zoolinnean/article/194/2/416/6280384)
(New Pleistocene rhinoceros species from 709 ± 68 kya archaeological deposits in Luzon Island, Philippines. Dental features reveal existence of unsuspected Asian Pleistocene clade, Nesorhinus gen. nov. Phylogenetic results suggest island-hopping dispersal for Nesorhinus from Asian mainland to Luzon via Taiwan by Late Miocene or later)

Anton, S.C. (1997)- Developmental age and taxonomic affinity of the Mojokerto child, Java, Indonesia. American J. Physical Anthropology 102, 4, p. 497-514.
(Mojokerto child (Perning I), Java, discovered in 1936, has been assigned to Australopithecus and multiple species of Homo modjokertensis, etc. Developmental age range probably 4-6 years)

Anton, S.C. (1999)- Cranial growth in *Homo erectus*: how credible are the Ngandong juveniles? American J. Physical Anthropology 108, 2, p. 223-236.
(Ngandong 5 and 9 skulls are adults, 8 an older juvenile and 2 is a juvenile. Adult cranial contours and pattern of contour development similar between Ngandong adults and other H. erectus adults. Nothing to suggest that Ngandong transitional in vault shape between H. erectus and H. sapiens, despite relatively large brain)

Anton, S.C. (2001)- Cranial evolution in Asian *Homo erectus*. Geological Research Development Centre (GRDC), Bandung, Special Publ. 27, p. 39-46.

Anton, S.C. (2002)- Evolutionary significance of cranial variation in Asian *Homo erectus*. J. Physical Anthropology 118, 4, p. 301-323.
(Principal components analysis of calvarial shape suggest regional differentiation between N Asian and SE Asian H. erectus. Most recent SE Asian fossils (e.g. Ngandong) conform to SE Asian pattern)

Anton, S.C. (2003)- Natural history of *Homo erectus*. Yearbook Physical Anthropology, Wiley, 46, p. 126-170.
(online at: http://studylib.net/download/8851017)
(In Indonesia earliest hominid crania (Sangiran) few in number, often badly deformed and ~1.6- 1.8 Ma old. More common hominids between 0.9- 1.5 Ma, with cranial capacity of ~800-1000cc. Later Indonesian hominids (Ngandong, Sambungmacan) <100ka in age, and lack upper facial, mandibular or dental remains. Cranial size 900- >1200cc, average >1000cc. Main differences between earlier and later Indonesian H. erectus: brain size

increase, increases in vault height, and decreases in postorbital constriction. H. erectus, notable for increased body size, originated around Plio-Pleistocene boundary in Africa and quickly dispersed into W and E Asia.)

Anton, S.C. & E.R. Middleton (2023)- Making meaning from fragmentary fossils: Early *Homo* in the Early to early Middle Pleistocene. J. Human Evolution 179, 103307, p. 1-56.
(Extensive re-evaluation of worldwide fossil record for early Homo (principally Homo erectus, Homo habilis, and Homo rudolfensis), from early diversification and dispersal in E Pleistocene (~2.0 Ma) to ultimate demise of H. erectus in early M Pleistocene. Includes occurrences in Indonesia.)

Anton, S.C., F. Spoor, C.D. Fellmann & C.C. Swisher (2007)- Defining *Homo erectus*: size considered. In: W. Henke et al. (eds.) Handbook of Paleoanthropology, Springer Verlag, 3, 11, p. 1655-1693.

Anton, S.C. & C.C. Swisher (2003)- Early dispersals of *Homo* from Africa. Annual Review of Anthropology 33, p. 271-296.
(Hominin presence outside Africa started at ~1.6-1.8 Ma (Homo erectus). Includes discussion on reliability of ages of Sangiran hominid ('poo-poo everything'; JTvG). Earliest hominins at Sangiran older than 1.0 Ma and probably 1.3 Ma or older)

Aplin, K. & K.M. Helgen (2010)- Quaternary murid rodents of Timor Part I: New material of *Coryphomys buehleri* Schaub, 1937, and description of a second species of the genus. Bull. American Museum Natural History 341, p. 1-80.
(online at: http://digitallibrary.amnh.org/handle/2246/6077)
(Archeological excavations in E Timor in 1968- 2002 provided new material of Late Pleistocene and recently extinct gigantic murine (rat) Coryphomys (originally described by Hooijer 1965))

Arifin, K. (2017)- Terminal Pleistocene and Early Holocene human occupation in the rainforests of East Kalimantan. In: P.J. Piper et al. (eds.) New perspectives in Southeast Asian and Pacific Prehistory, Chapter 6, p. 97-124.
(online at: https://press-files.anu.edu.au/downloads/press/n2320/pdf/ch06.pdf)
(Excavations at two rock shelters and cave in Berau region, NE Kalimantan, produced evidence for occupation of tropical rainforest along U Birang River by human foragers from at least end of Last Glacial Period)

Argue, D., C.P. Groves, M.S.Y. Lee & W.L. Jungers (2017)- The affinities of *Homo floresiensis* based on phylogenetic analyses of cranial, dental, and postcranial characters. J. Human Evolution 107, p. 107-133.
(online at: www.sciencedirect.com/science/article/pii/S0047248417300866)
(Analyses of multiple morphological characteristics suggest H. floresiensis (~65-90ka) is closest to early hominins (>1.75 Ma; Homo habilis?), suggesting probably long-surviving relict of early hominin lineage, with hitherto unknown migration out of Africa. Not recent descendants of either H. erectus or H. sapiens)

Ariens-Kappers, C.U. & K.H. Bouman (1939)- Comparison of the endocranial casts of the *Pithecanthropus erectus* skull found by Dubois and von Koenigswald's *Pithecanthropus* skull. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 45, p. 30-40.

Arif, J. (1998)- Morphological study of a *Homo erectus* skull (Skull IX) from Sangiran, Central Jawa, Indonesia. Thesis Kyoto University. p. 1-60.

Arif, J., H. Baba, M.E. Suparka, Y. Zaim & T. Setoguchi (2001)- Preliminary study of *Homo erectus* skull IX (Tjg-1993.05) from Sangiran, Central Java, Indonesia. Bull. National Science Museum, Tokyo, D, 27, p. 1-17.
(online at: https://www.kahaku.go.jp/research/publication/anthropology/download/27/BNSM_D2701.pdf)
(Similar to Arif et al. (2002) paper below, and Kaifu et al. (2011))

Arif, J., Y. Kaifu, H. Baba, M.E. Suparka, Y. Zaim & T. Setoguchi (2002)- Preliminary observation of a new cranium of *Homo erectus* (Tjg-1993.05) from Sangiran, Central Jawa. Anthropological Science 110, 2, p. 165-177.

(online at: https://www.jstage.jst.go.jp/article/ase1993/110/2/110_2_165/_pdf/-char/en)
(New well-preserved and most complete hominid skull IX found in 1993 from Bapang (Kabuh) Fm at Tanjung village, Sangiran region, C Java. Initially believed to be from Upper Tuff of Bapang Fm (Sartono & Tyler, 1993), then from Middle Sangiran Fm (Pucangan) Fm (Tyler & Sartono, 2001), but here relocated to basal or middle part of Bapang (Kabuh) Fm)

Arif, J., M.R. Schurr & R. Kapid (2008)- Evolutionary molar size reduction in Sangiran (Javanese), Early-Middle Pleistocene *Homo erectus* as an effect of palaeo-climatic change. J. Indian Ocean Archaeology 5, p. 10-23.
(online at: https://www.researchgate.net/profile/Johan-Arif/publication/370493081_Evolutionary_Molar_Size_Reduction_in_Sangiran_Javanese_Early-Middle_Pleistocene_Homo_erectus_As_an_Effect Etc.)(Secular molar size reduction in H.erectus's population from Sangiran from Grenzbank to Bapang Fm assemblages possibly associated with paleoclimate change between 1.0- 0.7 Ma)

Aubert, M., A. Brumm, M. Ramli, T. Sutikna, E.W. Saptomo, B. Hakim, M. J. Morwood, G. D. van den Bergh et al. (2014)- Pleistocene cave art from Sulawesi, Indonesia. Nature 514 , p. 223-227. *(ms online at: https://www.researchgate.net/publication/266683667_Pleistocene_cave_art_from_Sulawesi_Indonesia)*
(U-series dating of coralloid speleothems associated with 12 human hand stencils and two figurative animal depictions from cave sites in Maros karsts of S Sulawesi suggest earliest dated image of 39.9 ka, is oldest known hand stencil in world. A painting of babirusa ('pig-deer') dated as >35.4 ka is among earliest dated figurative depictions worldwide. Humans produced rock art by ~40 ka at opposite ends of Pleistocene Eurasia (see also Oktaviana et al., 2026))

Aubert, M., A. Brumm & P.S.C. Tacon (2017)- The timing and nature of human colonization of Southeast Asia in the late Pleistocene. A rock art perspective. Current Anthropology 58, Supplement 17, p. S553-S566.
(online at: https://www.journals.uchicago.edu/doi/epdf/10.1086/694414)
(Dating of rock art in S Sulawesi caves suggests early hunter-gatherers in SE Asia date back to at least ~40 ka)

Audley-Charles, M.G. & D.A. Hooijer (1973)- Relation of Pleistocene migrations of pygmy stegodonts to island arc tectonics in Eastern Indonesia. Nature 241, p. 197-198.
(Pleistocene pygmy stegodonts in Sulawesi, Flores and Timor, areas now separated by deep seas. Dwarf Stegodon populations coexisted in Flores and Timor, and apparently wandered back and forth in Pleistocene)

Aziz, F. (1983)- Notes on a new *Meganthropus* S.33 from the Sangiran Dome, Central Java. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 4, p. 56-60.

Aziz, F. (1989)- *Macaca fascicularis* (Raffles) from Ngandong, East Java. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 5, p. 50-56.
(Macaque fossil tooth from Pleistocene of Ngandong along Solo River, site of Homo soloensis. Supports earlier interpretations of open-country habitat)

Aziz, F. (1990)- Pleistocene mammal faunas of Sulawesi and their bearings to paleozoogeography. Ph.D. Thesis, Kyoto University, Japan, p. 1-106.

Aziz, F. (2000)- A new insight on the Pleistocene fauna of Sangiran and other hominid sites in Java. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 10, p. 49-57.
(Discussion of the De Vos 1982 and Sondaar 1982 Java mammalian biostratigraphy scheme)

Aziz, F. (2000)- The Pleistocene endemic fauna of the Indonesian Archipelago. Tropics 10, 1, p. 135-143.
(online at: www.jstage.jst.go.jp/article/tropics/10/1/10_1_135/_pdf/-char/ja)
(Brief review of Asiatic fauna migrations to Java since 1.5 Ma. Some crossed Wallace Line by sweepstake dispersal. They adapted to island environments and developed into unbalanced endemic island faunas)

Aziz, F. (2001)- New insight on the Pleistocene fauna of Sangiran and other hominid sites in Java. In: T. Simanjuntak et al. (eds.) Sangiran: man, culture, and environment in Pleistocene times, Proc. International Colloquium on Sangiran, Solo- Indonesia, Solo 1998, Jakarta, National Research Centre of Archaeology, Yayasan Obor Indonesia, Jakarta, p. 260-271.
(Same paper as Aziz (2000))

Aziz, F. (2001)- Hominid fossils housed at the Geological Research and Development Center, Bandung, Indonesia. In: A scientific life: papers in honor of Prof. Dr. T. Jacob, Bigraf Publishing, Yogyakarta, p. 53-66.

Aziz, F. (2002)- New discovery of a hominid skull from Cemeg, Sambungmacan, Central Java: an announcement. Jurnal Sumber Daya Geologi (JSDG) 7, 125, p. 2-7.
(Displaced hominid skull from Solo River sediments near Cemeng, W of Trinil, C Java, here called Sambungmacan 4. Probably Homo erectus of Pleistocene age)

Aziz, F. (2013)- Peran Badan Geologi dalam penelitian manusia purba. Geomagz 3, 2, p. 23-25.
('The role of the Geological Agency in ancient human research')

Aziz, F. & H. Baba (eds.) (2013)- *Homo erectus* in Indonesian. Recent progress of the study and current understanding, Centre for Geological Survey, Bandung, p. 1-106.
(Collection of papers)

Aziz, F., H. Baba & S. Narasaki (1994)- Preliminary report on recent discoveries of fossil hominids from the Sangiran area, Java. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 4, 29, p. 11-16.
(New hominid skull and mandible fragments, found by local collectors in E part of Sangiran Dome, C Java)

Aziz, F., H. Baba & N. Watanabe (1996)- Morphological study on the *Homo erectus* Sangiran 17 skull based on the new reconstruction. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 8, p. 11-25.

Aziz, F. & J. de Vos (1989)- Rediscovery of the Wadjak Site (Java, Indonesia). J. Anthropological Society of Nippon 97, 1, p. 133-144.
(online at: www.jstage.jst.go.jp/article/ase1911/97/1/97_1_133/_pdf)
(Site of Wajak cave at Gunung Lawa near Tjermee/ Campur Darat villages in C Java, with 'Wajak man' hominid remains and originally excavated by Dubois in 1890, still exists (latest Pleistocene or Holocene; see Storm et al., 2013)

Aziz, F. & J. de Vos (1999)- The fossil faunas from the Citarum area, West Java, Indonesia. In: J.W.F. Reumer and J. de Vos (eds.) Elephants have a snorkel!, Papers in honor of Paul Y. Sondaar, DEINSEA 7, p. 21-32.
(online at: https://natuurtijdschriften.nl/pub/538618/DEIN1999007001003.pdf)
(Four fossil mammal faunas from Citarum area, Bandung Basin, W Java: (1) Banuraja, with Sus, Panthera, Manis palaeojavanicus, etc., with age of ~0.8 Ma; (2) Cipatik/Cililin lake deposits with fossil fish of possibly Pleistocene age. (3) Ciharuman with fossils dated as ~29.6-42.3 k; (4) Cipeundeuy with Elephas maximus, Rhinoceros sondaicus, Rusa, Elephas maximus with age of ~0.42- 0.36 Ma)

Aziz, F., A.S. Hadiputro & H. Insani (2023)- Atlas fosil vertebrata koleksi Museum Geologi. Badan Geologi, Bandung, p. 1-96. *(online at: https://geologi.esdm.go.id/publikasi/laporan-dan-buku/atlas-fosil-vertebrata-koleksi-museum-geologi)*
('Atlas of vertebrate fossils in the collections of the Geological Museum'. In Indonesian and English. On fossil localities on Java, SW Sulawesi, Flores, etc. and examples of fossil materials in Bandung collected since early 1900s. With overview of Quaternary mammals in Indonesia, biostratigraphy, and descriptions of fossils from Java, Sulawesi and Flores in Geological Museum, Centre for Geological Survey, Bandung)

Aziz. F. & I. Kurniawan (2014)- Atlas of Indonesia *Homo erectus*. LIPI Press, Jakarta, p. 1-98.

Aziz, F., I. Kurniawan & E. Setiyabudin (2013)- The *Homo erectus* collection of the Centre for Geological Survey. In: F. Aziz & H. Baba (eds.) *Homo erectus* in Indonesian- Recent progress of the study and current understanding, Center for Geological Survey, Special Publ., Chapter 2, p. 9-42.
(Well illustrated listing of sizable collection of Homo erectus fossils from C Java (Sangiran, Sambungmacan, Mojokerto, Ngandong), collected since 1931)

Aziz, F. & I. Saefudin (1996)- An isolated tooth of orang-utan (*Pongo pygmaeus*) from the Sangiran area, Central Java, Indonesia. Publ. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 8, p. 47-50.

Aziz, F., P.Y. Sondaar, J. de Vos, G.D. van den Bergh & Sudijono (1995)- Early dispersal of man on islands of the Indonesian Archipelago: facts and controls. Anthropological Science 103, 4, p. 349-368.
(online at: https://www.jstage.jst.go.jp/article/ase1993/103/4/103_4_349/_pdf)
(Migration of hominids across sea barriers much earlier than Neolithic. Distance of island to mainland and food supply limiting factors on human colonization of islands. On islands of SE Asia giant rats may have served as food supply for Paleolithic Man. M Pleistocene faunal turnover on Flores. Java Homo erectus s.s. associated with 3 mammalian assemblages Ci Saat (~1.2 Ma), Trinil HK (Grenzbank- lower Bapang/ Kabuh Fm of Sangiran; ~1.0 Ma) and Kedung Brubus (U Bapang/ Kabuh Fm between middle and lower tuff of Sangiran; ~0.8 Ma). Ngandong fauna with Homo erectus soloensis <0.8 Ma. Punung and Wajak faunas with H. sapiens. On Flores two endemic island faunas: (1) dwarf Stegodon and giant tortoise Geochelone, and (2) younger fauna with Stegodon trigonocephalus, giant rats and hominid lithic industry)

Aziz, F., P.Y. Sondaar, J.J.M. Leinders & J. de Vos (1989)- Fossil faunas and early man of Java. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 6, p. 1-3.
(Brief summary of succession of seven Pleistocene mammal/ hominid assemblages of Java)

Aziz, F., P.Y. Sondaar, G.D. van den Bergh & J. de Vos (1995)- *Homo erectus* in S.E. Asia: time space and migration routes, II. The Java case. In: J. Gibert et al. (eds.) Proc. Int. Conference The hominids and their environment during the Lower and Middle Pleistocene of Eurasia, Orce 1995, p. 363-368.

Aziz, F. & G.D. van den Bergh (1995)- A dwarf *Stegodon* from Sambungmacan (Central Java, Indonesia). Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 98, 3, p. 229-241.
(New dwarf Stegodon jaw fragment found in 1992 in shallow marine Kalibeng Limestone at Sambungmacan. Age may be latest Pliocene or earliest Pleistocene. Site may have been margin of a paleo-island in Kendeng zone. Overlying clastics at this site yielded 1973 'Solo Man' Homo erectus skull cap discovery and stone tools)

Aziz, F., G.D. van den Bergh, M.J. Morwood, D.R. Hobbs, I. Kurniawan, J. Collins & Jatmiko (2009)- Excavations at Tangi Talo, central Flores, Indonesia. In: F. Aziz et al. (eds.) Pleistocene geology, palaeontology and archaeology of the Pleistocene Soa Basin, Central Flores, Indonesia, Chapter 3, Geological Survey Centre Bandung, Special Publ. 36, p. 41-58.
(Results of 1999 excavations at Tangi Talo, the oldest fossil site in the Soa basin, Flores. Basal tuffaceous interval of Ola Bula Fm with pygmy Stegodon sondaari, giant tortoise and Varanus komodoensis (Komodo dragon). Deposited shortly after massive volcanic eruption at ~900,000 yrs BP. No in situ stone artefacts found. Site probably predates arrival of hominins in basin (earlier excavations here by Sondaar et al. in 1992)

Baab, K.L. (2010)- Cranial shape in Asian *Homo erectus*: geographic, anagenetic, and size-related variation. In: C.J. Norton & D.R. Braun (eds.) Asian Paleoanthropology: from Africa to China and beyond, Springer, Chapter 6, p. 57-79.
(No strong support for linear progression in neurocranial skull shape from Sangiran to Ngandong via Sambungmacan/Ngawi)

Baab, K.L. (2016)- The place of *Homo floresiensis* in human evolution. J. Anthropological Sciences 94, p. 5-18.
(online at: www.isita-org.com/jass/Contents/2016vol94/Baab/26829572.pdf)

(Two evolutionary scenarios for small-bodied Homo floresiensis on Flores in Late Pleistocene: (1) H. floresiensis was dwarfed descendent of H. erectus, or (2) remnant of older lineage, perhaps descended from H. habilis. Could be either)

Baab, K.L. (2021)- Reconstructing cranial evolution in an extinct hominin. Proc. Royal Society (London), B, 288, 20202604, p. 1-9.
(online at: https://royalsocietypublishing.org/doi/10.1098/rspb.2020.2604)
(Homo erectus cranial fossils sorted into six major palaeodemes from sites across Africa, Eurasia and Asia spanning 1.8-0.1 Ma. H. erectus had higher individual and group variation than Homo sapiens)

Baab, K.L., K.P. McNulty & K. Harvati (2013)- *Homo floresiensis* contextualized: a geometric morphometric comparative analysis of fossil and pathological human samples. Plos One 8, 7, e69119, p. 1-11.
(online at: http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0069119)
(Geometric morphometric analyses of LB1 cranium shows Homo floresiensis is distinct from healthy modern humans and from those with pathological conditions (hypothyroidism, Laron syndrome, microcephism), and is most similar to Homo erectus (but smaller))

Baab, K.L. & Y. Zaim (2017)- Global and local perspectives on cranial shape variation in Indonesian *Homo erectus*. Anthropological Science 125, 2, p. 67-83.
(online at: https://www.jstage.jst.go.jp/article/ase/125/2/125_170413/_pdf/-char/en)
(Skull shapes of Homo erectus from Sangiran, Ngandong, Sambungmacan and Ngawi compared to H. erectus from outside of Java. Asian H. erectus fossils can be distinguished from African/Georgian ones. Late Indonesian H. erectus from sites like Ngandong, distinct from all other H. erectus groups, including older C Java fossils. Younger Sangiran fossils more closely approach Ngandong/Sambungmacan/Ngawi pattern)

Baba, H. & F. Aziz (1991)- Hominid skull fragments found in 1979 from Sangiran, Central Java. Bull. National Science Museum, Tokyo, D, 17, p. 1-8.
(online at: https://www.kahaku.go.jp/research/publication/anthropology/download/17/BNSM_D1701.pdf)
(Two new hominid skull fragments with typical Homo erectus characteristics))

Baba, H. & F. Aziz (1992)- Human tibial fragment from Sambungmachan, Java. In: T. Akazawa et al. (eds.) The evolution and dispersal of modern humans in Asia, Tokyo, p. 349-361.
(Shin bone fragment from C Java)

Baba, H., F. Aziz, Y. Kaifu, G. Suwa, R.T. Kono & T. Jacob (2003)- *Homo erectus* calvarium from the Pleistocene of Java. Science 299, 5611, p. 1384-1388.
(online at: https://ftp.soest.hawaii.edu/engels/Stanley/Textbook_update/Science_299/Baba-03.pdf)
(Homo erectus calvarium from Sambungmacan, C Java (Sambungmacan 4; Sm 4). Overall morphology intermediate between earlier and later Javanese Homo erectus)

Baba, H., F. Aziz & S. Narasaki (2000)- Restoration of the face of Javanese *Homo erectus* Sangiran 17 and re-evaluation of regional continuity in Australasia. In: W. Dong (ed.) Proc. 1999 Beijing International Symposium on Paleoanthropology, Acta Anthropologica Sinica 19 (Supplement), p. 34-40.
(Facial characteristic of restored Sangiran 17 skull do not support hypothesis of Thorne and Wolpoff of regional continuity between H. erectus and Late Pleistocene Australians)

Baba, H., F. Aziz, S. Narasaki, Sudijono, Y. Kaifu, A. Supriojo, M. Hyodo, E.E. Susanto & T. Jacob (2000)- A new hominid incisor from Sangiran, Central Java. J. Human Evolution 38, 6, p. 855-862.
(New hominid lower left central tooth found in 1997, in level between lower and upper tuffs in Bapang Fm, near village of Bukuran, ESE Sangiran dome area (latest E Pleistocene))

Baba, H., F. Aziz, S. Narasaki, Sudijono, Y. Kaifu, I. Saefudin & E.E. Susanto (2004)- Frontal bone fragment of *Homo erectus* from Sangiran, Java. Human Evolution 19, 3, p. 197-201.

(Homo erectus frontal bone fragment found in 1994 in Brangkal River floor, Sangiran area. Original stratigraphic level not known, possibly from 'Grenzbank')

Baba, H., F. Aziz & N. Watanabe (1990)- Morphology of the fossil hominid tibia from Sambungmacan, Java. Bull. National Science Museum, Tokyo, D15, p. 9-18.
(online at: http://ci.nii.ac.jp/naid/110000008554/en)
(Hominid tibial (shin bone) fragment Sm 2 collected in 1977 with other vertebrate bones from Sambungmacan, presumably from Kabuh equivalent beds. Rel. advanced character)

Bacon, A.M. & P.O. Antoine (2024)- Environments, terrestrial ecosystems and mammalian species: an overview of Southeast Asia in the Late Pleistocene. In: J. Louys et al. (eds.) Quaternary Palaeontology and Archaeology of Sumatra, Terra Australis 56, ANU Press, Canberra, Chapter 7, p. 145-182.
(online at: https://press-files.anu.edu.au/downloads/press/n12084/pdf/ch07.pdf)
(Sunda Shelf surface in late M- Late Pleistocene affected by tectonic, eustatic and climatic factors, with alternating phases of exposure and flooding. Analysis of mammalian assemblages from W Sumatra caves during two possible periods of faunal dispersion: ~130 ka (MIS 6-5 transition) and ~71 ka (MIS 4. Overall, Sundaland always less rich in large-bodied herbivore diversity. Etc.)

Bacon, A.M., K. Westaway, P.O. Antoine, P. Duringer, A. Blin, F. Demeter, J.L. Ponche, J.X. Zhao, L.M. Barnes et al. (2015)- Late Pleistocene mammalian assemblages of Southeast Asia; new dating, mortality profiles and evolution of the predator-prey relationships in an environmental context. Palaeogeogr. Palaeoclim. Palaeoecology, 422, p. 101-127.
(manuscript online at: https://espace.library.uq.edu.au/view/UQ:352803/UQ352803_OA.pdf?dsi_version =f77691238d4065e9f08494fd6b4d696d)
(On mammal remains in Late Pleistocene karst deposits in N Laos, N Vietnam, Punung (C Java) and Sibrambang (W Sumatra))

Badoux, D.M. (1959)- Fossil mammals from two fissure deposits at Punung (Java) with some remarks on migrations and evolution of mammals during the Quaternary in South East Asia. Doctoral Thesis, University of Utrecht, p. 1-151.
(Description of 'Punung mammal fauna' from karst hills of S Mountains, E Java, collected by Von Koenigswald in 1930's. (see also Storm et al. (2005) for details of locality, paleoenvironment (rainforest) and age (~0.1 Ma))

Balme, J. & S. O'Connor (2014)- Early modern humans in Island Southeast Asia and Sahul- Adaptive and creative societies with simple lithic industries. In: R. Dennell & M. Porr (eds.) Southern Asia, Australia, and the search for human origins, Cambridge University Press, Chapter 13, p. 164-174.
(online at: https://www.researchgate.net/publication/268219198_Early_modern_humans_in_Island_Southeast _Asia_and_Sahul_Adaptive_and_creative_societies_with_simple_lithic_industries)
(Review of possible settlement patterns of early modern humans across E Indonesia islands, New Guinea and northern coastal Australia, around 45 ka B.P. Earliest settlers had simple stone and bone tools and lived mainly of marine and coastal fish and shellfish. Etc.))

Balzeau, A. (2005)- Specificites des caracteres morphologiques internes du squelette cephalique chez *Homo erectus*. Doctoral Thesis, Museum National Histoire Naturelle, Paris, p. 1-394.
(online at: http://hopsea.mnhn.fr/pc/thesis/PhD_Balzeau2005.pdf)
('Internal morphologic characteristics of Homo erectus skull')

Balzeau, A. (2013)- Thickened cranial vault and parasagittal keeling: correlated traits and autapomorphies of *Homo erectus*? J. Human Evolution 64, 6, p. 631-644.

Balzeau, A. & P. Charlier (2016)- What do cranial bones of LB1 tell us about *Homo floresiensis*? J. Human Evolution 93, p. 12-24.
(No support for attribution of holotype of Homo floresiensis (LB1) from Liang Bua, Flores, to H. sapiens)

Balzeau, A., D. Grimaud-Herve & T. Jacob (2005)- Internal cranial features of the Mojokerto child fossil (East Java, Indonesia). J. Human Evolution 48, p. 535-553.

Balzeau, A., T. Jacob & E. Indriati (2002)- Structures craniennes internes de l'*Homo erectus* Sambungmacan 1 (Java, Indonesie). Comptes Rendus Palevol 1, p. 305-310)
(online at: https://www.sciencedirect.com/science/article/pii/S1631068302000489)
('Interior skull structures of the Sambungmacan Homo erectus'. CT data to reconstruct skull endocast suggest morphological character of Sambungmacan 1 is close to Ngandong hominids)

Bandet, Y., F. Semah, S. Sartono & T. Djubiantono (1989)- Premier peuplement par les mammiferes d'une region de Java Est, a la fin du Pliocene: age de la faune du Gunung Butak, pres de Kendungbrubus (Indonesie). Comptes Rendus Academie Sciences, Paris 308, p. 867-870.
(online at: https://core.ac.uk/download/pdf/39865312.pdf)
('First population by mammals of a region of E Java at the end of the Pliocene: age of the Gunung Butak fauna near Kedungbrubus'. Along flanks of Gunung Butak marine sediments abruptly overlain by coarse and slightly transported volcanic breccia. Fossil mammals appear just above, in fluviatile and lahar beds. Top of basal breccia dated 1.87 Ma. Emerged tongues of land due to volcanoes colonized by mammals by latest Pliocene)

Bandi, H.G. (1951)- Die Obsidianindustrie der Umgebung von Bandung in Westjava. Sudsee-Studien, Museum fur Volkerkunde, Basel, Felix Speiser Memorial Volume, p. 127-161.
('The obsidian industry in the surroundings of Bandung in West Java'.Studies of 'Mesolithic', bronze-age obsidian microlithic tools from Bandung area, collected by W. Rothpletz in 1940s, mainly from Dago area)

Barker, G. (2022)- The early settlement of Island Southeast Asia. In: C.F.W. Higham & N.C. Kim (eds.) The Oxford Handbook of Early Southeast Asia, Chapter 4, p. 92-123.

Barker, G., H. Barton, P. Beavitt, M. Bird, P. Daly, C. Doherty, D. Gilbertson, C. Hunt et al. (2014)- Prehistoric foragers and farmers in South-East Asia: Renewed investigations at Niah Cave, Sarawak. Proc. Prehistoric Society 68, p. 147-164.
(Niah cave in Sarawak has been used by humans since at least 45,000 years ago)

Barker, G., H. Barton, M. Bird, P. Daly, I. Datan, A. Dykes, L. Farr, D. Gilbertson, Haarrisson et al. (2007)- The `human revolution' in lowland tropical Southeast Asia: the antiquity and behavior of anatomically modern humans at Niah Cave (Sarawak, Borneo). J. Human Evolution 52, 3, p. 243-261.
(Anatomically modern humans in Niah cave, Sarawak, N Borneo. New dates and lithostratigraphy relate Deep Skull to evidence of human activity between ~46,000- 34,000 years ago)

Barker, G., C. Hunt, H. Barton, C. Gosden, S. Jones, L. Lloyd-Smith, L. Farr et al. (2017)- The 'cultured rainforests' of Borneo. Quaternary International 448, p. 44-61.
(Investigation of the archaeology and palaeoecology of the Niah Caves of Sarawak shows 50,000-year record of Homo sapiens' interactions with rainforest on coastal lowlands)

Barker, G. & M.B. Richards (2013)- Foraging-farming transitions in island Southeast Asia. J. Archaeological Method and Theory 20, p. 256-280.
(online at: www.researchgate.net/profile/Martin-Richards-2/publication/233854184_Foraging-Farming_Transitions_in_Island_Southeast_Asia/links/5453c4df0c Etc)
(Still much debate on when and why Homo sapiens in SE Asia switched from hunting-gathering to farming. Much of this discussion based on Niah cave excavations in N Borneo)

Bartstra, G.J. (1974)- Notes about Sangiran (Java, Indonesia). Quartar 25, p. 1-11.
(online at: www.quartaer.eu/pdfs/1974/1974_01_bartstra.pdf)
(Brief review of Sangiran dome Pleistocene stratigraphy, ~10km N of Solo, and localities of hominid fossils. Author believes that many of the hominid stone tools ("flake industries") of Sangiran are pseudo-artifacts. There are also some real artifacts at Sangiran, but their presumed M Pleistocene age is doubted. Most visitors

to Sangiran in late 1930s-1950s (De Terra, Movius, van Heekeren) appear to agree that the artifacts come from the (basal) Notoporu Beds volcanic breccias)

Bartstra, G.J. (1976)- Contributions to the study of the Palaeolithic Patjitan culture, Java, Indonesia. Thesis Rijksuniversiteit te Groningen, Brill, Leiden, p. 1-121.
(Abstract online at: https://pure.rug.nl/ws/portalfiles/portal/14561086/File0004.PDF)
(Study of Southern Mountains (Gunung Sewu) of C Java geology and geomorphology, incl. Miocene Wonosari limestone karst terranes, Pleistocene river terraces, etc., based on 1972-1973 fieldwork. Pacitanian Paleolithic artifacts, first(?) discovered by Von Koenigswald in 1935 in Baksoko River terrace gravels in Southern Mountains of C Java. Mammals in nearby Punung fissure deposit of M Pleistocene age (Djetis and Trinil associations; Hooijer appendix))

Bartstra, G.J. (1977)- The height of the river terraces in the transverse Solo valley in Java. Modern Quaternary Research in Southeast Asia 3, Balkema, Rotterdam, p. 143-155.
(Quaternary terraces in transverse valley of Solo River N of Ngawi, previously studied by De Terra (1943), Sartono (1976), etc. Highest terrace up to 40-50m above present Solo river level (T1; poorly preserved), second terrace at ~20m above rive (T2), third ~5m above river (T3; best preserved; location of villages). Ngandong Solo Man fossils in T2/20m terrace)

Bartstra, G.J. (1977)- Walanae Formation and Walanae terraces in the stratigraphy of South Sulawesi (Celebes, Indonesia). Quartar 27/28, p. 21-30.
(online at: https://journals.ub.uni-heidelberg.de/index.php/qu/article/view/80461/74507)
(Geologic setting of terraces of Walanae River, which contain Pleistocene vertebrate fossils (Archidiskodon-Celebochoerus fauna of Hooijer, 1948), and stone artefacts (Tjabenge industry). Author believes stone artifacts to be younger than S Sulawesi fossils)

Bartstra, G.J. (1978)- The age of the Djetis Beds in East and Central Java. Antiquity 52, 204, p. 56-58.
(On age of Jetis Fauna (incl. Homo erectus) from Mojokerto area)

Bartstra, G.J. (1978)- The Patjitan culture: a preliminary report on new research. In: F. Ikawa-Smith (ed.) Early Palaeolithic in South and East Asia, Mouton Publishers, The Hague, p. 29-36.
(Paleolithic 'Pacitanian' relatively advanced stone tools from along Baksoko River near Punung, S coast of E Java, first described by Von Koenigswald (1936). 1500 additional artifacts collected in 1972 excavations. Three fluvial terraces, all with stone artifacts but unfossiliferous. Pacitan culture may be work of Homo soloensis)

Bartstra, G.-J. (1978)- Note on new data concerning the fossil vertebrates and stone tools in the Wallanae valley in South Sulawesi (Celebes). Modern Quaternary Research in Southeast Asia 4, p. 71-77.

Bartstra, G.J. (1982)- The river-laid strata near Trinil, site of *Homo erectus*, Java, Indonesia. Modern Quaternary Research in Southeast Asia 7, Balkema, Rotterdam, p. 97-130.
(Historic review of stratigraphic interpretations of Pleistocene Trinil Beds of C Java, the site of first Pithecanthropus and of abundant vertebrate fossils collected by Dubois, Selenka, etc. Two fluvial sand horizons along Solo River at Trinil: (1) Lower horizon (Kabuh Beds) of M and Late Pleistocene age; (2) upper horizon of Late Pleistocene Solo River terrace deposits. Sands composed of andesitic material and unconformably overlie Late Neogene marine marls)

Bartstra, G.J. (1982)- *Homo erectus erectus*: the search for his artifacts. Current Anthropology 23, 3, p. 318-320.
(online at: https://www.rug.nl/research/portal/files/66717894/2742319.pdf)
(Years of fieldwork in C Java between 1977-1981 failed to find conclusive evidence for E-M Pleistocene tools used by Homo erectus. Many stone artifacts found across E Java, but all younger than Homo erectus and may be from modern Homo. Older small crude flakes interpreted as hominid implements 'may just be stones')

Bartstra, G.J. (1983)- The fauna from Trinil, type locality of *Homo erectus*: a reinterpretation, Comment I: The vertebrate bearing deposits of Kedungbrubus and Trinil, Java, Indonesia. Geologie en Mijnbouw 62, 2, p. 329-336.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0TXdLZ3dtNVloTHc/view)
(De Vos et al. (1982) suggested classic Pleistocene vertebrate stratigraphy of Java as established by Von Koenigswald is incorrect, calling Jetis·fauna younger than Trinil fauna. On stratigraphic grounds this view is untenable and Van Koenigswald's stratigraphy is essentially correct. See also Hooijer 1983))

Bartstra, G.J. (1983)- Some remarks upon fossil man from Java, his age, and his tools. Bijdragen Taal Land en Volkenkunde 139, 4, Leiden, p. 421-434.
(online at: www.kitlv-journals.nl/index.php/btlv/article/view/2001/2762)
(Review of stratigraphy and hominid fossils of C Java. Solo River deposits near Trinil two units: (1) upper unit typical modern Solo river terrace deposits with Homo soloensis, (2) lower unit Pleistocene clays, silts, sands and gravels, with Homo erectus)

Bartstra, G.J. (1984)- Dating the Pacitanian: some thoughts. In: P. Andrews & J.L. Franzen (eds.) The early evolution of Man with special emphasis on South East Asia and Africa, Courier Forschungsinstitut Senckenberg 69, p. 253-258.

Bartstra, G.J. (1985)- Sangiran, the stone implements of Ngebung and the Paleolithic of Java. Modern Quaternary Research in Southeast Asia 9, Balkema, Rotterdam, p. 99-113.
(Review of Late Pleistocene archaeological stone industries from Ngebung top-gravel, in Sangiran Dome (also described by Von Koenigswald, 1935))

Bartstra, G.J. (1987)- Late *Homo erectus* or Ngandong man of Java. Palaeohistoria 29, p. 1-7.
(online at: https://ugp.rug.nl/Palaeohistoria/article/download/24867/22315)
(Review of Ngandong man ('Homo soloensis') from 'High Terrace' deposits of Solo river in Kendeng zone of Java, Generally regarded as last representative of Homo erectus in SE Asia. Associated with rich mammal fauna, mainly bovids, indicative of open forest environment. Oldest Solo terrace sediments in C Java must date from beginning of N-ward directed drainage pattern and probably Late Pleistocene in age. U-series ages from Ngandong bone samples mainly between ~40-80 ka. No in-situ stone implements, but nearby surface finds of small chalcedony artefacts ('Ngandong Industry') (chalcedony= silicified coral limestone; Bartstra 1989))

Bartstra, G.J. (1989)- Sangiran for the archaeologist. A short guide for students. Berkala Arkeologi 10, 2, p. 1-20.
(online at: https://berkalaarkeologi.kemdikbud.go.id/index.php/berkalaarkeologi/article/view/540/488)
(Review of the Sangiran Dome site in C Java, where famous Pithecanthropus and Meganthropus (Homo erectus) fossils were found)

Bartstra, G.J. (1994)- Indonesia in the period of *Homo habilis* and *Homo erectus*. In: S.J. de Laet et al. (eds.) History of humanity: prehistory and the beginnings of civilization, 1, Chapter 8, UNESCO, Paris, p. 275-292.
(online at: https://ia801701.us.archive.org/21/items/history-of-humanity-1/History-of-Humanity%201.pdf)
(Popular reviews of Java Pleistocene hominid sites at Trinil and Sangiran, vertebrate stratigrapy, handaxes of Pacitan and Homo erectus, etc.)

Bartstra, G.J. (1994)- Indonesia in the period of *Homo sapiens neanderthalensis.* In: S.J. de Laet et al. (eds.) History of humanity: prehistory and the beginnings of civilization, 1, Chapter 17, UNESCO, Paris, p. 446-456.
(online at: https://ia801701.us.archive.org/21/items/history-of-humanity-1/History-of-Humanity%201.pdf)
(Popular reviews of two hominin sites (1) Ngebung Hill in Sangiran, C Java, and (2) Ngandong, a younger site in Solo Riverterrace deposits in E Java. Discovered in 1931;and hominid fossils were decribed as Homo soloensis by Oppenoorth (debated whether H. soloensis is advanced Homo erectus or a Neanderthal (Von Koenigswald; less likely?)

Bartstra, G.J. (1997)- A fifty years commemoration: fossil vertebrates and stone tools in the Walanae valley, South Sulawesi, Indonesia. Quartaer 47/48, p. 29-50.
(online at: www.quartaer.eu/pdfs/1997/1997_02_bartstra.pdf)
(Review of history of discovery of Pleistocene mammals and hominid artifacts along the Walanae River in S Sulawesi in 1947 by archeologist H.R. van Heekeren and works of vertebrate paleontologist D.A. Hooijer)

Bartstra, G.J. (1998)- Short history of the archaeological exploration of the Maros caves in South Sulawesi. In: G.J. Bartstra (ed.) Bird's Head approaches; Irian Jaya studies; a programme for interdisciplinary research, Modern Quaternary Research in Southeast Asia, Balkema, Rotterdam, 15, p. 193-210.
(online at: https://www.papuaerfgoed.org/sites/default/files/collectie/files/2006-05/Bartstra_1998_Birds.pdf)
(Review of archaeological research in caves of Maros kast area of S Sulawesi, known for its prehistoric Toalean culture. The first European scientists to unearth prehistoric remains in Maros caves were the Swiss Sarasin cousins in 1902, who assigned to stone tools to the prehistoric Toalean culture. Additional work in area by Van Stein Callenfels in late 1933 and 1937 at Lamoncong, Van Heekeren in 1936-1937 and 1950. Modern archeological work since late 1960s)

Bartstra, G.J. & Basoeki (1989)- Recent work on the Pleistocene and the Palaeolithic of Java. Current Anthropology 30, 2, p. 241-244.
(online at: https://pure.rug.nl/ws/portalfiles/portal/66717792/2743557.pdf)
(1989 status review, mainly on C Java Pleistocene stone artifacts)

Bartstra, G.J., M. Basoeki & B. Santosa Azis (1976)- Solo Valley research 1975 Java, Indonesia. Modern Quaternary Research in Southeast Asia 2, Balkema, Rotterdam, p. 23-36.
(Brief review of Pleistocene geology along Solo River in Trinil and Ngawi areas. Discovery of stone implements in high terrace gravels)

Bartstra, G.J. & D.A. Hooijer (1992)- New finds of fossil vertebrates from Sulawesi, Indonesia. Lutra 35, p. 113-122.

Bartstra, G.J., D.A. Hooijer, B. Kallupa & M.A. Akib (1992)- Notes on fossil vertebrates and stone tools from Sulawesi, Indonesia, and the stratigraphy of the northern Walanae depression. Palaeohistoria (University of Groningen) 33/34, p. 1-18.
(online at: https://ugp.rug.nl/Palaeohistoria/article/view/25054/22512)
(Pleistocene Archidiskodon-Celebochoerus vertebrate fauna and artifacts of the Cabenge Industry. Fossils and artifacts may not be contemporaneous. Singkang embayment/ Tempe depression separates SW peninsula of Sulawesi from rest of island and was covered by sea until recently)

Bartstra, G.J., S.G. Keates, Basoeki & B. Kallupa (1991)- On the dispersion of *Homo sapiens* in Eastern Indonesia: the Palaeolithic of South Sulawesi. Current Anthropology 32, 3, p. 317-321.
(online at: https://pure.rug.nl/ws/files/66717477/2743781.pdf)
(Homo sapiens reached Sulawesi around 50,000 B.P. (Walanae Fm); Homo erectus never reached Sulawesi)

Bartstra, G.J., S. Soegondo & A. van der Wijk (1988)- Ngandong man: age and artifacts. J. Human Evolution 17, 3, p. 325-337.
(Homo soloensis found in 1930's in High Terrace of Solo River near Ngandong, with associated small stone implements. Th/U ages for Ngandong vertebrate remains suggest Upper Pleistocene age (~30-100 ka?))

Basilia, P., J.J. Miszkiewicz, J. Louys, U.P. Wibowo & G.D. van den Bergh (2023)- Insights into dwarf stegodon (*Stegodon florensis florensis*) palaeobiology based on rib histology. Annales de Paleontologie 109, 4, 102654, p. 1-12.
(Stegodon florensis florensis example of dwarf insular stegodon from E Middle Pleistocene bonebeds of Mata Menge, So'a Basin, in Flores. First insights into Stegodon rib bone histology, suggesting long-lived individuals in advanced age)

Basilia, P., J.J. Miszkiewicz, K. Nganvongpanit, J. Zaim, Y. Rizal, Aswan, M.R. Puspaningrum et al. (2023)- Bone histology in a fossil elephant (*Elephas maximus*) from Pulau Bangka, Indonesia. Historical Biology 35, 8, p. 1356-1367.
(online at: https://repository.naturalis.nl/pub/801043/Basilia-2022-Bone-histology-in-a-fossil-elephant-A.pdf)
(Something with elephantoid bone structure)

Basilia, P., J.J. Miszkiewicz, J. Zaim, Y. Rizal, Aswan, M.R. Puspaningrum, A. Trihascaryo, G.J. Price & J. Louys (2024)- Investigating super osteons in fossil Asian elephant (*Elephas maximus*) bone from Bangka Island, southeastern Sumatra. In: L. Louys et al. (eds.) Quaternary palaeontology and archaeology of Sumatra, Australian National University Press, Canberra, p. 183-198.
(online at: https://press-files.anu.edu.au/downloads/press/n12084/pdf/ch08.pdf)
(Study of bone structure in fossil elephant fragments from Late Pleistocene alluvial deposits on Bangka Island)

Bednarik, R.G. (1997)- The initial peopling of Wallacea and Sahul. Anthropos 92, p. 355-367.
(online at: www.ifrao.com/wp-content/uploads/2015/02/97Wallacea.pdf)
(Review of Pleistocene migrations of hominids into Indonesia and Australia)

Bednarik, R.G. (2000)- Pleistocene Timor: some corrections. Australian Archaeology 5, p. 16-20.
(Review of Pleistocene mammal fossils (Stegodon, giant tortoise) and some questionable relicts of Pleistocene hominids on Timor)

Bednarik, R.G. (2002)- The maritime dispersal of Pleistocene humans. Migration and Diffusion 3, 10, p. 6-33.
(online at: www.ifrao.com/wp-content/uploads/2014/06/dispersal.pdf)

Bello, S.M. & C.B. Stringer (2025)- The newly uncovered Madura Strait fossil assemblage and its role in Pleistocene hominin dispersals in Southeast Asia. Quaternary Environments and Humans, 100070, p. 1-2 *(in press)*
(online at: https://www.sciencedirect.com/science/article/pii/S2950236525000143)
(Introduction to special volume with four papers on new discovery of Pleistocene mammals in Madura Straits N of Surabaya)

Bellwood, P. (1987)- The prehistory of island Southeast Asia: a multidisciplinary review of recent research. J. World Prehistory 1, 2, p. 171-224.
(Sundaland region in W of SE Asia archipelago and isolated islands of Wallacea in E witnessed complex trajectories of human movement and evolution during Pleistocene)

Bellwood, P. (1997)- *Homo erectus* in Sundaland. In: Prehistory of the Indo-Malaysian Archipelago, Chapter 2, ANU Press, p. 39-68.
(online at: http://press-files.anu.edu.au/downloads/press/p80041/pdf/ch0273.pdf)

Bellwood, P.S. (1998)- Bridging Sunda and Sahul: The archaeological significance of the Aru Islands, southern Moluccas. In: G.J. Bartstra (ed.) Bird's Head approaches; Irian Jaya studies; a programme for interdisciplinary research, Modern Quaternary Research in Southeast Asia 15, Balkema, Rotterdam, p. 157-177.
(online at: https://www.papuaerfgoed.org/sites/default/files/collectie/files/2006-05/Bartstra_1998_Birds.pdf)

Bellwood, P.S. (2007)- Prehistory of the Indo-Malaysian Archipelago. Australia National University (ANU) Press, Canberra, p. 1-384.
(*online at: http://epress.anu.edu.au/pima/pdf/pima-whole.pdf*
(3rd edition of 1985 textbook)

Bellwood, P.S. (2017)- First Islanders: prehistory and human migration in island Southeast Asia, Wiley-Blackwell, p. 1-384.
(Recent review of history of hominids in SE Asia)

Bellwood, P.S. (2017)- *Homo erectus* and *Homo floresiensis*- Archaic hominins in island Southeast Asia. In: P. Bellwood et al. (eds.) First Islanders: prehistory and human migration in island Southeast Asia, Wiley, Chapter 3, p. 34-85.
(Discussion of lithic stone tool industries reportedly associated with Homo erectus in various regions of island SE Asia. Two categories: (1) 'chopper/chopping-tool industries' characterized by Java Pacitanian industry and supposedly work of Homo erectus; (2) 'pebble and flake industries' more characteristic of early Homo sapiens)

Bellwood, P.S., G. Nitihaminoto, G. Irwin, Gunadi, A. Waluyo & D. Tanudirjo (1998)- 35,000 years of prehistory in the northern Moluccas. In: G.J. Bartstra (ed.) Bird's Head approaches; Irian Jaya studies; a programme for interdisciplinary research, Modern Quaternary Research in Southeast Asia 15, Balkema, Rotterdam, p. 233-275.
(online at: https://www.papuaerfgoed.org/sites/default/files/collectie/files/2006-05/Bartstra_1998_Birds.pdf)
(Review of archeological records of Homo sapiens in the N Moluccas, dating back to >32,000 years))

Berghuis, H.W.K., Y. Kaifu, U.P. Wibowo, T. van Kolfschoten, I.Sutisna, S. Noerwidi, S. Adhityatama, G. van den Bergh, E. Pop et al. (2025)- The late Middle Pleistocene *Homo erectus* of the Madura Strait, first hominin fossils from submerged Sundaland. Quaternary Environments and Humans 3, 2, 100068, p. 1-11
(online at: https://www.sciencedirect.com/science/article/pii/S295023652500012X)
(Discovery of two hominin cranial fragments from submerged Sunda Shelf, retrieved during dredging work in Madura Strait, N of Surabaya, off NE Java coast. Specimens derive from sandy fill of late M Pleistocene paleovalley of Solo River and consist of two frontal and parietal fragments, resembling H. erectus crania from Sambungmacan. Madura Strait hominins probably part of MIS6 population that lived along Solo River, flowing E-eastward over exposed shelf area of Madura Strait, during rel. dry late M Pleistocene)

Berghuis, H.W.K., G. van den Bergh, T. van Kolfschoten, U.P. Wibowo,I. Kurniawan, S. Adhityatama, I. Sutisna et al. (2025)- First vertebrate faunal record from submerged Sundaland: the late Middle Pleistocene, hominin-bearing fauna of the Madura Strait. Quaternary Environments and Humans 3, 2, 100047, p. 1-48.
(online at: https://www.sciencedirect.com/science/article/pii/S2950236524000458)
(First vertebrate fossils from subsea Sundaland, from lowstand valley of Solo River in Madura Strait, N of Surabaya, OSL-dated to ~162 and ~ 119 ka (late M Pleistocene). At least 36 species: reptiles (turtles, crocodilians, varanids, pythons), sharks (incl. extinct Hemipristis serra) and stingrays. Lowland plains populated by herbivores (Hippopotamidae, Bovidae, Cervidae, Proboscidea (Stegodon, Elephas), suggesting grass-dominated environment similar to contemporaneous fossil assemblage of Ngandong (Java) (but with more estuarine species, fewer carnivores). Also rare Varanus komodoensis, Macaca sp. and Homo erectus)

Berghuis, H.W.K., T. van Kolfschoten, S. Adhityatama, S.R. Troelstra, S. Noerwidi, R.A.Suriyanto, U.P. Wibowo et al. (2022)- The eastern Kendeng Hills (Java, Indonesia) and the hominin-bearing beds of Mojokerto, a re-interpretation. Quaternary Science Reviews 295,107692, p. 1-25.
(online at: www.sciencedirect.com/science/article/pii/S0277379122003237)
(Re-interpretation of 1000m thick exposed marine-to-fluvial Late Pliocene-Pleistocene deposits in paleo-embayment in E Kendeng zone. Fluvial conglomerate with Mojokerto Homo erectus skullcap is tied to MIS14, with age of ~550ka (younger by 1 Myrs or more than previous age interpretations!). All earlier radiometric and paleomagnetic work discarded as wrong (this radical change in age is based mainly on sequence stratigraphic reasoning in a non-marine series, with no support from any new independent age data; JTvG)

Berghuis, H.W.K., T. van Kolfschoten, S. Adhityatama, S.R. Troelstra, S. Noerwidi, R.A.Suriyanto, U.P. Wibowo et al. (2023)- Response to the comment by Huffman and Zaim. Quaternary Science Reviews 307, 108059, p. 1-3.
(online at: https://www.sciencedirect.com/science/article/pii/S0277379123001075)
(Response to Critical discussion of Berghuis et al. (2022) paper on stratigraphic context and age of the Homo erectus child skull from Mojokerto, E Java)

Berghuis, H.W.K., T. van Kolfschoten, U.P. Wibowo, I. Kurniawan, S. Adhityatama, I. Sutisna et al. (2025)- The taphonomy of the Madura Strait fossil assemblage, a record of selective hunting and marrow processing by late Middle Pleistocene Sundaland hominins. Quaternary Environments and Humans 100055, p. 1-21. *(in press)*
(online at: https://www.sciencedirect.com/science/article/pii/S2950236524000537)
(Late M Pleistocene Madura Strait bones assemblage reflect fluvial size-sorting, but no significant rounding, pointing to short transportation distances. Skeletal remains accumulated in fluvial valley fill or in overlying transgressive lag during subsequent transition to estuarine conditions (linked to peak-highstand conditions around MIS5e (~123 ka)). Extensive fragmentation of ruminant limb bones points to hominin bone battering and marrow processing. Cut marks on turtle bones show that hominins also fed on these aquatic species)

Berghuis, H.W.K., A. Veldkamp, S. Adhityatama, S.L. Hilgen, I. Sutisna, D.H. Barianto, E.A.L. Pop, E.A.L. Reimann, D. Yurnaldi et al. (2021)- Hominin homelands of East Java: revised stratigraphy and landscape reconstructions for Plio-Pleistocene Trinil. Quaternary Science Reviews 260, 106912, p. 1-27.
(online at: www.sciencedirect.com/science/article/pii/S0277379121001190)
(Restudy of Pleistocene stratigraphy of Trinil area. Seven fluvial terraces identified and correlated with terraces of Kendeng Hills (e.g. hominin-bearing Ngandong terrace). Five stratigraphic units: Kalibeng deep marine, Padas Malang shallow marine, Batu Gajah volcanic breccias and black clays, Trinil ('Kabuh') fluvial and volcanoclastic (all S-dipping pre-terrace deposits) and Solo Formations. Between ~550 and 350 ka area part of Ngawi Lake Basin. After 350 ka role was taken over by perennial Solo River)

Berghuis, H.W.K., A. Veldkamp, S. Adhityatama, T. Reimann, A. Versendaalf, I. Kurniawan, E. Pop, T. van Kolfschoten & J.C.A. Joordens (2025)- A late Middle Pleistocene lowstand valley of the Solo River on the Madura Strait seabed, geology and age of the first hominin locality of submerged Sundaland. Quaternary Environments and Humans 3, 2, 100042, p. 1-18.
(online at: https://www.sciencedirect.com/science/article/pii/S2950236524000409)
(Marine sand extraction work in Madura Strait, N of port of Surabaya, yielded vertebrate-rich sandstones. >6000 fossils retrieved from dredged sand. Fossiliferous sandstones are from Solo River lowstand valley, OSL-dated to ~162 and ~119 ka, linking it to lowstand of MIS6. Top of valley fill marine sandstones, probably during MIS5e highstand. Similar age as Solo terrace of Ngandong, with youngest record of Homo erectus)

Berry, E.W. (1916)- The environment of the ape man. The Scientific Monthly 3, 2, p. 161-169.
(Early discussion of environmental setting of Pleistocene Pithecanthropus erectus (Java man). Recognized that during Pleistocene glacial lowstands Sumatra and Java were connected with Borneo and Malay Peninsula, forming landmass. Java Pleistocene mammal faunas of Siamese and Indian affinity and believed to have migrated from SE Asia mainland in NW)

Bettis, E.A., A.K. Milius, S.J. Carpenter, R. Larick, Y. Zaim, Y. Rizal, R.L. Ciochon, S.A. Tassier-Surine, D. Murray, Suminto & S. Bronto (2009)- Way out of Africa: Early Pleistocene paleoenvironments inhabited by *Homo erectus* in Sangiran, Java. J. Human Evolution 56, 1, p. 11-24.
(online at: www.academia.edu/3580806/Way_out_of_Africa_Early_Pleistocene_paleoenvironments_in Etc.)
(Stratigraphy and paleosols at Sangiran, C Java, document environments of Homo erectus in E Pleistocene. Earliest human immigrants encountered low-relief lake-margin landscape dominated by moist grasslands with open woodlands in driest positions. By 1.5 Ma, large streams filled lake and landscape became more riverine. Long-term shift toward regional drying or longer dry seasons through E Pleistocene)

Bettis, E.A., Y. Zaim, R.R. Larick, R.L. Ciochon, Suminto, Y. Rizal, M. Reagan & M. Heizler (2004)- Landscape development preceding *Homo erectus* immigration into Central Java, Indonesia: the Sangiran Formation Lower Lahar. Palaeogeogr. Palaeoclim. Palaeoecology 206, p. 115-131.
(online at: https://www.academia.edu/3580813/Landscape_development_preceding_Homo_erectus_immigration_into_Central_Java_Indonesia_the_Sangiran_Formation_Lower_Lahar)
(Sangiran Lower Lahar Unit debris flow age 1.90 Ma, terminates Late Pliocene shallow marine sedimentation. LLU likely originated from sector collapse on neighboring volcanic edifice.)

Bettis, E.A., Y. Zaim & Y. Rizal (2009)- Plio-Pleistocene climatic and volcanic controls on high to moderate accommodation space systems in the Solo Basin, Central Java, Indonesia. AAPG Hedberg Conference Variations in fluvial-deltaic and coastal reservoirs deposited in tropical environments, Jakarta 2009, p. 1-3.
(online at: www.searchanddiscovery.com/abstracts/pdf/2010/hedberg_indonesia/abstracts/ndx_bettis.pdf)
(Extended abstract. Late Pliocene- Pleistocene marginal marine, lacustrine and fluvial sediments exposed in Sangiran Dome interpreted in terms of interactions between tectonics and climate change over past 2 My. n Abrupt change from limnic -plaudal to fluvial environments marks contact between Sangiran and overlying Bapang (Kabuh) Fm).BapangFm 30-50 m thick, three upward-fining fluvial cycles. Overlying 20-40 m thick Pojarjar (Notoporo) Fm fills large channels incised into Bapang Fm; with common fine-grained volcaniclastics and two laharic diamictons, Upper and Uppermost Lahars)

Beyer, H.O. (1947)- Outline review of Philippine archaeology by islands and provinces. The Philippine J. Science 77, p. 205-374.

Beyer, H.O. & D.J. Steinberg (1957)- New finds of fossil mammals from the Pleistocene strata of the Philippines. National Research Council of the Philippines, Quezon City, Bull. 41, p. 220-238.

Bibi, F. & G. Metais (2016)- Evolutionary history of large herbivores of South and Southeast Asia (Indomalayan Realm). In: F.S. Ahrestani & M. Sankaran (eds.) The ecology of large herbivores in South and Southeast Asia, Springer Verlag, p. 15-88.

Bilsborough, A. (2005)- *Homo erectus* revisited: aspects of affinity and diversity in a Pleistocene hominin species. Anthropologie 43, 2-3, p. 129-158.
(No convincing morphological case for differentiating early African specimens (H. ergaster) or 'Meganthropus' material from Homo erectus. Long interval between early (Sangiran- Trinil- Kedung Brubus) and latest H. erectus (Ngandong- Sambungmacan). Regional trends include increase in brain size and cranial robusticity, dental reduction, etc. Homo floresiensis probably derived from SE Asian Homo erectus via transilience event and selection for endemic dwarfing)

Bird, M.I., D. Taylor & C. Hunt (2005)- Palaeoenvironments of insular Southeast Asia during the last glacial period; a savanna corridor in Sundaland? Quaternary Science Reviews 24, 20-21, p. 2228-2242.
(online at: www.researchgate.net/publication/313759617_Paleoenvironments_of_insular_Southeast_Asi Etc.)
(Geomorphology, palynology, biogeography and vegetation/climate modelling suggests N-S 'savanna corridor' through Sundaland continent during Last Glacial Period at time of lowered sea-level. Minimal interpretation of 50-150 km wide zone of open savanna vegetation along divide between S China and Java Seas, forming land bridge between Malay Peninsula, Sumatra, Java and Borneo and served as barrier to dispersal of rainforest-dependent species between Sumatra and Borneo. Savanna corridor may have provided convenient route for rapid early dispersal of modern humans through region and on into Australasia)

Blain, H. (2012)- An environmental tale from Pleistocene Java. Reconstructing dietary niche and palaeoenvironment by applying stable isotope analysis to selected fossil fauna From Trinil (Java, Indonesia). Master of Arts Thesis, University of Leiden, Faculty of Archaeology, p. 1-61.
(online at: https://openaccess.leidenuniv.nl/handle/1887/19730)
(Dietary and water resource niches for selected fossil bovids from Trinil HK, Java determined from stable isotopes of carbonate fraction of teeth and bones of fossil aquatic fauna (Trinil and Sangiran) suggests Pleistocene environment of site more diverse environment than just grassland)

Bocherens, H., F. Schrenk, Y. Chaimanee, O. Kullmer, D. Morike, D. Pushkina & J.J. Jaeger (2017)- Flexibility of diet and habitat in Pleistocene South Asian mammals: implications for the fate of the giant fossil ape *Gigantopithecus*. Quaternary International 434, p. 148-155.
(Giant fossil ape Gigantopithecus blacki from SE Asia survived until ~100,000 years ago. Known only from isolated teeth and lower jaw fossils. Carbon isotopes of tooth enamel from N Thailand suggest Gigantopithecus was forest-dweller with vegetarian diet. Demise possibly due to forest reduction during glacial periods)

Boediharto, R. (1964)- New finds of vertebrate layers in the Wonogiri and Wonosari areas, Central Java. Geological Survey Indonesia Bull. 1, 2, p. 47-49.

Boivin, N., D.Q. Fuller, R. Dennell, R. Allaby & M.D. Petraglia (2013)- Human dispersal across diverse environments of Asia during the Upper Pleistocene. Quaternary International 300, p. 32-47.
(New, more complex out-of-Africa scenario involving multiple exits, varying terrestrial routes, a sub-divided African source population, slower progress to Australia, and interbreeding with archaic varieties of Homo)

Bonatz, D., D. Neidel & M.L. Tjoa-Bonatz (2006)- The megalithic complex of highland Jambi: an archaeological perspective. Bijdragen Taal Land en Volkenkunde 162, 4, 490-522.
(online at: https://brill.com/view/journals/bki/162/4/article-p490_4.xml?language=en)
(Summary of 2003 and later German archeological work on carved megaliths scattered across Jambi/Barisan highlands. Most likely age believed to be 11th-13th centuries AD, during time of Srivijaya Kingdom hegemony in eastern Sumatran lowlands.)

Bonde, N. & B. Westergaard (2004)- Progress in hominid classification: cladistic approaches. In: Miscelanea en homenaje a Emiliano Aguirre, Paleoantropologia, p. 37-57.
(Elegant general review of hominid evolution)

Borel, A., R. Cornette & M. Baylac (2017)- Stone tool forms and functions: a morphometric analysis of modern humans' stone tools from Song Terus Cave (Java, Indonesia). Archaeometry 59, 3, p. 455-471.
(Stone industries from beginning of Holocene of SE Asia difficult to characterize typo-technologically)

Borel, A., C. Gaillard, M.H. Moncel, R. Sala, E. Pouydebat, T. Simanjuntak & F. Semah (2013)- How to interpret informal flakes assemblages? Integrating morphological description, usewear and morphometric analysis gave better understanding of the behaviors of anatomically modern human from Song Terus (Indonesia). J. Anthropological Archaeology 32, 4, p. 630-646.
(Analysis of thousands of Holocene (~11-5 ka) hominid stone tools from upper ('Keplek') levels of Song Terus cave in Southern Mountains of Central Java)

Bouteaux, A. (2005)- Paleontologie, paleoecologie et taphonomie des mammiferes du Pleistocene moyen ancien du site a hominides de Sangiran (Java central, Indonesie). Doct. Thesis Museum National Histoire Naturelle, Paris, p. 1-368.
(online at: http://hopsea.mnhn.fr/pc/thesis/PhD_Bouteaux2005.pdf)
('Paleontology, paleoecology and taphonomy of the E-M Pleistocene mammals from the Sangiran hominid site, Central Java')

Bouteaux, A. (2008)- Etude taphonomique d'assemblages fauniques de sites a *Homo erectus* du dome de Sangiran (Pleistocene moyen, Java central, Indonesie). Annales de Paleontologie 94, 4, p. 229-243.
('Taphonomic study of faunal assemblages of Homo erectus sites at Sangiran Dome (M Pleistocene, Central Java, Indonesia'. Most bone assemblages from H. erectus sites come from fluvial volcanic-sedimentary Kabuh layers (E-M Pleistocene). Herbivores dominate assemblages (large bovids like Bubalus palaeokerabau or Bibos palaesondaicus and smaller cervids like Axis lydekkeri). Carnivores are rare. High degree of fragmentation of fossils related to fissuration and fluviatile transport)

Bouteaux, A. & A.M. Moigne (2010)- New taphonomical approaches: the Javanese Pleistocene open-air sites (Sangiran, Central Java). Quaternary International 223-224, p. 220-225.
(online at: https://www.academia.edu/37335828/New_taphonomical_approaches_The_Javanese_Pleistocene_open_air_sites_Sangiran_central_Java_)
(Excavations in Sangiran Dome produced numerous mammal fossils, including Homo erectus. Bones most common in M Pleistocene fluvial Kabuh Fm volcanoclastics, dominated by teeth and extremities of large bovids and smaller cervids, mostly in fragments. Modification of assemblages by water action. Carnivores and traces of their actions rare. Anthropic influence at Ngebung 2 site, with occurrence of lithic artifacts)

Bouteaux, A., A.M. Moigne & T. Jacob (2008)- Palaeontology, palaeoecology and taphonomy of Middle Pleistocene: mammals in the hominid site of Sangiran dome. In: E. Indriati (ed.) Recent Advances on Southeast Asian palaeoanthropology and archaeology, Int. Seminar on Southeast Asian Paleoanthropology, Yogyakarta, p. 160-168.

Bouteaux, A., A.M. Moigne, F. Semah & T. Jacob (2007)- Les assemblages fauniques associes aux sites a *Homo erectus* du dome de Sangiran (Pleistocene moyen, Java, Indonesie). Comptes Rendus Palevol 6, 3, p. 169-179.
(online at: https://www.sciencedirect.com/science/article/pii/S1631068306001485)
('The faunal assemblages associated with Homo erectus sites at Sangiran (M Pleistocene, Java)'. Homo erectus in fluvial deposits outcropping in several localities. Thirteen taxa of M Pleistocene mammals determined. Lithic tools rare at these sites. Mechanical action of water responsible for accumulations)

Bouteaux, A., A.M. Moigne & K. Setiagama (2008)- Etudes archeozoologiques de sites javanais du Pleistocene: les sites de plein air du dome de Sangiran (Java central) et le site en grotte de Song Terus (Java est). In: Archaeozoology of the Near East VIII, Travaux Maison de l'Orient et de la Mediterranee, 49, p. 79-97.
(online at: www.persee.fr/doc/mom_1955-4982_2008_act_49_1_2702)
('Archeozoologic studies at Pleistocene sites of Java: Sangiran Dome (C Java) and Song Terus cava (E Java)')

Braches, F. & R. Shutler (1984)- The Philippines and Pleistocene dispersal of mammals in island Southeast Asia. Philippine Quarterly Culture and Society 12, 2, p. 106-115.
(Presence of Pleistocene mammalian faunas on Luzon and Mindanao led Von Koenigswald (1935) to propose Pleistocene migration route from mainland SE Asia to Borneo/ Java through Philippines. However, Luzon probably island fauna and Philippines probably played no major role in Pleistocene dispersal of 'Sino-Malayan faunas' to Java)

Brandon-Jones, D. (1998)- Pre-glacial Bornean primate impoverishment and Wallace's Line. In: R. Hall & J.D. Holloway (eds.) Biogeography and geological evolution of SE Asia, Backhuys Publishers, Leiden, p. 393-403.
(online at: http://searg.rhul.ac.uk/searg_uploads/2016/01/Brandon-Jones.pdf)

Brandt, R.W. (1976)- The Hoabinhian of Sumatra: some remarks. Modern Quaternary Research in Southeast Asia 2, p. 49-52.
('Hoabinhian' stone tools from Medan area, N Sumatra. Called 'sumatraliths', made of andesite. Named after E Holocene stone artifact assemblages from N Vietnam)

Brasseur, B. (2009)- Dynamique et histoire des depots du Pleistocene inferieur et moyen ancien du dome de Sangiran (Java central, Indonesie): caracterisation des surfaces d'occupation a *Homo erectus*. Ph.D. Thesis, Museum National Histoire Naturelle, Paris, p. 1-360. *(Unpublished)*
(online at: www.researchgate.net/publication/326655192_Dynamique_et_Histoire_des_depots_du_Plei Etc.)
('Dynamics and history of lower and early M Pleistocene deposits of Sangiran dome (central Java, Indonesia): characterization of Homo erectus occupation layers'. Thick Quaternary sediments of Sangiran dome with oldest human fossils dated to 1.5 Ma, but mainly in Bapang/Kabuh Fm fluvio-volcanic layers dated 1- 0.7 Ma (also with tektite horizon). Ngebung 2 hill with preserved human occupation surface dating to 0.8 Ma. Proposes reconstruction of rivers pathways and development at ~1.0 Ma of alluvial fan coming from N and linked to active tectonic phase. Frequent weathering and reworking of volcanic tuffaceous material may explain wide range of radiometric dates in hominid bearing series. Several mudflows rapidly covered Ngebung 2 H. erectus occupation surface)

Brasseur, B. (2020)- A younger “earliest human migration” to Southeast Asia. New dating suggests that hominids arrived at the Sangiran dome later than had been thought. Science 367, 6474, p. 147-148.
(Comment on Matsu’ura et al. (2020). For 25 years, two competing chronological frameworks for Sangiran sedimentary sequence: (1) “younger” chronology for hominid fossil layers (from 1.3- 1.1 Ma to 0.7- 0.6 Ma); and (2) “older” chronology (1.7- 1.6 Ma until 1 Ma). Matsu’ura et al. (2020) support “younger” model, with new dating of 3 tuffs: below hominid-bearing layers (1.34 Ma), Grenzbank (<0.97 Ma) and lowermost Bapang

Fm (0.88 Ma). Also Perning hominin age rejuvenated to <1.49 Ma. "Younger" chronology for Sangiran hominids implies humans migrated from Asia to SE Asia ~300,000 years later than previous estimates)

Brasseur, B., M.A. Courty, B. Deniaux, N. Fedoroff, B. Poreda & F. Semah (2007)- The geodynamic context of the ca. 0.8 Ma layers in the Sangiran Dome (Central Java. Indonesia): traces of the fall-event linked to the Australasian tektites strewn field? In: N.R. Catto (ed.) 17th INQUA Congress, The tropics: heat engine of the Quaternary, Cairns 2007, Quaternary International 167-168, Supplement, p. *(Abstract only)*

Brasseur, B., F. Semah, A.M. Semah & T. Djubiantono (2011)- Approche paleopedologique de l'environnement des hominides fossiles du dome de Sangiran (Java central, Indonesie). Quaternaire (Paris) 22, 1, p. 3-11.
(online at: https://journals.openedition.org/quaternaire/5815)
('Palaeopedological approach of the environment of fossil hominids from the Sangiran dome (Central Java, Indonesia)'. Study of paleo-soils in Pleistocene of Sangiran Dome area, C Java. Six paleosoil types. First fully terrestrial level identified at base of U Pucangan Fm, corresponding to development of open landscape on earlier sites of wide coastal swamps. Higher up, environments indicative of seasonal climate with long dry season, alternating with periods of more humid palustrine conditions. Recurrent aridity proxies in Grenzbank and Kabuh series (with most commont hominid fossils). Soils reflect long dry season and open vegetation landscape, in agreement with stratigraphical and palynological observations)

Brasseur, B., F. Semah, A.M. Semah & T. Djubiantono (2015)- Pedo-sedimentary dynamics of the Sangiran dome hominid bearing layers (Early to Middle Pleistocene, Central Java, Indonesia): a palaeopedological approach for reconstructing '*Pithecanthropus*' (Javanese *Homo erectus*) palaeoenvironment. Quaternary International 376, p. 84-100.
(online at: https://www.academia.edu/78543820/Pedo_sedimentary_dynamics_of_Sangiran_dome_hominid_bearing_layers_L_M_Pleistocene_Java_central_Indonesia_a_paleopedological_approach_of_Pithec Etc.)
(Paleosols in Pleistocene of Sangiran dome, C Java. Base of Upper Sangiran (= Pucangan) Mb earliest continental deposits with fresh-water molluscs, corresponding to development of open landscape with wide coastal marshes and mangroves, with rain forest cover on hinterland. Higher in stratigraphic succession, seasonal climate with long dry season alternating with periods of more humid palustrine conditions. From U Sangiran Mb to lower Bapang (= Kabuh Fm) Mb, erosion of soil cover caused accumulation of pedosediments in topographic depressions. Recurrent aridity proxies in paleosols of Bapang (= Kabuh) Fm, reflecting long dry season and open vegetation landscape)

Broadfield, D., R.L. Holloway, K. Mowbray, A. Silver & S. Marquez (2001)- The endocast of Sambungmacan 3 (Sm3): a new *Homo erectus* from Java. The Anatomical Record 262, 4, p. 369-379.
(online at: https://onlinelibrary.wiley.com/doi/10.1002/ar.1047)
(New fossil calvaria, Sambungmacan3 (Sm3; see also Marquez et al., 2001) one of most advanced and complete endocasts from Java. Sm3 unique morphology, with features similar to both Indonesian and Chinese Homo erectus, but also significant characters reminiscent of later hominins)

Brongersma, L.D. (1935)- Notes on some recent and fossil cats, chiefly from the Malay archipelago. Zoologische Mededelingen 18, p. 1-90.
(online at: www.repository.naturalis.nl/document/149407)

Brongersma, L.D. (1937)- On fossil remains of a Hyaenid from Java. Zoologische Mededelingen 20, p. 186-202.
(online at: www.repository.naturalis.nl/document/150150)
(On hyaenid fossils from Kedung Brubus, C Java, in Dubois collection)
(Study of Recent cats in Indonesian region and on Pleistocene fossils in Dubois collection from Trinil, C Java)

Brongersma, L.D. (1937)- Notes on fossil and prehistoric remains of Felidae from Java and Sumatra. Comptes Rendus XIIe Congres International de Zoologie, Lisbon 1935, p. 1855-1865.

Brongersma, L.D. (1941)- On the remains of carnivora from cave deposits in Java and Sumatra, with notes on recent specimens, I. Zoologische Mededelingen, 23, p. 114-148.
(online at: www.repository.naturalis.nl/document/149372)
(Descriptions of Late Pleistocene carnivore fossils from C Java caves: dog from Gua Lawa (=Wajak man locality) and a marten/weasel (Martes) from Gua Jimbe)

Brongersma, L.D. (1941)- De verzameling van Indische fossielen (Collctie Dubois). De Indische Gids 63, March 1941, p. 97-116.
('The Indies fossils collection (Collection Dubois)')

Brongersma, L.D. (1958)- On an extinct species of the genus *Varanus* (Reptilia, Sauria) from the island of Flores. Zoologische Mededelingen 36, 7, p. 113-125.
(online at: www.repository.naturalis.nl/document/149846)
(Late Pleistocene(?) lizard Varanus hooijeri n.sp. from cave deposits on Flores, collected by T.L. Verhoeven. Associated with Mesolithic flake and blade industry)

Bronson, B. & T. Asmar (1975)- Prehistoric investigations at Tianko Panjang Cave, Sumatra: an interim report. Asian Perspectives 18, 2, p. 128-145.
(Excavation in 1974 of obsidian artifacts in cave in W Jambi, which was first tested by Zwierzycki in early 1920s)

Brothwell, D.R. (1960)- Upper Pleistocene human skull from Niah caves, Sarawak. Sarawak Museum Journal 9, p. 323-349.

Brown, P. (1992)- Recent human evolution in East Asia and Australasia. Philosophical Transactions Royal Society London, B, 337, p. 235-242.
(online at: www.peterbrown-palaeoanthropology.net/Brown%201992%20235-242.pdf)

Brown, P. (1994)- Cranial vault thickness in Asian *Homo erectus* and modern *Homo sapiens*. In: J.L. Lorenz (ed.) In: J.L. Franzen (ed.) 4th Int. Conference 100 years of *Pithecanthropus*. The *Homo erectus* problem, Frankfurt 1991, Courier Forschungsinstitut Senckenberg 171, p. 33-46.
(Thickened cranial vault bone argued to distinguished Homo erectus from H. sapiens, but considerable overlap with modern Australian aboriginal populations and (Chinese) archaic Homo sapiens)

Brown, P. (2001)- Chinese Middle Pleistocene hominids and modern human origins in East Asia. In: L. Barham & K.R. Brown (eds.) Human roots- Africa and Asia in the Middle Pleistocene. University of Bristol, Western Academic Specialist Press, Chapter 10, p. 135-147.
(online at: www.peterbrown-palaeoanthropology.net/Brown%202001%20Chinese%20mid%20Pleist.pdf)

Brown, P. (2012)- LB1 and LB6 *Homo floresiensis* are not modern human (*Homo sapiens*) cretins. J. Human Evolution 62, p. 201-224.
(online at: https://www.peterbrown-palaeoanthropology.net/Brown%202012.pdf)
(Late Pleistocene Homo floresiensis from Liang Bua cave, Flores, associated with stone artefacts and bones of Stegodon. Recent arguments that characteristics of H. floresiensis consistent with dwarfism and delayed development in modern human (Homo sapiens) cretins deemed invalid: no modern human skeleton known with attributes of H. floresiensis)

Brown, P. & T. Maeda (2009)- Liang Bua *Homo floresiensis* mandibles and mandibular teeth: a contribution to the comparative morphology of a new hominin species. J. Human Evolution 57, 5, p. 571-596.
(online at: https://www.peterbrown-palaeoanthropology.net/Brown%20and%20Maeda%202009.pdf)
(Morphological and metrical comparisons of mandibles of Homo floresiensis from Liang Bua place them outside H. sapiens and H. erectus ranges of variation. Mandibles, cranial and postcranial anatomy, limb proportions and functional anatomy of wrist and shoulder in many respects closer to African early Homo or Australopithecus than to later Homo)

Brown, P., T. Sutikna, M. Morwood, R.P. Soejono, Jatmiko, E.W. Saptomo et al. (2004)- A new small-bodied hominin from the Late Pleistocene of Flores, Indonesia. Nature 431, p. 1055-1061.
(online at: https://www.peterbrown-palaeoanthropology.net/nature02999.pdf)
(Discovery of new, tiny hominid Homo floresiensis from Liang Bua cave on Flores)

Brumm, A., F. Aziz, G.D. van den Bergh, M.J. Morwood, M.W. Moore, I. Kurniawan, D.R. Hobbs & R. Fullagar (2006)- Early stone technology on Flores and its implications for *Homo floresiensis*. Nature 441, 7093, p. 624-628.
(C Flores Soa Basin sites contain stone artefacts associated with Stegodon florensis, Komodo dragon, rat, etc., dated as 840-700 ka. Apparent technological continuity with those excavated from Late Pleistocene at Liang Bua cave, 50 km to W, dated as 95-74 and 12 ka, and associated with small-bodied Homo floresiensis)

Brumm, A., D. Bulbeck, B. Hakim, B. Burhan, A.A. Oktaviana, I. Sumantri, J.X. Zhao, M. Aubert, R. Sardi et al. (2021)- Skeletal remains of a Pleistocene modern human (*Homo sapiens*) from Sulawesi. PloS One 16, 9, e0257273, p. 1-27.
(online at: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0257273)
(First Pleistocene human skeletal remains from Sulawesi, at Leang Bulu Bettu, SW Sulawesi. Dated to 25-16 ka)

Brumm, A., B. Hakim, M. Ramli, M. Aubert, G.D. van den Bergh, Bo Li, B. Burhan + 22 others! (2018)- A reassessment of the early archaeological record at Leang Burung 2, a Late Pleistocene rock-shelter site on the Indonesian island of Sulawesi. PloS One 13, 4, e0193025, p. 1-43.
(online at: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0193025)
(Leang Burung 2 limestone rock shelter early human occupation site in Late Pleistocene of Maros karst area of S Sulawesi. New stratigraphic information collected in 2007 and 2011-2013)

Brumm, A., G.M. Jensen, G.D. van den Bergh, M.J. Morwood, I. Kurniawan, F. Aziz & M. Storey (2010)- Hominins on Flores, Indonesia, by one million years ago. Nature 464, 7289, p. 748-752.
(online at: www.academia.edu/10968013/Hominins_on_Flores_Indonesia_by_one_million_years_ago)
(Wolo Sege, a new site in Soa Basin, with in- situ stone artefacts stratigraphically below previously discovered Mata Menge site. Ignimbrite overlying artefact layers erupted 1.02 Ma, providing new minimum age for hominins on Flores and predates disappearance from Soa Basin of 'pygmy' Stegodon and giant Geochelone)

Brumm, A., I. Kurniawan, M.W. Moore, Suyono, R. Setiawan, Jatmiko, M.J. Morwood & F. Aziz (2009)- Early Pleistocene stone technology at Mata Menge, Central Flores, Indonesia. In: F. Aziz et al. (eds.) Pleistocene geology, palaeontology and archaeology of the Pleistocene Soa Basin, Central Flores, Indonesia, Chapter 4, Pusat Survei Geologi, Bandung, Special Publ. 36, p. 119-137.
(E Pleistocene stone tool assemblage from Mata Menge in Soa Basin, Flores, is oldest Palaeolithic stone assemblage in well-dated stratigraphic context in SE Asia. 91% of 459 artefacts made from volcanic rocks)

Brumm, A., M.C. Langley, M.W. Moore, B. Hakim, M. Ramli, I. Sumantri, B. Burhan, A.M. Saiful, L. Siagian, Suryatman et al. (2017)- Early human symbolic behavior in the Late Pleistocene of Wallacea. Proc. National Academy of Sciences USA (PNAS) 114, 16, p. 4105-4110.
(onlne at: https://www.pnas.org/doi/10.1073/pnas.1619013114)
(Leang Bulu Bettue cave and rock-shelter in SW Sulawesi with relicts of Late Pleistocene (~30-22 ka) Homo sapiens. Include previously unknown items of personal ornamentation, portable art, etc., fashioned from body parts of endemic animals)

Brumm, A. & M.W. Moore (2012)- Biface distributions and the Movius Line: a Southeast Asian perspective. Australian Archaeology 74, p. 32-46.
(online at: http://ro.uow.edu.au/cgi/viewcontent.cgi?article=7783&context=scipapers)
(Movius Line suggests that 'true' Acheulean biface stone tools, especially handaxes, are common only in Africa and W Eurasia, but bifaces relatively common in SE Asia (e.g. Pacitan/Java, Walanae River/Sulawesi, etc.))

Brumm, A, M.W. Moore, G.D. van den Bergh, I. Kurniawan, M.J. Morwood & F. Aziz (2010)- Stone technology at the Middle Pleistocene site of Mata Menge, Flores, Indonesia. J. Archaeological Science 37, p. 451-473.
(Ola Bula Fm (1.0- 0.65 Ma) at Mata Menge on Flores, yielded largest stone artefact assemblage from bedded and securely dated M Pleistocene in SE Asia. Stone flaking technology described here for first time. Reduction sequence based on the centripetal or 'radial' reduction of transported blanks)

Brumm, A., G.D. van den Bergh, M. Storey, I. Kurniawan, B.V. Alloway, R. Setiawan, E. Setiyabudi, R. Grun et al. (2016)- Age and context of the oldest known hominin fossils from Flores. Nature 534, 7606, p. 249-253.
(Excavations in fluvial valley-fill sandstone at M Pleistocene (0.7 Ma) Mata Menge site, Soa Basin, C Flores, yielded hominin fossils ancestral to Late Pleistocene Homo floresiensis. Hominins inhabited savannah-like open grassland habitat with wetland component, in relatively dry climate, Hominin fossils occur alongside remains of insular fauna (Stegodon florensis, giant rat Hooijeromys nusatenggara, Varanus komodoensis, etc.) and simple stone technology)

Bulbeck, D. (2004)- South Sulawesi in the corridor of island populations along East Asia's Pacific Rim. In: S.G. Keates & J. Pasveer (eds.) Quaternary Research in Indonesia, Chapter 12, Modern Quaternary Research in Southeast Asia 18, Balkema, Leiden, p. 221-258.

Bulbeck, D. & I. Caldwell (2000)- Land of iron. The historical archaeology of Luwu and the Cenrana valley. Results of the origin of complex society in South Sulawesi Project (OXIS). Centre for South-East Asian Studies, University of Hull, p. 1-141.
(online at: www.academia.edu/7476872/Land_of_Iron_The_Historical_Archeology_of_Luwu_and_the_Cenrana_Valley)

Bulbeck, D. & B. Marwick (2022)- Stone industries of mainland and island Southeast Asia. In: C.F.W. Higham & N.C. Kim (eds.) The Oxford Handbook of Early Southeast Asia, Chapter 5, p. 124-C5._125.
(online at: https://faculty.washington.edu/bmarwick/PDFs/Bulbeck-and-Marwick-2021-Oxford.pdf)
(Archaeological sites with concentrations of flaked stones associated with anatomically modern Homo sapiens, are widespread across Southeast Asia. Two main types of lithic industries: (1) core-based (knapping raw stone to produce cores, used for detaching flakes); and (2) cobble-based (flaking surface of cobbles). Quality of locally available stone underpinned distinction. Dominance of core-based industries in Island SE Asia and cobble-based industries in Mainland SEt Asia (with exceptions such as cobble-based industries in N Sumatra, Niah Caves (Sarawak), and N Moluccas))

Bulbeck, D., M. Pasqua & A. di Lello (2000)- Culture history of the Toalean of South Sulawesi, Indonesia. Asian Perspectives (University of Hawaii Press) 39, 1-2, p. 71-108.
(online at: https://scholarspace.manoa.hawaii.edu/bitstream/10125/17135/1/AP-v39n1-2-71-108.pdf)
(Review of the 'Toalean': microlitihic stone tool assemblages from cave/ rock shelter deposits fills in Miocene limestones in SW Sulawesi, with age ranges between ~8000 and 1500 B.P. First excavated and named in 1902 by P & F. Sarasin. Later work by P. van Stein Callenfels in 1930s, R. van Heekeren in 1940s and Indonesian-Australian expeditions since 1969. Etc.etc.)

Bulbeck, D., I. Sumantri & P. Hiscock (2004)- Leang Sakapao 1, a second dated Pleistocene site from South Sulawesi, Indonesia. In: S.G. Keates & J. Pasveer (eds.) Quaternary Research in Indonesia, Chapter 8, Modern Quaternary Research in Southeast Asia 18, p. 111-128.
(Rock shelter at base of limestone cliff in SW Sulawesi with evidence of Late Pleistocene (~31-20 ka) human habitation (stone artefacts, pottery, etc.)))

Burhan, B., B. Hakim, I. Sumantri, A.M. Saiful, A.A. Oktaviana, R. Sardi, Hasliana, M. Ramli, L. Siagian et al. (2025)- A near-continuous archaeological record of Pleistocene human occupation at Leang Bulu Bettue, Sulawesi, Indonesia. PLoS One 20, 12, e0337993, p. 1-51.
(online at: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0337993)

(Well-illustrated review of deep-trench excavations at Leang Bulu Bettue limestone cave/ rock-shelter complex in Maros-Pangkep karst region of S Sulawesi .With two phases of M? Pleistocene to late Holocene archeological record: Phase I, with cobble-based core and flake stone artefact production (> 132 ka). fauna dominated by extant dwarf bovids (Bubalus sp., anoas) and now-extinct proboscideans, and evidence for animal butchery. Followed around 40 ka by Phase II, with distinct archaeological signature, including first evidence for burning, artistic expression and symbolic culture. Possibly reflects replacement of archaic hominins by modern humans)

Busk, G. (1869)- Notice of the discovery at Sarawak in Borneo of the fossilized teeth of *Rhinoceros* and of a Cervine ruminant. Proc. Zoological Society of London 37, 1, p. 409-416.

Chia, S., L. Yondri & T. Simanjuntak (2007)- The origins of the obsidian artifacts from Gua Pawon, Dago and Bukit Karsamanik in Bandung, Indonesia. Amerta 25, 1, p. 48-57.
(online at: https://jurnalarkeologi.kemdikbud.go.id/index.php/amerta/article/view/680)
(Obsidian artifacts found in Gua Pawon, Dago and Bukit Karsamanik in Bandung came from well-known sources of Gunung Kendan in Nagreg, Kampung Rejeng in Garut or elsewhere (age of tools ~7-10 ka))

Chia, S., L. Yondri & T. Simanjuntak (2011)- Obsidian sourcing in Bandung, Indonesia. Asian Perspectives 49, 1, p. 148-156.
(online at: https://scholarspace.manoa.hawaii.edu/bitstream/10125/29094/1/AP_V49No1_chia.pdf)
(same paper as above)

Choi, K. & D. Driwantoro (2007)- Shell tool use by early members of *Homo erectus* in Sangiran, central Java, Indonesia: cut mark evidence. J. Archaeological Science 34, 1, p. 48-58.
(1.6- 1.5 Ma old cut marks on Pleistocene bovid bones from Black Clay in Pucangan Fm in Sangiran, inflicted by thick clamshell flakes (presumably during butchery by Homo erectus). This documents first use of tools in Sangiran and oldest evidence of shell tool use in world)

Ciochon, R.L. (2010)- Divorcing hominins from the *Stegodon-Ailuropoda* fauna: new views on the antiquity of hominins in Asia. In: J.G. Fleagle et al. (eds.) Out of Africa I: The first hominin colonization of Eurasia, Chapter 8, Springer, p. 111-126.
(online at: www.researchgate.net/publication/226797194_Divorcing_Hominins_from_the_Stegodon-Ail Etc.)
(Pleistocene Stegodon-Ailuropoda (= Panda) fauna of S China and peninsular SE Asia contains ape species previously attributed to early hominins, but no clear evidence. Early hominins may have inhabited parts of S China without forest, but not with heavily forested, humid-climate adapted Stegodon-Ailuropoda mammalian fauna. Homo erectus likely arrived in Java between 1.8-1.6 Ma, but at ~900 ka hominins and most other contemporary large mammals seem to have left area)

Ciochon, R.L. & O.F. Huffman (2018)- Java Man. In: S. Smith (ed.) Encyclopedia of Global Archaeology, Springer International, p. 1-8.
(online at: https://www.researchgate.net/publication/326681417_Java_man)
(Review of 'Java Man', the popular name for Pleistocene Homo erectus of Java. H. erectus widely distributed in Old World, but in SE Asia limited to 100 skeletal specimens from C and E Java. Discussions of Trinil, Sangiran and Ngandong and Mojokerto hominid sites)

Ciochon, R.L., F. Huffman, E.A. Bettis, Y. Zaim, Y. Rizal & Aswan (2009)- Rediscovery of the *Homo erectus* bed at Ngandong: Site formation of a late Pleistocene hominin site in Asia. American J. Physical Anthropology, Supplement 48, p. 110. *(Abstract only?)*

Clarkson, C, S. Haberle & S. O'Connor (2023)- 40,000 years of technological continuity and change at Madja Kuru 2 Timor-Leste. Quaternary Science Reviews 320, 108340, p. 1-15.
(online at: https://www.sciencedirect.com/science/article/pii/S0277379123003888)

(Timor-Leste known for long-term continuity in Homo sapiens lithic manufacturing techniques that span the last 44,000 years. Confirmation of this pattern at Matja Kuru 2 limestone cave in NE-most Timor Leste close to Lake Ira Lalaro. With earliest known evidence for obsidian and possible inter-island transport in Wallacea)

Cooper, A., Stringer. C.B. (2013)- Did the Denisovans cross Wallace's Line? Science 342, 6156, p. 321-323.
(online at: https://www.researchgate.net/publication/258056301_Did_the_Denisovans_Cross_Wallace's_Line)
(DNA data suggest highest % of Denisovan DNA in modern human populations are in New Guinea and Wallacea (E Indonesia) and in Australian Aborigines, suggesting interbreeding of indigenous? Denisovan humans (who may have arrived after 600,000 years ago) with later-arriving Homo sapiens)

Corlett, R.T. (2010)- Megafaunal extinctions and their consequences in the tropical Indo-Pacific. In: S.G. Haberle et al. (eds.) Altered ecologies: fire, climate and human influence on terrestrial landscapes, Terra Australis 32, ANU Press, Chapter 8, p. 117-131.
(online at: www.jstor.org/stable/pdf/j.ctt24h8rj.10.pdf)
(Global Quaternary Megafauna Extinction (QME) event eliminated 2/3 of all mammal genera, with most well-dated extinctions occurring between ~50-30 ka. Java probably had fully modern fauna by 120 ka. In Indo-Pacific, hominin impacts probably major factor behind most large vertebrate extinctions and range restrictions in the past 130 kyrs and probably earlier ones)

Corvinus, G. (2003)- *Homo erectus* in East and Southeast Asia, and the questions of the age of the species and its association with stone artifacts, with special attention to handaxe-like tools. Quaternary International 117, 1, p. 141-151.
(Many fossil remains of H. erectus found In C Java, but not sure which tools belonged to H. erectus. Sangiran and Ngandong industries of small flakes provisionally connected with H. erectus soloensis. Handaxe-like tools from Pacitan, Java and Cabenge, Sulawesi are of uppermost Pleistocene age and work of modern humans)

Cosijn, J. (1931)- Voorloopige mededeeling omtrent het voorkomen van fossiele beenderen in het heuvelterrein ten Noorden van Djetis en Perning (Midden Java). Verhandelingen Geologisch-Mijnbouwkundig Genootschap Nederland Kolonien, Geologische Serie 9, 2, p. 113-119.
('Preliminary communication on the occurrence of fossil bones in the hill country N of Jetis and Perning, C Java'. Localities N of Mojokerto. Bone-bearing layers in folded volcanoclastic sediments with Stegodon, Cervus lydekkeri, Sus, Cervus, Crocodilus, etc.Underlying marine claystones with corals and molluscs. Bone beds deemed similar to Trinil beds, and considered as in Pliocene age (now known as 'Djetis fauna', always viewed by Von Koenigswald as one of the older mammal assemblages on Java, older than Trinil, and of Early Pleistocene age; JTvG))

Cosijn, J. (1932)- Tweede mededeeling over het voorkomen van fossiele beenderen in het heuvelland ten Noorden van Djetis en Perning (Java). Verhandelingen Geologisch-Mijnbouwkundig Genootschap Nederland Kolonien, Geologische Serie 9, 3, p. 135-148.
('Second communication on the occurrence of fossil bones in the hill country N of Jetis and Perning, C Java'. More details on fossil mammal localities. No paleontology (latest Pliocene molluscs summarized in Martin (1932))

Covert, H.H., M.W. Hamrick, T. Dzanh & K.C. McKinney (2001)- Fossil mammals from the Late Miocene of Vietnam. J. Vertebrate Palaeontology 21, p. 633-636.

Croft, D.A., L.R. Heaney, J. Flynn & A.P. Bautista (2006)- Fossil remains of a new, diminutive *Bubalus* (Artiodactyla: Bovidae: Bovini) from Cebu Island (Philippines). J. Mammalogy 87, 5, p. 1037-1051.
(online at: https://academic.oup.com/jmammal/article/87/5/1037/894807)
(Small Pleistocene- Holocene Bubalus (Bubalus cebuensis, sp. nov .) likely result of island dwarfing. First fossil mammal reported from Cebu Island and only non-proboscidean from Negros-Panay Philippine Faunal Region)

Curnoe, D., I. Datan, J.X. Zhao, C. Leh Moi Ung, M. Aubert, M.S. Sauffi, G.H. Mei, R. Mendoza & P.S.C. Tacon (2018)- Rare Late Pleistocene-early Holocene human mandibles from the Niah Caves (Sarawak, Borneo). PloS One 13, 6, e0196633, p. 1-17.
(online at: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0196633)
(Uranium-series dating and descriptions of three partial human mandibles from Niah Caves recovered during excavations by Harrisson in 1957: ~30-28 ka, ~11.0-10.5 ka and ~10.0-9.0 ka (= minimum ages))

Curnoe, D., I. Datan, P.S.C. Tacon, C.L.M. Ung & M.S. Sauffi (2016)- Deep skull from Niah Cave and the Pleistocene peopling of Southeast Asia. Frontiers in Ecology and Evolution 4, 75, p. 1-17.
(online at: http://journal.frontiersin.org/article/10.3389/fevo.2016.00075/full)
(Late Pleistocene Deep Skull from Niah Cave in Sarawak the oldest (>50 ka) anatomically modern human from island SE Asia)

Dammerman, K.W. (1934)- On prehistoric mammals from the Sampoeng cave, Central Java. Treubia 14, p. 477-486.
(online at: https://e-journal.biologi.lipi.go.id/index.php/treubia/article/view/2517/2159)
(Diverse fauna found in Sampoeng cave/rock shelter E of Ponorogo, C Java (first described by Van Es (1929) and more systematically excavated by P.V. van Stein Callenfels around 1930). Associated with Melanesian-like human remains and stone tools (N.B.: Erdbrink (1954) noted that Dammermann failed to report the common Varanus remains from this site, which he believed were an important part of the cave dwellers' diet))

Danisworo, C. (1987)- Lithostratigraphy and magnetostratigraphy of the Quaternary deposits in the Sangiran area, Central Java, Indonesia. Doct. Thesis, Vrije Universiteit, Brussel, p. *(Unpublished)*

Danisworo, C. (1992)- Magnetostratigraphy of Plio-Pleistocene deposits in the Sangiran area, Central Java. Proc. 21st Annual Meeting Indonesian Association Geologists (IAGI), Yogyakarta, 2, p. 477-485.
(online at: https://www.iagi.or.id/web/digital/50/21th-Vol-II-(7-10-Des-1992)-107-116.pdf)
(Normal and reversed magnetic polarities identified in Late Pliocene- Pleistocene in Sangiran Dome. Gauss-Matuyama boundary (2.48 Ma) just above Balanus Limestone of Puren Fm (U Kalibeng Fm). Normal polarity in Cemoro (Pucangan) Fm claystone probaly Olduvai event (~1.87-1.67 Ma). Base Jaramillo event (0.97 Ma) below 'Grenzbank' at base Bapang (Kabuh) Fm. Brunhes- Matuyama reverse to normal transition (0.73 Ma) between TB3 and TB4 tuff horizons in Bapang (Kabuh) Fm (see also Hyodo et al. 1988, 2011, Sunardi 2010))

Danisworo, C. (2001)- Stratigraphic position (in the Quaternary stratigraphy) and the age of *Pithecanthropus erectus* VIII discovered in the Sangiran area, Central Java. Proc. 30th Annual Conv. Indonesian Association Geologists (IAGI) and 10th Reg. GEOSEA Congress, Yogyakarta, Majalah Geologi Indonesia (IAGI) 16, Special Edition, p. 131-139.
(Pithecanthropus erectus VIII skull considered to be from within Jagan Tuff Member, lower Bapang Fm., between 0.97- 0.90 Ma old, or in Pleistocene Jaramillo magnetic event)

Das, I. (1997)- A reassessment of *Hardella isoclina* Dubois, 1908 (Testudines: Bataguridae) from the Trinil beds of the Javan Pleistocene. Herpetological Journal 7, 2, p. 71-73.
(online at: https://www.thebhs.org/publications/the-herpetological-journal/volume-7-number-2-april- Etc.)
(Fossil turtle excavated from Trinil, C Java, and described by E. Dubois in 1908; holotype and sole example of this fossil turtle is at the Leiden Nationaal Natuurhistorisch Museum (NNM; now Naturalis). Here attibuted to genus Mauremys (called Duboisemys isoclina (Dubois, 1908) by Karl et al., (2019) and viewed as aquatic river turtle)

Dawson, M.R. (1971)- Fossil mammals of Java I. Notes on Quaternary Leporidae (Mammalia, Lagomorpha) from Central Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B74, p. 27-32.
(On fossil rabbits/ hares from Pleistocene of Sangiran, C. Java)

D'Cunha, M.G.T., A. Montenegro & J.S. Field (2021)- Modeling water crossings leading to the arrival of early Homo in Sulawesi, Indonesia, via paleoclimate drift experiments. J. Archaeological Science: Reports 40, A, 103194, p. 1-12.
(Lithic tools found at Talepu, Sulawesi, in Wallacea suggest that unidenfied hominins (but predating Homo sapiens; Denisovans?) occupied Sulawesi between ~ 118-194 ka. Borneo most likely departure point. Mindanao departures more likely to arrive in Sulawesi than those from Banda Arc)

De Almeida, A. & G. Zbyszewski (1967)- A contribution to the study of the prehistory of Portuguese Timor lithic industries. In: W. G. Solheim (ed.) Archaeology at the Eleventh Pacific Science Congress. (Asian and Pacific Archaeology, Series 1, Honolulu), p. 55-67).
(One of first records of human stone artifacts from E Timor, from Lene Hara cave. Made from jasper and chert (later more systematic excavations at this locality reported by O'Connor et al. 2002))

De Bruyn, M., B. Stelbrink, R.J. Morley, R. Hall, G.R. Carvalho, C.H. Cannon, G. van den Bergh, E. Meijaard, I. Metcalfe, L. Boitani, L. Maiorano, R. Shoup & T. von Rintelen (2014)- Borneo and Indochina are major evolutionary hotspots for Southeast Asian biodiversity. Systematic Biology 63, 6, p. 879-901
(manuscript online at: https://ro.uow.edu.au/articles/journal_contribution/Borneo_and_Indochina_are_major_evolutionary_hotspots_for_Southeast_Asian_biodiversity/27787422?file=50557956)
(SE Asia (in particular Borneo and Indochina) major 'evolutionary hotspots' for diverse range of fauna- flora. Most of SE Asia's biodiversity result of accumulation of immigrants and in situ diversification. Colonization events comparatively rare from younger emergent islands like Java, which show increased immigration events)

Delfino, M. & J. De Vos (2010)- A revision of the Dubois crocodylians, *Gavialis bengawanicus* and *Crocodylus ossifragus*, from the Pleistocene *Homo erectus* beds of Java. J. Vertebrate Paleontology 30, p. 427-441.
(online at: https://www.researchgate.net/publication/232667880_A_Revision_of_the_Dubois_Crocodylians_Gavialis_Bengawanicus_and_Crocodylus_Ossifragus_from_the_Pleistocene_Homo_Erectus_Beds_of_Java)
(Revision of two extinct Javanese crocodylian species Gavialis bengawanicus Dubois 1908 and Crocodylus ossifragus Dubois 1908 (= C. siamensis Schneider 1801). Both found in Trinil, with Stegodon- Homo erectus fauna, which is considered to be result of E Pleistocene dispersal from Siwaliks Hills via Siva-Malayan route)

Delfino, M. & J. De Vos (2014)- A giant crocodile in the Dubois Collection from the Pleistocene of Kali Gedeh (Java). Integrative Zoology 9, 2, p. 141-147.
(Unpublished crocodylian specimen collected by Dubois in latest E Pleistocene of Kali Gedeh tentatively referred to genus Crocodylus. ~1m long lower jaw indicate total length of ~6-7m)

Delson, E., K. Harvati, D. Reddy, L.F. Marcus, K. Mowbray, G.J. Sawyer, T. Jacob & S. Marquez (2001)- The Sambungmacan 3 *Homo erectus* calvaria: a comparative morphometric and morphological analysis. Anatomical Record 262, 4, p. 380-397.
(online at: https://onlinelibrary.wiley.com/doi/10.1002/ar.1048)
(Sambungmacan (Sm) 3 calvaria, discovered on Java in 1977, presumably dredged from bank of Solo River near Sambungmacan. It was illegally removed from Indonesia in 1998 and appeared in New York City natural history shop in early 1999, now at Gadjah Mada University. Age unknown, but Sm 3 skull features intermediate between classic Homo erectus and more modern humans, with greatest similarity to specimens from Ngandong)

De Lumley, H., F. Semah & T. Simanjuntak (1993)- Les outils du Pithecanthrope. Les dossiers d'Archeologie 184, p. 62-68.
('The tools of Pithecanthropus')

De Neve, G.A. (1949)- Note on the occurrence of the remains of a Cetacean in the alluvial tin deposits of the island of Billiton. Chronica Naturae 105, 4, p. 118-120.
(online at: http://colonial.library.leiden.edu/....)
(Vertebra of fossil whale in Quaternary tin ore-bearing deposits near Manggar, Belitung. Related to fin whale Balaenoptera)

De Neve, G.A. (1949)- Opmerking over de fossiele wervel gevonden in Billiton. Chronica Naturae 105, 11, p. 293-294.
('Remark on the fossil vertebra found on Billiton'. Second whale vertebra fossil from Manggar, Belitung, probably from (Quaternary) marine shell layer first identified by Verbeek at ~2m(?) below sea level)

De Neve, G.A. (1950)- Notitie over vertebraat-fragmenten uit de proefboring Kebajoran. De Ingenieur in Indonesia 1950, 5, p. IV.23-IV.25.
(online at: https://colonialarchitecture.eu/islandora/object/uuid%3A6cdd1184-7f8c-4d56-8823-3cc86d06abef/datastream/PDF/view)
('Note on vertebrate fragments from the test well Kebajoran'. Vertebrate remains from 51,50 m. depth in water well drilled by the Geological Survey Water Supply department) near Kampong Kebajoran, SW of Jakarta, include jaw fragment of Sus brachygnatus with two molars, and indicate M-Pleistocene age of river deposit)

Deninger, K. (1910)- Uber einen Affenkiefer aus den Kendeng-Schichten von Java. Centralblatt Mineralogie Geologie Palaontologie 1910, p. 1-3.
(online at: www.biodiversitylibrary.org/item/192869page/25/mode/1up)
('On a monkey jaw from the Kendeng beds of Java'. Pleistocene fossil jaw with molars collected by Elbert near Saradan, Madiun District (younger than Trinil bone bed). Assigned to Inuus nemestrinus (macaque family))

Dennell, R.W. (2004)- Hominid dispersals and Asian biogeography during the Lower and early Middle Pleistocene, c. 2.0- 0.5 Mya. Asian Perspectives 43, 2, p. 205-226.

Dennell, R.W. (2005)- The Solo (Ngandong) *Homo erectus* assemblage: a taphonomic assessment. Archaeology in Oceania 40, 3, p. 81-90.
(Homo erectus site near base of Solo River terrace deposits at Ngandong, excavated by Oppenoorth of Geological Survey in 1931-1933, differs from other sites with hominin remains in fluvial deposits, because 12 crania are present, but few other skeletal elements (fluvial disarticulation, large carnivores or headhunters?). Most of >25,000 mammalian fossils excavated from Ngandong and stored in Bandung now lost)

Dennell, R.W. (2009)- The Palaeolithic settlement of Asia. Cambridge University Press, p. 1-548.

Dennell, R.W. (2014)- Hallam Movius, Helmut de Terra, and the line that never was; Burma 1938. In: K. Boyle et al. (eds.) Living in the landscape: essays in honour of Graeme Barker, McDonald Institute Archaeological Research, Cambridge, p. 11-34.
(online at: www.researchgate.net/publication/303157763_Hallam_Movius_Helmut_de_Terra_and_the Etc.)
(About the 'Movius Line', a supposed Early Paleolithic paleobiogeographic dividing line)

Dennell, R.W., J. Louys, H.J. O'Regan & D.M. Wilkinson (2014)- The origins and persistence of *Homo floresiensis* on Flores: biogeographical and ecological perspectives. Quaternary Science Reviews 96, p. 98-107.
(online at: www.academia.edu/11413646/The_origins_and_persistence_of_Homo_floresiensis_on_Flor Etc.)
(Stone artifacts suggest hominids arrived on Flores before 1 Ma and small hominin species (Homo floresiensis) lived on Flores in Late Pleistocene. Flores was always island, at least 19 km from other islands on Sunda Shelf, suggesting early hominids reaching Flores were capable of using watercraft)

Dennell, R.W. & M.D. Petraglia (2012)- The dispersal of *Homo sapiens* across southern Asia: how early, how often, how complex? Quaternary Science Reviews 47, p. 15-22.
(Timing and the paths of colonization of S Asia by Homo sapiens poorly known. Dispersal from E Africa between 60- 40 ka, but U Pleistocene population history of S Asia likely complex)

Dennell, R.W. & W. Roebroeks (2005)- An Asian perspective on early human dispersal from Africa. Nature 438, 7071, p. 1099-1104.

De Terra, H. (1943)- Pleistocene geology and early man in Java. In: Research on early man in Burma, Transactions American Philosophical Society, N.S. 32, 3, p. 437-464.
(Review of work by Dubois, Von Koenigswald, etc. on Pleistocene stratigraphy and hominid sites of Java, incl. Mojokerto, Sangiran, Trinil, Ngandong, Pacitan. Includes study of Solo River Quaternary terraces from observations during 1938 fieldtrip with Teilhard de Chardin, Movius and Von Koenigswald)

De Terra, H. (1949)- Geology and climate as factors of human evolution in Asia. In: W.W. Howell (ed.) Early man in the Far East, Symposium, American Assoc. Physical Anthropologists, Chicago 1946, p. 7-15.

Detroit, F. (2000)- The period of transition between *Homo erectus* and *Homo sapiens* in East and Southeast Asia: new perspectives by the way of geometric morphometrics. In: W. Dong (ed.) Proc. 1999 Beijing Int. Symposium on paleoanthropology, Acta Anthropologica Sinica 19 (Suppl.), p. 75-81.
(Comparison of morphometric components of human skulls from ~2.5 Ma- present. Always clear distinction between H. erectus and H. sapiens cranial architecture. Ngandong specimens (= 'H. soloensis') sometimes considered archaic Homo sapiens, but typical Homo erectus architecture)

Detroit, F. (2002)- Origin et evolution des *Homo sapiens* en Asie du Sud-Est: descriptions et analyses morphometriques de nouveaux fossiles. Doct. Thesis Museum National Histoire Naturelle, Paris, p. 1-445.
(online at: http://hopsea.mnhn.fr/pc/thesis/Detroit_2002_PhD_thesis.pdf)
(' Origin and evolution of Homo sapiens in SE Asia: overview and morphometric analyses of new fossils'. Study of U Pleistocene- Holocene Homo sapiens populations from Malaysia, Java (Wajak, Tulungagung, Gua Lawa, Gunung Sewu, etc.), Flores, Thailand (Moh Khiew), Palawan (Tabon). Two periods in Pleistocene evolution of Homo: (1) time of Homo erectus, marked by some endemism in SE Asia archipelago, (2) intensified migrations with first arrival of H. sapiens in M Pleistocene)

Detroit, F. (2006)- *Homo sapiens* in Southeast Asian archipelagos: the Holocene fossil evidence with special reference to funerary practices in East Java. In: H.T. Simanjuntak et al. (eds.) Proc. Symposium Austronesian diaspora and the ethnogeneses of people in Indonesian archipelago, Solo, Indonesian Inst. Science (LIPI), Jakarta, p. 186-204.
(On Homo sapiens fossils from Gunung Sewu area, Southern Mountains of Java (Song Terus, Song Keplek, Goa Braholo), and their funeral practices)

Detroit, F., J. Corny, E.Z. Dizon & A.S. Mijares (2013)- ""Small Size" in the Philippine human fossil record: Is it meaningful for a better understanding of the evolutionary history of the Negritos?. Human Biology 85, 1 (Special Issue on revisiting the “Negrito” hypothesis), 3, p. 45-63.
(online at: http://digitalcommons.wayne.edu/humbiol/vol85/iss1/3)

Detroit, F., A.S. Mijares, J. Corny, G. Daver, C. Zanolli, E. Dizon, E. Robles, R. Grun & P.J. Piper (2019)- A new species of *Homo* from the Late Pleistocene of the Philippines. Nature 568, 7751, p. 181-186.
(ms online at: https://hal.archives-ouvertes.fr/hal-02296712/file/Detroit_&_al_2019_Nature_postprint.pdf)
(Hominin foot bone from Callao Cave (N Luzon) dated to 67 ka, providing earliest evidence of human presence in Philippines. Twelve additional hominin elements in same layer combination of primitive and derived morphological features, different from other species and warrants attribution to new species Homo luzonensis)

De Visser, J.A. (2008)- The extinct genus *Hexaprotodon* Falconer & Cautley, 1836) (Mammalia, Artiodactyla, Hippopotamidae) in Asia: paleoecology and taxonomy. Ph.D. Thesis University of Utrecht, p. 1-390.
(online at: https://dspace.library.uu.nl/bitstream/handle/1874/31366/visser.pdf?sequence=2)
(Study of Asian (‘Siwalik’) Pleistocene hippopotamid Hexaprotodon, including H. sivajavanicus Dubois 1908 in Java (Kali Glagah, Kedung Brubus and Ngandong. Two sub-species described from Java (Hippopotamus sivalensis koenigswaldi Hooijer 1942 and H.sivalensis soloensis Hooijer 1950) viewed as younger synonyms and parts of natural variability. Javanese Hexaprorodon tends to be smaller than H. from Siwaliks, India)

De Vos, J. (1983)- The *Pongo* faunas from Java and Sumatra and their significance for biostratigraphical and paleo-ecological interpretations: Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B86, p. 417-425.
(Fossil faunas from Punung, Java, and Sumatran caves. Biostratigraphically intermediate between Ngandong and Wadjak faunas, and both indicative of humid forest climate)

De Vos, J. (1985)- De Collectie Dubois. Cranium 1985, p. 26-32.
(online at: http://natuurtijdschriften.nl/download?type=document&docid=523096)
(Brief review of life of E. Dubois and his Pithecanthropus discoveries on Java. Very little on other fossils in Dubois collection)

De Vos, J. (1985)- Faunal stratigraphy and correlation of the Indonesian hominid sites. In: E. Delson (ed.) Ancestors, the hard evidence, A.R. Liss, New York, p. 215-220.
(Review of succession of Pleistocene mammalian faunas from E Java, as also described by Sondaar (1984). From old to young: Satir, Ci Saat, Trinil HK, Kedung Brubus, Ngandong, Punung and Wajak. Oldest faunas rel. poor island faunas. Kedung Brubus fauna reflects greatest interchange with Asian mainland. Oldest hominids in Sangiran of Trinil HK or Ci Saat age)

De Vos, J. (1994)- *Homo modjokertensis*; vindplaats, ouderdom en fauna. Cranium 11, 2, p. 103-107.
(online at: http://natuurtijdschriften.nl/download?type=document&docid=523348)
('Homo modjokertensis; locality, age and fauna'. Discussion of Swisher et al. (1994), who suggest new 1.81 Ma radiometric age of Homo modjokertensis beds, indicating Homo erectus-like fossils may have originated in Asia, not in Africa as generally assumed. Magnetostratigraphic work, associated mammals and uncertainty about exact level of H. modjokertensis skull suggest Swisher's Ar/Ar age and conclusions may be wrong, and H. mojokertensis probably younger)

De Vos, J. (1995)- The migration of *Homo erectus* and *Homo sapiens* in SE Asia and the Indonesian Archipelago. In: J.R.F. Bower & S. Sartono (eds.) Human evolution in its ecological context, Proc. Pithecanthropus Centennial Congress, Leiden 1993, vol. 1, Evolution and ecology of *Homo erectus*, DSWO Press, p. 239-260.

De Vos, J. (1996)- Faunal turnovers in Java in relation to faunas of the continent. Odontologie, Association for comparative Odontology, 1, p. 32-36.
(Commonly quoted, but hard to find paper)

De Vos, J. (2002)- A century of Dutch paleo-anthropological research in Indonesia. In: H. Vermeulen & J. Kommers (eds.) Tales from academia; history of anthropology in the Netherlands. Niccos, Nijmegen Studies in Development and Cultural Change 40, 2, p. 1095-1116.

De Vos, J. (2004)- The Dubois collection: a new look at an old collection. In: C.F. Winkler Prins & S.K. Donovan (eds.) VII Int. Symposium 'Cultural Heritage in Geosciences, Mining and Metallurgy: Libraries - Archives - Museums, Leiden 2003, Scripta Geologica, Special Issue, 4, p. 267-285.
(online at: www.repository.naturalis.nl/document/148593)
(Description of large Dubois collection in Leiden Naturalis museum and its role in studies of Java Pleistocene stratigraphy, mammal faunas, faunal migrations, hominid evolution, etc.)

De Vos, J. (2009)- Receiving an ancestor in the phylogenetic tree. Neanderthal Man, *Homo erectus* and *Homo floresienses*: l'histoire se repete. J. History of Biology 42, 2, p. 361-379.
(Each of the three above hominid species generated widespread discussions and criticism when they were first introduced. It took/ will take one generation before the meaning of these finds was/ will be accepted)

De Vos, J. (2013)- Mid-Pleistocene of Southern Asia. In: S.A. Elias (ed.) Encyclopedia of Quaternary Science, Elsevier, 4, 2n Edition of 2007 book, p. 651-663.

(online at: https://www.researchgate.net/publication/288160779_VERTEBRATE_RECORDS_Mid-Pleistocene_of_Southern_Asia)
(SE Asia Pleistocene land fossils 3 groups:(1) E-M Pleistocene from open sites of Siwaliks, etc. in India–Pakistan, Irrawadi (Myanmar), Nakorn Sawan (Thailand), Satir, Ci saat, Trinil, Kedung Brubus, and Ngandong from Java: only extinct species, mainly open woodlands, including Homo erectus; (2) M-L Pleistocene records from caves/ fissures of S China, Vietnam, Laos, Thailand, Sumatra, Java and Borneo:most species still extant, with Pongo (orangutan) and probably with late Pleistocene H. sapiens; (3) Late Pleistocene- E Holocene from Wallacea (Sulawesi, Flores, Timor, Sumba), Sangihe and Philippines: unbalanced endemic island faunas, with few genera of small proboscideans, pig, large mammals (incl. giant rats, but no carnivores))

De Vos, J. (2014)- The history of palaeoanthropological research in Asia: reasons and priorities for future cooperation in research and preservation of sites and collections. In: N. Sanz (ed.) Human origin sites and the World Heritage Convention in Asia, UNESCO World Heritage Papers 39, p. 68-82.
(online at: http://unesdoc.unesco.org/images/0022/002291/229174e.pdf)
(Review of history of paleoanthropology in SE Asia and principal localities in Indonesia)

De Vos, J. & F. Aziz (1987)- Note on two upper canines of *Megantereon* sp. (Mammalia, Felidae) from the Pleistocene of Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B90, p. 57-63.
(Teeth of sabre-tooth cat, found by Indonesian-Japanese team in 'Grenzbank' Bed in Sangiran Dome, C Java, and reported as Homotherium zwierzyckii. These canines belong to Megantereon sp.)

De Vos, J. & F. Aziz (1989)- The excavations by Dubois (1891-1900), Selenka (1906-1908) and the Geological Survey by the Indonesian-Japanese Team (1976-1977) at Trinil (Java, Indonesia). J. Anthropological Society of Nippon 97, 3, p. 407-420.
(online at: https://www.jstage.jst.go.jp/article/ase1911/97/3/97_3_407/_pdf)
(Reconstructions of Dubois' (poorly documented) excavations along both sides of the Solo River at Trinil in 1891-1893 (which were continued until 1900 by sergeants Kriele and De Winter after Dubois' return to The Netherlands in 1895), and compared to Selenka Expedition work of 1906-1908. Both teams excavated down to the bottom of the main bone layer ('Hauptknochenschicht'; ~1.0 Ma)))

De Vos, J., F. Aziz & P.Y. Sondaar (1993)- Les faunes quaternaires de Java. In: F. Semah & D. Grimaud-Herve (eds.) Le Pithecanthrope de Java, a la decouverte du chainon manquant, Les dossiers d'Archeologie 184, p. 56-61.
('The Quaternary faunas of Java'. Age of Ngandong hominids (H. soloensis or 'advanced H. erectus) ~80,000 to 250,000 years)

De Vos, J., F. Aziz, E. Setiabudi, G.D. van den Bergh & E.Y. Patriani (2007)- A new vertebrate fossil locality near Sumberdadi, Mojokerto (East Java, Indonesia). Proc. International Senckenberg conference, Late Neogene and Quarternary biodiversity and evolution: regional developments and interregional correlations, Weimar 2006, 2, Courier Forschungsinstitut Senckenberg (CFS) 259, p. 175-180.
(36 vertebrate fossils from new locality in sand quarry near Sumberdadi, ~30 km N of Mojokerto, E Java, incl. Stegodon trigonocephalus cf. ngandongensis, Bibos palaeosondaicus, Axis lydekkeri, Rusa sp. and crocodile remains. Advanced stage of Stegodon suggests late M- Late Pleistocene, comparable to Ngandong fauna)

De Vos, J., F. Aziz, P.Y. Sondaar & G.D. van den Bergh (1995)- *Homo erectus* in S.E. Asia: time space and migration routes, III. Migration routes and evolution. In: J. Gibert et al. (eds.) Proc. Int. Conference The hominids and their environment during the Lower and Middle Pleistocene of Eurasia, Orce 1995, p. 369-381.
(Brief review of Pleistocene mammal fossil occurrences in SE Asia. E-M Pleistocene migration via Siva-Malayan route from Siwaliks via Birma to Java brought in Homo erectus in M Pleistocene. During Late Pleistocene Sunda Shelf became connected with continent, causing migration from China, Vietnam, Cambodia via 'Sino-Malayan' corridor, bringing in Homo sapiens and leading to extinction of Homo erectus)

De Vos, J. & A. Bautista (2001)- An update on the vertebrate fossils from the Philippines. National Museum Papers 11, p. 58-62.

De Vos, J. & A. Bautista (2001)- 2003. Preliminary notes on the vertebrate fossils from the Philippines. Proc. Society of Philippine Archaeologists 1, p. 46-62

De Vos, J., A. Bouteaux & A. Bautista (2007)- The mammalian faunas chronology in island Southeast Asia. In: A.M. Semah & K. Setiagama (eds.) First Islanders; human origins patrimony in Southeast Asia, p. 81-84.
(online at: http://hopsea.mnhn.fr/pc/brochures/2007HOPseaFI.pdf)
(Brief review of Pleistocene mammal fauna evolution of SE Asia. Parts of Java first emerged above sea level at ~1.8 Ma. First mammals to reach Java unbalanced island fauna (Satir Fauna of Bumiayu and Sangiran, with hippos (Hexaptotodon), cervids and mastodon (Sinomastodon bumiajuensis)). At 1.2 Ma better connection with Java, with arrival of more balanced Ci Saat fauna, including tigers, and more diverse Trinil Fauna with Homo erectus no later than 1.0 Ma. Around 0.8 Ma new migration to Java (Kedung Brubus Fauna, with first Elephas). Next fauna is Ngandong Fauna that includes Solo Man, a Homo erectus with larger brain size than older forms, that lived until ~0.2 Ma. Final faunas Punung Fauna (~120ka) and Wajak Fauna, both tropical rainforest assemblages, with Homo sapiens)

De Vos, J., S. Sartono, S. Hardja-Sasmita & P.Y. Sondaar (1982)- The fauna from Trinil, type locality of *Homo erectus*: a reinterpretation. Geologie en Mijnbouw 61, 2, p. 207-211.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0ZlN5anhJdHF5YWs/view)
(Revision of Von Koenigswald's mammal stratigraphy. Fauna at Trinil type locality rel. low in species and is older than 'Jetis-fauna' of Von Koenigswald 1934 (= Kedung Brubus fauna). Many endemic species, suggesting island setting (see also comments of Bartstra (1983) defending Von Koenigswald's work, and discussion of Hooijer and Kurten (1984))

De Vos, J. & P.Y. Sondaar (1982)- The importance of the 'Dubois collection' reconsidered. Modern Quaternary Research in Southeast Asia 7, Balkema, Rotterdam, p. 35-63.
(Review of Dubois collection of Pleistocene mammals from C Java, mainly from excavations at Trinil along Solo River and Kedung Brubus near Madiun in 1890's. Material in Naturalis Museum, Leiden, since. Much of material not collected in stratigraphic context. With locality maps and listings of species)

De Vos, J. & P.Y. Sondaar (1994)- Dating hominid sites in Indonesia. Science 266, p. 1726-1727.
(Question correctness of Swisher et al. (1994) older-than-generally-accepted new radiometric ages of pumice associated with Java hominids at Mojokerto and Sangiran sites of C Java (1.8-1.66 Ma instead of 'conventional' 0.97-0.73 Ma))

De Vos, J., P.Y. Sondaar, G.D. van den Bergh & F. Aziz (1994)- The *Homo* bearing deposits of Java and its ecological context. In: J.L. Franzen (ed.) 4th International Conference 100 years of *Pithecanthropus*. The *Homo erectus* problem, Frankfurt 1991, Courier Forschungsinstitut Senckenberg 171, p. 129-140.
(online at: www.researchgate.net/publication/281085976_The_Homo_bearing_deposits_of_Java_and_i Etc.)
(Discussion of Quaternary faunal succession of Java: (1) Satir (lower Kali Klagah series in Bumiayu; island fauna), (2) Ci Saat (upper Kali Glagah, Sangiran upper Black Clay; 1.2 Ma), (3) Trinil H.K. (1 Ma; with first Homo erectus; large amounts of bovids suggest rel. dry, glacial period), (4) Kedung Brubus Fauna (= upper Bapang Fm at Sangiran, ~0.8 Ma; also rel. dry climate), (5)Ngandong (with Homo erectus soloensis), (6) Punung (Late Pleistocene; common primates indicates humid tropical forest environment; last interglacial?), (7) Wajak (Wajak cave; Holocene with Homo sapiens). All Pleistocene Java hominids are Homo erectus, but with several subspecies)

De Vos, J., L.W. van den Hoek Ostende & G.D. van den Bergh (2007)- Patterns in insular evolution of mammals: a key to island palaeogeography. In: W. Renema (ed.) Biogeography, time, and place: distributions, barriers, and islands, Chapter 10, Springer, p. 315-345.
(online at: https://www.researchgate.net/publication/227064513_Patterns_in_Insular_Evolution_of_Mammals_A_Key_to_Island_Palaeogeography)

(Dwarfism and gigantism in unbalanced faunas on islands attest that only a few non-flying mammals were able to reach island. Pygmy elephants and giant rats evolved in isolation of these insular environments. Includes review of Pleistocene mammal island faunas of SE Asia)

De Vos, J. & T.L. Vu (2001)- First settlements: relations between continental and insular Southeast Asia. In: F. Semah et al. (eds.) Proc. International Symposium Origine des peuplements et chronologie des cultures paleolithiques dans le sud-est asiatique, Paris 1998, 24, p. 225-249.

Di Geronimo, I. & S. Sartono (1990)- Sangiran (Java, Indonesia): Upper Pliocene- Pleistocene molluscan environments. Buletin Jurusan Geologi (ITB) 20, p. 33-35.

DIRSP (Dutch-Indonesian Joint-Research Group on Sedimentology and Paleontology of South Sulawesi) (1995)- The geology and stratigraphy of the vertebrate-bearing deposits in the Sengkang Basin: The terrestrial faunal evolution of South Sulawesi during the Plio-Pleistocene. Geological Research Development Centre (GRDC), Bandung, Special Publ. 18, p. 1-112.

Djubiantono, T. (1986)- Etude sedimentologique et paleomagnetique des derniers depots marins de la depression de Solo (Java, Indonesie). Memoire de D.E.E., Museum National Histoire Naturelle, Paris, p.
('Sedimentological and paleomagnetic study of the last marine deposits of the Solo depression (Java, Indonesia')

Djubiantono, T. (1992)- Les derniers depots marins de la depression de Solo (Java Central, Indonesie): chronostratigraphie et paleogeographie. Doct. Thesis, Museum National Histoire Naturelle, Paris, p. 1-208. *(Unpublished)*
('The final marine deposits of the Solo Depression, Central Java; chonostratigraphy and paleogeography'. Kaliuter region of C Java at S flank of Kendeng Hills with thick regressive marine facies. Marine regression deatd as ~2.4 Ma. Two folding phases in Kendeng zone, one at 0.7 Ma, one before 1 Ma. Paleogeographic evolution of Solo Depression and associated fossiliferous sites)

Djubiantono, T. (1993)- Umur alat batu Kedungcumpleng di daerah Kaliuter, Solo, Jawa Tengah. Proc. 22nd Annual Conv. Indonesian Association Geologists (IAGI), Bandung, 1, p. 292-303.
(online at: https://www.iagi.or.id/web/digital/61/31.pdf)
('Age of the Kedungcumpleng stone tools in the Kaliuter area, Solo, Central Java'. Pleistocene correlations and chronostratigraphy and correlations of sections in Kaliuter area, 30km N of Solo. Kedungcumpleng site in Jaramillo paleomagnetic episode (0.87-0.97 Ma), and considered site with oldest hominid stone tools in Java)

Djubiantono, T. & F. Semah (1991)- Lower Pleistocene marine-continental transitional beds in the Solo depression and their relation to the environment of the Pucangan hominids. In: P. Bellwood (ed.) Indo-Pacific Prehistory 1990, Bull. Indo-Pacific Prehistory Association (IPPA) 11, p. 7-13.
(Online at: http://ejournal.anu.edu.au/index.php/bippa/article/view/595/584)
(Kaliuter River section 10 km N of Sangiran shows transition from Pliocene folded marine Lower KalibengFm marls and Late Pliocene- E Pleistocene Upper Kalibeng Fm regressive series, unconformably overlain by unfolded M-U Pleistocene non-marine series with hominids. 'Grenzbank' is unconformity surface)

Djubiantono, T. & F. Semah (1993)- L'ile de Java et son peuplement. In: F. Semah et al. (eds.) Le Pithecanthrope de Java, Les dossiers d' Archeologie 184, p. 12-19.
('Java island and its colonization by humans'.Grenzbank conglomerate in Sangiran linked to uplift and subsequent erosion of ranges surrounding Solo Depression (Kendeng hills and S Mountains))

Djubiantono, T. & F. Semah (1993)- L'evolution de la region de Solo au Quaternaire. In: Le Pithecanthrope de Java, Les dossiers d'Archeologie 184, p. 46-49.

Djubiantono, T., F. Semah & A.M. Semah (1992)- Chronology and palaeoenvironment of Plio- Pleistocene deposits in the Solo Depression (Central Java): the Kaliuter area and its relations with the ancient Javanese

settlements. Proc. VIth National Archaeological Congress, Association of Indonesian Archaeologists, Malang 1992, II, p. 191-242.
(C Java around Lw-M Pleistocene boundary (1.0- 0.8 Ma) dramatic volcano-tectonic activity, involving uplift of ranges around Solo Depression: Kendeng zone last major phase of folding and probably S Mountains uplift)

Djubiantono, T., F. Semah, A.M. Semah, H. Saleki, C. Falgueres & G. Feraud (1994)- Pertanggalan radiometri pada lapisan pengandung *Homo erectus* di Ngebung (Jawa Tengah, Indonesia) hasil pendahuluan. Proc. 23rd Annual Conv. Indonesian Association Geologists (IAGI), Jakarta, 1, p. 184-191.
('Radiometric dating of the Homo erectus-bearing layer in Ngebung (Central Java, Indonesia), preliminary results'. Ngebung site at NW side of Sangiran Dome with rather variable results from different radiometric dating methods just above 'Grenzbank' layer: Ar/Ar of amphibole from tuff ~811 ± 25 ka; U/Th and ESR much younger)

Downing, K.F. (1995)- A large tropical rat from the late Pliocene Walanae Formation, Sulawesi, Indonesia. In: 55th Annual Meeting Soc. Vertebrate Paleontology, Pittsburgh 1995, J. Vertebrate Paleontology 15, Supplement to No. 3, p. 26A-27A. *(Abstract only)*
(First small mammal fossils recovered from Sulawesi, in Late Pliocene Walanae Fm near Lakibong, SW Sulawesi: lower molars of large rat)

Downing, K.F., G.G. Musser & L.E. Park (1998)- The first fossil record of small mammals from Sulawesi, Indonesia; the large murid, *Paruromys dominator*, from the Late(?) Pliocene Walanae Formation. In: Y. Tomida et al. (eds.) Advances in vertebrate paleontology and geochronology. National Science Museum, Tokyo, Monograph 14, p. 105-121.
(Discovery of first small mammal fossils from Sulawesi: two teeth of large rat species, identified as Paruromys dominator, from Walanae Fm at Lakibong, SW Sulawesi)

Drawhorn, G.M. (2024)- Dubois and beyond: The historical background of cave exploration in Sumatra. In: J. Louys et al. (eds.) Quaternary Palaeontology and Archaeology of Sumatra, Terra Australis 56, ANU Press, Canberra, Chapter 4, p. 77-97.
(online at: https://press-files.anu.edu.au/downloads/press/n12084/pdf/ch04.pdf)
(Historic review of five phases of exploration and human use of caves in Sumatra, including cave excavations by E. Dubois in late 1800s)

Drieenhuizen, C. & F. Sysling (2021)- Java Man and the politics of natural history: an object biography. Bijdragen Taal Land en Volkenkunde (J. Humanities Social Sciences SE Asia and Oceania) 177, 2-3, p. 290-311.
(online at: https://brill.com/view/journals/bki/177/2-3/article-p290_9.xml?language=en)
(Review of hominid artifacts of Java Man from Indonesia in Dutch museums)

Dubois, E. (1888)- Over de wenschelijkheid van een onderzoek naar de diluviale fauna van Ned. Indie, in het bijzonder van Sumatra. Natuurkundig Tijdschrift voor Nederlandsch-Indie 48, p. 148-165.
('On the desirability of an investigation into the diluvial (= Pleistocene) fauna of the Netherlands Indies'. 'Officer of Health' Eug. Dubois in Padang argues that Pleistocene mammals from Java are still little known (except Martin (1887) paper), and can be compared to Siwalik-fauna of India. Fossil faunas which can be expected in caves also barely investigated. Also argues that great apes are limited to the Tropics, and that ancestors of humans should be expected in the Tropics, possibly in the three larger Sunda islands)

Dubois, E. (1891)- Voorlopig bericht omtrent het onderzoek naar de Pleistocene en Tertiaire vertebraten-fauna van Sumatra en Java, gedurende het jaar 1890. Natuurkundig Tijdschrift voor Nederlandsch-Indie 51, p. 93-100.
('Preliminary report on investigation of Pleistocene and Tertiary vertebrate fauna of Sumatra and Java during the year 1890'. Mentions excavation of Lida Ajer cave in Padang Highlands, with rich Late Pleistocene rainforest mammalian fauna, incl. orang-utan, and two human teeth (see also Hooijer 1948, Westaway et al. 2017))

Dubois, E. (1894)- *Pithecanthropus erectus*. Eine menschenaehnliche Uebergangsform aus Java. Landsdrukkerij, Batavia, p. 1-64.
(online at: https://ia802704.us.archive.org/28/items/Pithecanthropus00Dubo/Pithecanthropus00Dubo.pdf)
(also in Jaarboek Mijnwezen Nederlandsch Oost-Indie 1895, Wetenschappelijk Gedeelte, p. 5-77)
('Pithecanthropus erectus a transitional human-like transitional form from Java' Classic first description of 'Java man'/ Homo erectus, based on skull cap, femur (upper thigh bone)and molar from Trinil)

Dubois, E. (1896)- On *Pithecanthropus erectus*: a transitional form between Man and the apes. Journal of the Anthropological Institute of Great Britain and Ireland 25, p. 240-255.
(English language abstract of Dubois (1894) about Homo erectus discovery at Trinil, Java, with extensive discussions with members of the Royal Anthropological Institute)

Dubois, E. (1896)- On *Pithecanthropus erectus*, a transitional form between man and the apes. Transactions Royal Dublin Society, Ser. 2, 6, p. 1-18.

Dubois, E. (1907)- Eenige van Nederlandschen kant verkregen uitkomsten met betrekking tot de kennis der Kendengfauna (fauna van Trinil). Tijdschrift Koninklijk Nederlands Aardrijkskundig Genootschap (2) 24, p. 449-458.
('Some results from the Netherlands side regarding the knowledge of the Kendeng (Trinil) fauna'. Brief review of Pleistocene mammal fossils. Most bones in coarser lapilli tuffs, suggesting killing by volcanic eruptions. Fragmentary nature of fossils probably due to crocodiles)

Dubois, E. (1908)- Das geologische Alter der Kendeng- oder Trinil-Fauna. Tijdschrift Koninklijk Nederlands Aardrijkskundig Genootschap (2), 25, p. 1235-1270.
('The geologic age of the Kendeng or Trinil fauna'. Suggests most likely age of hominid bearing beds is Late Pliocene (subsequent workers all assume Pleistocene age))

Dubois, E. (1920)- De proto-Australische fossiele mensch van Wadjak (Java), I-II. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 19, p. 88-105 and p. 866-887.
('The Proto-Australian fossil man of Wadjak (Java)'. Dutch version of Dubois 1922).

Dubois, E. (1922)- The Proto-Australian fossil man of Wadjak, Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 23, 7, p. 1013-1051.
(online at: https://ia802802.us.archive.org/14/items/proceedingsofsec232koni/proceedingsofsec232koni.pdf)
(On 'Wajak Man', from slopes of Gunung Lawa limstone scarp near Wajak, 2 km SSW of Tjampur Darat, E Java (believed by Dubois to be of Pleistocene age and ancestor of Australian aborigenes, but age too young for that? (6-10 ka or younger; Storm 1995))

Dubois, E. (1924)- On the principal characters of the cranium and the brain, the mandible and the teeth of *Pithecanthropus erectus*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 27, 3, p. 265-278.
(Dutch version of Dubois 1926 in Proc. KNWAW 29, 5)

Dubois, E. (1924)- Figures of the calvarium and endocranial cast, a fragment of the mandible and three teeth of *Pithecanthropus erectus*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 27, 5, p. 459-464.
(Illustration of mandible found by Dubois in Kedung Brubus, E Java, in 1890, but not described before (see also Tobias 1966))

Dubois, E. (1926)- *Manis palaejavanica*, het reuzenschubdier van de Kendeng fauna. Verslagen Koninklijke Akademie van Wetenschappen, Amsterdam, 35, 8, p. 949-958.
(Dutch version of Dubois 1926)

Dubois, E. (1926)- *Manis palaejavanica*, the giant pangolin of the Kendeng fauna. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 29, 9, p. 1233-1243.
(online at: https://dwc.knaw.nl/DL/publications/PU00015393.pdf)
(Description of large Pleistocene ant eater from Kedung Brubus, 40 km ESE of Trinil, C Java. Possibly suggesting drier climate than today)

Dubois, E. (1934)- New evidence of the distinct organization of *Pithecanthropus*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 37, 3, p. 139-145.
(online at: https://dwc.knaw.nl/DL/publications/PU00016532.pdf)

Dubois, E. (1935)- On the gibbon-like appearance of *Pithecanthropus erectus*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 38, 6, p. 578-585.
(online at: https://dwc.knaw.nl/DL/publications/PU00016738.pdf)
(Thighbones of Pithecanthropus erectus suggest close affinity with gibbon group of apes. In this paper Dubois no longer believes Pithecanthropus is the 'missing link' human ancestor)

Dubois, E. (1935)- The sixth (fifth new) femur of *Pithecanthropus erectus*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 38, 8, p. 850-852.

Dubois, E. (1936)- Racial identity of *Homo soloensis* Oppenoorth (including *Homo modjokertensis* von Koenigswald) and *Sinanthropus pekinensis* Davidson Black. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 39, 10, p. 1180-1185.
(online at: https://dwc.knaw.nl/DL/publications/PU00016989.pdf)

Dubois, E. (1937)- On the fossil human skulls recently discovered in Java and *Pithecanthropus erectus*. Man (Royal Anthropological Insitute of Great Britain and Ireland) 37, 1, p. 1-7.
(Recently discovered Homo soloensis at Ngandong, E Java, is primitive Homo sapiens and proto-Australian. 'He has nothing in common with Pithecanthropus erectus')

Dubois, E. (1938)- The mandible recently described and attributed to *Pithecanthropus* by G.H.R. von Koenigswald, compared with the mandible of *Pithecanthropus erectus* described in 1924 by Eug. Dubois. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 41, 2, p. 139-147.
(online at: https://dwc.knaw.nl/DL/publications/PU00017147.pdf)
(Critique of information sent in a private letter of July 1937 by Von Koenigswald from Bandung, about discovery of new Pithecanthropus mandible fragment)

Dubois, E. (1940)- The fossil human remains discovered in Java by G.H.R. von Koenigswald and attributed by him to *Pithecanthropus erectus* in reality remains of *Homo wadjakensis* (syn. *Homo soloensis*). Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 43, 4, p. 494-496.
(online at: https://dwc.knaw.nl/DL/publications/PU00017432.pdf)
(Another of cranky-old Dubois criticisms of Von Koenigswald's work, here claiming that all of Von K's Pithecanthropus finds are 'doubtless' Homo soloensis.)

Dubois, E. (1940)- The fossil human remains discovered in Java by G.H.R. von Koenigswald and attributed by him to *Pithecanthropus erectus* in reality remains of *Homo sapiens soloensis*. Continuation. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 43, 7, p. 842-854.
(online at: https://dwc.knaw.nl/DL/publications/PU00017474.pdf)
(Part 2 of Dubois criticisms of Von Koenigswald's work. Interesting comment (p. 851): 'Man did not come into being in the Darwinian way, by gradual transformations required by the outer world power, but by inner world, autonomous, power')

Dubois, E. (1940)- The fossil human remains discovered in Java by G.H.R. von Koenigswald and attributed by him to *Pithecanthropus erectus* in reality remains of *Homo sapiens soloensis*. Conclusion. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 43, 10, p. 1268-1275.

(online at: https://dwc.knaw.nl/DL/publications/PU00017522.pdf)
(Part 3 of Dubois criticisms of Von Koenigswald's work)

Duller, G.A.T. (2001)- Dating methods: the role of geochronology in studies of human evolution in Southeast Asia and Australasia. Progress in Physical Geography 25, 2, p. 267-276.

Durband, A.C. (2004)- A test of the multiregional hypothesis of modern human origins using basicranial evidence from Indonesia and Australia. Ph.D. Thesis University of Tennessee, Knoxville, p. 1-236.
(online at: https://trace.tennessee.edu/utk_graddiss/2155)
(Features on cranial base in Ngandong hominids unique to that group. Autapomorphic characters in Ngandong population, in conjunction with previous work on Pleistocene paleoecology of Java, suggests multiple hominid species inhabited Java in Pleistocene. Also provides strong evidence of discontinuity between Indonesian Homo erectus and earliest Homo sapiens in Australasian fossil record)

Durband, A.C. (2006)- Craniometric variation within the Pleistocene of Java: the Ngawi 1 cranium. Human Evolution 21, 3, p. 193-201.
(online at: https://www.researchgate.net/publication/226150468_Craniometric_Variation_within_the_Pleistocene _of_Java_The_Ngawi_1_Cranium)
(Ngawi 1 homonid cranium discovered in 1987 submerged in Solo River near village of Selo puro, E Java. Described earlier by Widianto and Zeitoun (2003). Ngawi calvaria strong shape similarities to Ngandong and Sambungmacan fossils)

Durband, A. (2008)- Mandibular fossa morphology in the Ngandong and Sambungmacan fossil hominids. Anatomical Record 291, p. 1212-1220.

Durband, A. (2009)- Southeast Asian and Australian paleoanthropology: a review of the last century. J. Anthropological Sciences 87, p. 7-31.
(Review of study of hominids of SE Asia and Australia since work of Dubois. Populations represented by fossils from Sangiran and Ngandong, Java, went extinct without contributing genes to modern Australians)

Duval, M., K. Westaway, J. Zaim, Y. Rizal, Aswan, M.R. Puspaningrum, A. Trihascaryo, P.C. H. Albers, H.E. Smith et al. (2021)- New chronological constraints for the Late Pleistocene fossil assemblage and associated breccia from Ngalau Sampit, Sumatra. Open Quaternary 7, 9, p. 1-23.
(online at: https://www.openquaternary.com/articles/10.5334/oq.96/)
(First numerical ages for palaeontological assemblage and associated sediment from Ngalau Sampit cave, near Payakumbuh, first examined by M.E. Dubois in 1889. Three tooth samples Sus, cervid, bovid) U-series/ESR mean age of 105 ± 9 ka. Three breccia samples mean age 93 ± 6 ka.). Third site from Padang Highlands to be radiometrically dated, after Lida Ajer and Ngalau Gupin. Correlates to MIS 5, and Punung site in Java)

Earl of Cranbrook, A.P. Currant & G.W. Davison (2000)- Quaternary mammal fossils from Borneo: *Stegodon* and *Hippopotamus*. Sarawak Museum Journal 55, 76, p. 215-233.
(First report of Pleistocene? Stegodon and Hippopotamus fossils from Borneo. Unfortunately, paper retracted in SMJ 62 (2006); Stegodon and Hippotamus fossils described here were purchased in a curio shop in Kuching and were said to be from Kalimantan, but now believed to be from Sangiran, Java)

Eckhardt, R.B., M. Henneberg, A.S. Weller & K.J. Hsu (2014)- Rare events in Earth history include the LB1 human skeleton from Flores, Indonesia, as a developmental singularity, not a unique taxon. Proc. National Academy of Sciences USA (PNAS) 111, 33, p. 11961-11966.
(online at: www.pnas.org/content/111/33/11961.full.pdf
(Homo floresiensis is regional variant of Homo sapiens. Abnormal features of specimen LB1 not necessarily typical of whole assemblage (disputed by Westaway et al. 2015))

Edwards Mackinnon, E. (1991)- The Hoabinhian in the Wampu/Lau Biang Valley of Northeast Sumatra: An update, In: P. Bellwood (ed.) 14th Congress of the Indo-Pacific Prehistory Association, Yogyakarta 1990, Bull. Indo-Pacific Prehistory Association (IPPA) 10, p. 132-142.
(online at: https://journals.lib.washington.edu/index.php/BIPPA/article/view/11302)
(One of the more recent reviews of rel. large prehistoric shell mounds along NE coast of Sumatra N of Medan ('kjokkenmoddinger' of Witkamp (1920), Schurmann (1931), etc., and their vaying states of destruction by native shell mining for lime production. Shell mounds may be as old as 8000- 10,000 years and contain 'Hoabinhian' monofacial pebble-core stone tools ('Sumatraliths') and skeletons of humans with Austro-Melanoid features)

Elbert, J. (1908)- De nieuwste onderzoekingen over het *Pithecanthropus* vraagstuk. Natuurkundig Tijdschrift voor Nederlandsch-Indie 67, 3-4, p. 125-142.
(online at: https://ia800605.us.archive.org/27/items/mobot31753002489901/mobot31753002489901.pdf)
('The latest investigations on the Pithecanthropus question'. October 1907 presentation by J. Elbert, participant in the Selenka expedition. Pithecanthropus is intermediate between humans and apes, as suggested by Dubois. With discussion of Trinil stratigraphy. Age of Pithecanthropus is Pleistocene. Plant fossils and common unweathered feldspars in Trinil Beds suggest climate was 3- 6°C cooler)

Elbert, J. (1908)- Uber das Alter der Kendeng-Schichten mit *Pithecanthropus erectus* Dubois. Neues Jahrbuch Mineralogie Geologie Palaeontologie, Beilage Band 25, p. 648-662.
('On the age of the Kendeng beds with Pithecanthropus erectus Dubois'. Overview of Kendeng- Solo River Late Pliocene- Pleistocene stratigraphy. Recognizes two similar-looking fluvial packages at Trinil: lower is true Trinil beds with Pithecanthropus and of Early Pleistocene age ('Unterdiluvial'), the upper one represents younger Solo River terrace deposits)

Elbert, J. (1909)- Dubois' Altersbestimmung der Kendengschichten- ein Wort der Entgegnung. Centralblatt Mineralogie Geologie Palaontologie 1909, 17, p. 513-520.
('Dubois' age determination of the Kendeng beds- a reply'. Reply to Dubois 1908 paper)

Elbert, J. (1911)- Die Selenka'sche Trinil-Expedition und ihr Werk. Centralblatt Mineralogie Geologie Palaontologie 1911, 23, p. 736-741.
(online at: www.biodiversitylibrary.org/item/192769#page/9/mode/1up)
(Review of Selenka & Blankenhorn 1911 report on Trinil expedition and its results)

Elmaleh, A., J.P. Valet, X. Quidelleur, A. Solihin, H. Bouquerel, T. Tesson, E. Mulyadi, A. Khokhlov & D. Wirakusumah (2004)- Palaeosecular variation in Java and Bawean islands (Indonesia) during the Brunhes chron. Geophysical J. International 157, 1, p. 441-454.
(Paleomagnetic study from lava flows and dykes of Merapi and Merbabu, Bromo-Tengger, Lurus and Bawean Island. Ages mainly in Brunhes chron. Few reverse polarity flows probably emplaced during late Matuyama chron. Bawean leucite-bearing volcanics M Pleistocene age (0.3-0.8 Ma))

Erdbrink, D.P. (1954)- Mesolithic remains of the Sampung stage in Java: some remarks and additions. Southwestern J. Anthropology, Albuquerque, 10, 3, p. 294-303.
(Mesolithic stone implements from Punung (Pacitan) and from area S of Sukabumi, W Java, associated with the macrodont '(large teeth') Melanesoid-Australoid-Papuan people that populated Java before the arrival of non-Melanesoid people with a Neolithic culture. It is here called Sampung stage (later Sampungian by Van Heekeren), based on the type localy at the Guwo Lowo rock shelter at Sampung on the W slope of Wilis volcano, first discovered by Van Es in 1926 and excavated by Van Stein Callenfels (1932))

ESCAP (Economic Social Comm. for Asia and Pacifiic (1994)- Quaternary stratigraphy of Asia and the Paciific IGCP 296, ESCAP Atlas of Stratigraphy XIII, UN ESCAP Mineral Resources Development Series 63, p. 1-158.
(online at: https://repository.unescap.org/server/api/core/bitstreams/898632e4-9825-43c0-bd52-a43677e9034f/content)

Fadjar, K.S. (2006)- L'industrie osseuse de l'horizon Keplek Holocene de la grotte Song Terus, Punung, Java Est. Museum National Histoire Naturelle Paris, Departement Prehistoire, Memoire 6, p. 1-78.
('The bone industry of the Holocene Keplek horizon of Song Terus cave, Punung, E Java'. Gunung Sewu site)

Fae, M. (1996)- Lithobiostratigraphy and fossil hominids of the Sangiran-Krikilan area, Java (Yogyakarta, Indonesia). Memorie di Scienze Geologiche, Padova, 48, p. 143-153.
(Good review of 300-350m thick Late Pliocene- Pleistocene stratigraphy and hominid, mammal and mollusc distributions in Sangiran dome, C Java. From old to young: Shallow marine Late Pliocene Upper Kalibeng Fm, E Pleistocene Pucangan Fm Lower lahar and lacustrine black clay with diatomite beds and rich in Corbicula (first hominids possibly from top of formation), and M Pleistocene fluvial Kabuh Fm (with bulk of Homo erectus finds; tektite layer near top; common reworked marine microfauna). Capped by Notopuro Fm lahars with erosional base. Four mud volcanoes in center of dome, with rock fragments including Eocene Nummulites limestone. Reliable stratigraphic information lacking for most hominid fossils)

Fairbairn, A.S., G.S. Hope & G.R. Summerhayes (2006)- Pleistocene occupation of New Guinea's highland and subalpine environments. World Archaeology 38, 3, p. 371-386.
(online at: http://palaeoworks-dev.anu.edu.au/wp-content/uploads/2012/08/Fairbairnetal2006.pdf)
(Human colonization of New Guinea Highlands pre-dated 35 ka. Plant food use dates from at least 31 ka, i.e. in earliest millenia of human presence. Humans persisted in intermontane valleys through Late Glacial Maximum)

Fakhri (2018)- Vertebrate fauna from Gua Sambangoala, Southeast Sulawesi. In: S. O'Connor et al. (eds.) The archaeology of Sulawesi: current research on the Pleistocene to the historic period, Terra Australis 48, ANU Press, Canberra, Chapter 10, p. 153-170.
(online at: https://press-files.anu.edu.au/downloads/press/n4569/pdf/ch10.pdf)
(Mid-Holocene faunal remains in Gua Sambangoala rock shelter. Faunal assemblage mainly deposited between
~5500-4500 cal BP. Fauna of rock shelter included anurans, lizards, snakes, birds, bear cuscus, dwarf cuscus, bats, monkeys, rats, squirrels, civet cats, pig-deer (babirusa), Sulawesi warty pig (Sus celebensis) and Anoa)

Falgueres, C. (2001)- Dating layers and fossils in Sangiran Dome: methods and results.In: T. Simanjuntak et al. (eds.) Sangiran: man, culture, and environment in Pleistocene times, Proc. International Colloquium on Sangiran, Solo- Indonesia 1998, National Research Centre of Archaeology, Yayasan Obor Indonesia, Jakarta, p. 309-319.

Falgueres, C., F. Semah, H. Saleki, S. Hameau, H. Tu, G. Feraud, H. Simanjuntak & H. Widianto (2016)- Geochronology of early human settlements in Java: what is at stake? In: A. Borel et al. (eds.) Southeast Asia: Human evolution, dispersals and adaptation, Quaternary International 416, p. 5-11.
(online at: www.researchgate.net/publication/287973740_Geochronology_of_early_human_settlements Etc.)
(About challenges of chronological framework of early human settlements of Java, from oldest Lower Pleistocene Homo erectus island dwellers, to modern humans dispersals after Last Glacial Maximum)

Falk, D, C. Hildebolt, K. Smith, M.J. Morwood, T. Sutikna, P. Brown, Jatmiko, E. Saptomo et al. (2005)- The brain of LB1, *Homo floresiensis*. Science 308, 5719, p. 242-245.
(online at: https://www.peterbrown-palaeoanthropology.net/Falk%202005.pdf)
(Homo floresiensis type specimen (LB1) shull endocast was compared to great apes and hominids. LB1 not a microcephalic or pygmy. Brain size versus body size scales like an australopithecine, but endocast shape resembles Homo erectus)

Fauzi, M.R., M.M. Ansyori, D. Prastiningtyas, M.F.S. Intan, U.P. Wibowo, Wulandari, H. Rahmanendra, H. Widianto & T. Simanjuntak (2016)- Matar: a forgotten but promising Pleistocene locality in East Java. In: A. Borel et al. (eds.) Southeast Asia: Human evolution, dispersals and adaptation, Quaternary International 416, p. 183-192.
(Matar locality across from Ngandong on E bank of Solo River, 12km N of Ngawi, with fossils and lithic artifacts in terraces of poorly consolidated sand and gravels. Visited earlier by Bartstra et al. (1988). New

excavations yielded 4970 vertebrate fossils, incl. Stegodon trigonocephalus, Bubalus paleokarabau, Bibos paleosondaicus and Hexaprotodon sivalensis, typical of dry and open-woodland environment in M-U Pleistocene. Also 112 lithic flake-tools and some crude core-tools, similar to Ngandong)

Fauzi, M.R., M.F.S. Intan & T. Simanjuntak (2015)- Karakter teknologi litik *Homo erectus* progresif berdasarkan himpunan artefak dari Situs Matar, Bojonegoro. Kalpataru, Majalah Arkeologi 24, 1, p. 1-11.
(online at: https://jurnalarkeologi.kemdikbud.go.id/index.php/kalpataru/article/view/41/18)
('Lithic technology characteristic of progressive Homo erectus based on artifact assemblage from Matar Site, Bojonegoro'. New Matar site on E banks of Solo River (equivalent of 20m terrace of Ngandong?) with stone tool assemblage of 'progressive Homo erectus'. Flakes and massive tools such as bola, spheroidal, polyhedrons, and chopper-chopping tools)

Fauzi, M.R., T. Sutikna, F.S. Intan, S.E. Prasetyo, H. Widianto, F. Semah, T. Simanjuntak & H. Forestier (2025)- Further insight into the shape variability of handaxes from Indonesia. Archaeological Research in Asia 44, 100663, p.
(Study of 44 bifacial handaxes, which are rel. widespread across Indonesian archipelago, even beyond Wallace Line. Distribution of handaxes and Acheulean tools in Indonesia challenges outdated Movius Line theory of 1948)

Fauzi, M.R., A.A. Oktaviana, M.M. Ansyori, S. Noerwidi, D. Prastiningtyas, S.E. Prasetyo, Budiman et al. (2023)- The excavation of Gua Harimau's Western Gallery: a contribution to the Terminal Pleistocene-Early Holocene archaeological records in Sumatra. L'Anthropologie 127, 3, 103156, p. 1-20.
(Excavations at Harimau cave in 2009-2016 show three episodes of latest Pleistocene- Holocene human occupation: Preneolithic, Neolithic and Paleometalic (early metal age) with significant differences in artifacts)

Foo Shu Tieng (2015)- The Guar Kepah shell middens: evidence and questions. In: First SEAMEO SPAFA Conference Advancing Southeast Asia Archaeology, Chonburi, Thailand, Chapter 12, p. 114-139.
(online at: www.researchgate.net/publication/316957279_The_Guar_Kepah_Shell_Middens_Evidence Etc.)
(Good review of mid-Holocene edible shell middens with human artifacts at both sides of Malacca Straits, at Penang and NE Sumatra)

Forestier, H. (1999)- L'assemblage industriel de Song Keplek, Java Est (un nouveau regard sur l'outillage lithique de l'homme moderne au debut de l'Holocene en Indonesie). Bull. Ecole francaise d'Extreme-Orient 86, 1, p. 129-159.
(online at: www.persee.fr/docAsPDF/befeo_0336-1519_1999_num_86_1_3409.pdf)
('The industrial assemblage of Song Keplek, East Java; a new view on stone tools of modern man at the beginning of the Holocene in Indonesia'. Song Keplek stone tool assemblage from Punung area ~6000-4000 years old)

Forestier, H. (2007)- Les eclats du passe prehistorique de Sumatra: une tres longue histoire des techniques. Archipel 74, p. 15-44.
(online at: www.persee.fr/docAsPDF/arch_0044-8613_2007_num_74_1_3914.pdf)
(Long history of prehistoric stone tool making in Sumatra, incl. Acheulean tools from Ogan River tributaries in S Sumatra, possibly from Homo erectus, 'Hoabinhian' shell middens 30m wide x 5m high in NE Aceh, etc.)

Forestier, H. (2007)- Ribuan Gunung, ribuan alat batu- prasejarah Song Keplek, Gunung Sewu, Jawa Timur. Kepustakaan Populer Gramedia (KPG), p. 1-314.
(online at: https://horizon.documentation.ird.fr/exl-doc/pleins_textes/divers14-11/010039121.pdf)
('Thousands of Mountains, thousands of stone tools- prehistoric Song Keplek, Gunung Sewu, East Java'. On the ~5000 year-old (Paleolithic) types of stone tools of Song Keplek. Originally discovered by Von Koenigswald in 1934 (Revised Indonesian edition of 1998 dissertation at Museum National d'Histoire Naturelle, Paris))

Forestier, H. (2024)- Stone tools in Palaeolithic Sumatra, Indonesia: From *Homo erectus* to the Hoabinhian. In: J. Louys et al. (eds.) Quaternary Palaeontology and Archaeology of Sumatra, Terra Australis 56, ANU Press, Canberra, Chapter 10, p. 219-231.
(online at: https://press-files.anu.edu.au/downloads/press/n12084/pdf/ch10.pdf)

Forestier, H., D. Driwantoro, D. Guillaud, Budiman & D. Siregar (2006)- New data for the prehistoric chronology of South Sumatra. In: T. Simanjuntak et al. (eds.) Archaeology: Indonesian perspective, R.P. Soejono's Festschrift, LIPI Press, Jakarta, p. 177-192.
(online at: http://horizon.documentation.ird.fr/exl-doc/pleins_textes/divers17-07/010041012.pdf)

Forestier, H., M. Grenet, A. Borel & V. Celiberti (2017)- Les productions lithiques de l'Archipel indonesien. Journal of Lithic Studies 4, 2, p. 231-303.
(online at: http://journals.ed.ac.uk/lithicstudies/article/view/2544/3753)
('The lithic productions of the Indonesian Archipelago'. Extensive review of Pleistocene stone tool 'industries')

Forestier, H. & E. Patole-Edoumba (2000)- Les industries lithiques du Paleolithique tardif et du debut de l'Holocene en Insulinde. Aseanie 6, 1, p. 13-56.
(online at: www.persee.fr/doc/asean_0859-9009_2000_num_6_1_1683)
(Review of major Late Paleolithic lithic (stone tool) assemblages in Indonesia and Philippines)

Forestier, H., T. Simandjuntak, F. Detroit & V. Zeitoun (2010)- Unite et diversite prehistorique entre Java et Sumatra. Archipel 80, p. 19-44.
(online at: www.persee.fr/docAsPDF/arch_0044-8613_2010_num_80_1_4175.pdf)
('Prehistoric unity and diversity of Java and Sumatra'. Prehistory of Indonesia from 20000- 5000 BP. Java marked by technical heterogeneity in produced stone tools, Sumatra more homogenous technical choices, with unifacial pebble shaping which still belongs to Hoabinhian tradition)

Forestier, H., T. Simandjuntak & D. Driwantoro (2005)- Les premiers indices d'un facies Acheuleen a Sumatra-Sud, Indonesia. In: L. Faton (ed.) Asie du Sud-Est: de l'*Homo erectus* a l'*Homo sapiens*, Dossiers d'Archeologie 302, 16-17.
('The first indications of an Acheuleen facies in South Sumatra'. Paleolithic stone tools from Ogan River tributaries)

Forestier, H., T. Simanjuntak, D. Guillaud, D. Driwantoro, K. Wiradnyana, D. Siregar, R. Due Awe & Budiman (2005)- Le site de Togi Ndrawa, ile de Nias, Sumatra nord: les premieres traces d'une occupation hoabinhienne en grotte en Indonesie. Comptes Rendus Palevol 4, p. 727–733.
(The Togi Ndrawa site, Nias Island, Noth Sumatra: the first record of a Hoabinhian cave settlement in Indonesia'. Late Pleistocene- E Holocene classic Hoabinhian pebble artefacts, forest and coastal fauna and human bones in Togi Ndrawa cave, NE Nias)

Forestier, H., V. Zeitoun, A. Lenoble & C. Tiamtinkrit (2006)- Discovery of new old material in the basaltic region of Lampang (Northern Thailand): a techno-functional interpretation. In: Archaeology in Southeast Asia: from *Homo erectus* to the living traditions, 11th International Conference European Assoc. SE Asian Archaeologists (EurASEAA 2006), Bougon papers, p. 35-39.
(online at: https://www.researchgate.net/publication/281801991_Discovery_of_new_old_material_in_the_basaltic_region_of_Lampang_Northern_Thailand_a_techno-functional_interpretation)
('Technical analysis' of Pleistocene stone artefacts (choppers, chopping tools) from above basalt level of Ban Don Mun in Lampang province (N Thailand))

Frantz, L.A.F., A. Rudzinski, A.M S. Nugraha, A. Evin, J. Burton et al. (2018)- Synchronous diversification of Sulawesi's iconic artiodactyls driven by recent geological events. BioRxiv, p. 1-22.
(preprint online at: https://www.biorxiv.org/content/biorxiv/early/2018/01/04/241448.full.pd)

(Paleogeographical reconstructions with genetic and morphometric data from Sulawesi mammals Babirusa, Anoa and Sulawesi warty pig indicate almost synchronous expansion from central part of island after recent emergence of land on Sulawesi (~1-2 Ma))

Gaffney, D. (2021)- Pleistocene water crossings and adaptive flexibility within the *Homo* genus. J. Archaeological Research 29, p. 255-326
(online at: https://link.springer.com/article/10.1007/s10814-020-09149-7)
(Crossing of substantial bodies of straits, seas, and oceans to arrive at new landmasses has long been seen as innovation within our species, Homo sapiens, sometime after 60 ka. However, increasing evidence that older hominins, incl. Homo erectus, also made significant water crossings)

Gathorne-Hardy, F.J. & W.E.H. Harcourt-Smith (2003)- The super-eruption of Toba, did it cause a human bottleneck? J. Human Evolution 45, p. 227-230.
(No evidence to support hypothesis that Toba supereruption at 73.5 ka caused bottleneck in human, animal or plant populations)

Ge, Y. & X. Gao (2020)- Understanding the overestimated impact of the Toba volcanic super-eruption on global environments and ancient hominins. Quaternary International 559, p. 24-33.
(Although Toba 74 ka super-eruption was largest such event of Quaternary, it did not have a devastating impact on Earth climate and human activity, as suggested by some authors)

Ghosh, A.K. (1978)- Further note on Sangiran flake industry. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 81, 2, p. 211-214.

Glover, I.C. (1981)- Leang Burung 2: an Upper Palaeolithic rock shelter in South Sulawesi, Indonesia. Modern Quaternary Research in Southeast Asia 6, Balkema, Rotterdam, p. 1-38.
(Late Pleistocene (~30ka) flaked stone tools from shelter at base of Leang Burung limestone cliff E of Tompokbalang in Maros District, S Sulawesi)

Grenet, M., J. Sarel, R. Fauzy, A.A. Oktaviana, B. Sugiyanto, J.M. Chazine & F.X. Ricaut (2016)- New insights on the late Pleistocene- Holocene lithic industry in East Kalimantan (Borneo): the contribution of three rock shelter sites in the karstic area of the Mangkalihat peninsula. In: A. Borel et al. (eds.) Southeast Asia: Human evolution, dispersals and adaptation, Quaternary International 416, p. 126-150.
(Stone tools in three rock shelter sites in karst area of Mangkalihat Peninsula (Liang Abu, Liang Jon and Liang Pemalawan). Reduction sequences, flaking technology and typology of blanks and tools, persisted across Pleistocene/Holocene boundary for >20,000 years. Also apparent climatic stability over last 40,000 years)

Grimaud-Herve, D. (1986)- The parietal bone of Indonesian *Homo erectus*. Human Evolution 2, p. 167-181.
(Comparative study of parietal (skull) bones from Sangiran, Sambungmacan 1 and Ngandong. Some morphological metrical features allow separation of Sangiran and Ngandong samples. Sambungmacan 1, whose chronological age is not well established, appears closer to Ngandong men)

Grimaud-Herve, D. (1998)- Taxonomic position of the Sangiran 31 hominid. In: T. Simanjuntak et al. (eds.) Sangiran: man, culture, and environment in Pleistocene times, Proc. International Colloquium on Sangiran, Solo- Indonesia, Solo 1998, National Research Centre of Archaeology, Yayasan Obor Indonesia, Jakarta, p. 46-53.
(Sangiran 31 damaged cranial remains discovered by Sartono and Djubiantono in Pucangan Fm of Sangiran in 1979. Here deemed to a robust varaint of other known Homo erectus from this site)

Grimaud-Herve, D., A. Balzeau, H. Widianto, T. Djubiantono, F. Detroit, A.M. Moigne, A.M. Semah, A. Purnomo, M. Ansyori, B. Brasseur, T. Ingicco & F. Semah (2016)- Position of the posterior skullcap fragment from Sendang Klampok (Sangiran Dome, Java, Indonesia) among the Javanese *Homo erectus* record. Quaternary International 416, p. 193-209.

(New Homo erectus skullcap fragment fossil from 'Grenzbank' at Sendang Klampok in NNW Sangiran Dome, associated with rel. rich 'Trinil-type' mammal assemblage. Among more ancient hominids from Sangiran dome. Many characters closer to Ngandong hominins, different from those of Kabuh (Bapang) and Zhoukoudian Lower Cave specimens (?))

Grimaud-Herve, D. & J.L. Franzen (1994)- Evolution of the Javanese fossil hominid brain. In: In: J.L. Franzen (ed.) 4th Int. Conference 100 years of *Pithecanthropus*. The *Homo erectus* problem, Frankfurt 1991, Courier Forschungsinstitut Senckenberg, Frankfurt, 171, p. 61-68.

Grimaud-Herve, D., F. Valentin, F. Semah, A.M. Semah & H. Widianto (1994)- Le femur humain Kresna 11 compare a ceux de Trinil. Comptes Rendus Academie Sciences, Paris 318, II, p. 1139-1144.
('The human femur Kresna 11 compared to those from Trinil'. Thigh bone named Kresna 11 discovered in 1992 from the 'Grenzbank' horizon in Sangiran Dome shows several Homo erectus features, similar to Trinil (see also Puymerial et al. 2012))

Grimaud-Herve, D. & H. Widianto (1993)- Les Hominides de Java. In: F. Semah & D. Grimaud-Herve (eds.) Le Pithecanthrope de Java a la decouverte du chainon manquant, Les dossiers d'Archeologie 184, p. 30-45.
('The hominids of Java')

Grimaud-Herve, D. & H. Widianto (2001)- Les fossils humains decouverts a Java depuis les annees 1980. In: F. Semah et al. (eds.) Origin des peuplements et chronologie des cultures Paleolithiques dans le sud-est Asiatique. Semenanjung, Paris, p. 331-357.
('The fossil humans discovered in Java since the 1980's')

Grimaud-Herve, D., H. Widianto, F. Detroit & F. Semah (2012)- Comparative morphological and morphometric description of the hominin calvaria from Bukuran (Sangiran, Central Java, Indonesia). J. Human Evolution 63, p. 637-652.
(online at: www.academia.edu/24123718/Comparative_morphological_and_morphometric_descript Etc.)
(Description of heavily mineralized Homo erectus skull discovered in 1977 near Sendangbusik, Bukuran area, E side of Sangiran dome, C Java. In cross-bedded fluvio-volcanic sands of lower Kabuh (Bapang) Fm, ~14m above 'Grenzbank')

Grimaud-Herve, D., H. Widianto & T. Jacob (2000)- Two new human fossil remains discovered in Sangiran (Central Java, Indonesia). In: W. Dong (ed.) Proc. 1999 Beijing International Symposium on paleoanthropology, Acta Anthropologica Sinica 19 (Supplement), p. 41-45.
(Fragmented Homo erectus skull from Kabuh Beds at Grogol Wetan site. Ar-dating indicates 0.78 ± 0.29 Ma. Second hominid skull in 1996 from Kabuh Beds at Bukuran site)

Grine, F. & J.L. Franzen (1994)- Fossil hominid teeth from the Sangiran Dome (Java, Indonesia). In: J.L. Franzen (ed.) 4th International Conference 100 years of *Pithecanthropus*. The *Homo erectus* problem, Frankfurt 1991, Courier Forschungsinstitut Senckenberg, Frankfurt, 171, p. 75-103.

Groves, C.P. (1976)- The origin of the mammalian fauna of Sulawesi (Celebes). Zeitschrift fur Saugetierkunde 41, 4, p. 201-216.
(online at: www.landesmuseum.at/pdf_frei_remote/Zeitschrift-Saeugetierkunde_41_0201-0216.pdf)
(Discussion of present-day mammal faunas of Sulawesi and relation to classic faunal province boundaries suggested for Indonesia ((Wallace, Weber, Lydekker Lines))

Groves, C.P. (1984)- Mammal faunas and the palaeogeography of the Indo-Australian Region. In: P. Andrews & J.L. Franzen (eds.) The early evolution of man with special emphasis on Southeast Asia and Africa. Courier Forschungsinstitut Senckenberg 69, p. 267-273.

Groves, C.P. (1985)- Plio-Pleistocene mammals in island southeast Asia. Modern Quaternary Research in Southeast Asia 9, Balkema, Rotterdam, p. 43-54.

Groves, C.P. (2001)- Mammals in Sulawesi: where did they come from and when, and what happened to them when they got there? In: I. Metcalfe, J.M.B. Smith et al. (eds.) Faunal and floral migrations and evolution in SE Asia- Australia, A.A. Balkema, Lisse, p. 333-342. *(online at: https://www.academia.edu/296852/Faunal_and_Floral_Migrations_and_Evolution_In_SE_Asia_Australasia)*
(Present day faunal endemism suggests Sulawesi was cluster of islands until quite late in geologic time)

Grun, R. & A. Thorne (1997)- Dating the Ngandong humans. Science 276, p. 1575-1576.
(Critique of Swisher et al. 1994, 1996 dating results of Ngandong hominids. Consider the Solo high terrace to represent a mix of materials reworked from different levels, sites, and ages)

Gruwier, B.J. (2017)- The large vertebrate remains from Bindjai Tamieng (Sumatra, Indonesia). J. Indo-Pacific Archaeology 41, p. 22-29.
(online at: https://journals.lib.washington.edu/index.php/JIPA/article/download/15027/12536)
(Study of animal- and human remains excavated in 1928 by BPM geologist Schurmann from Early Holocene man-made Binjai Tamieng shell midden, along lower course of the Tamiang River in NE Sumatra coastal plain. Original site now completely destroyed, but part of Schurmann collection of vertebrates deposited in Naturalis Museum, Leiden, and briefly examined by Hooijer (1948). These include marine turtles, fish and artiodactyls (deer, pig), while crocodiles and a small whales were probably opportunistically hunted or scavenged. Also rhinoceros and human remains showing traces of disarticulation, possibly indicative of funerary rituals or cannibalism)

Gruwier, B.J. (2019)- *Homo erectus* palaeoecology in Java: a study of cervid post-cranial ecomorphology. Ph.D. Thesis Durham University, p. 1-391..
(online at: https://etheses.dur.ac.uk/13666/)
(Paleoenvironmental reconstructions of several Pleistocene sites from Java (Trinil, Kedung Brubus and Sangiran), based on morphologies of cervids (deer fossils). Trinil open woodland, but relatively wet substrate. Kedung Brubus open, but drier conditions, presumably coincided with glacial stage when Java was connected to Asian mainland. Material from Sangiran suggests open conditions, and either dry or wet substrates)

Gruwier, B., J. de Vos & K. Kovarovic (2015)- Exploration of the taxonomy of some Pleistocene Cervini (Mammalia, Artiodactyla, Cervidae) from Java and Sumatra (Indonesia): a geometric- and linear morphometric approach. Quaternary Science Reviews 119, p. 35-53.
(Study of deer fossils. Medium sized deer of genus Axis and large sized genus Cervus s. l. in Java. Axis lydekkeri and Axis javanicus are considered valid taxa. Large deer, such as Cervus kendengensis, Cervus stehlini and Cervus problematicus most likely of subgenus Rusa. Sumatran fossils members of subgenus Rusa)

Habinger, S.G., O. Chavasseau, J.J. Jaeger, Y. Chaimanee, A.N. Soe, C. Sein & H. Bocherens (2022)- Evolutionary ecology of Miocene hominoid primates in Southeast Asia. Nature Scientific Reports 12, 11841, p. 1-12.
(online at: https://www.nature.com/articles/s41598-022-15574-z)
(Paleoecology of Miocene and younger apes in SE Asia, like Late Miocene Khoratpithecus ayeyarwadyensis from Myanmar, similar to modern orangutans (Pongo):in woodland forest)

Hakim, B., U.P. Wibowo, G.D van den Bergh, D. Yurnaldi, R. Joannes-Boyau, A. Duli, Suryatman, R. Sardi et al. (2025)- Hominins on Sulawesi during the Early Pleistocene. Nature 646, p. 378-383.
(online at: https://www.nature.com/articles/s41586-025-09348-6)
(Previous excavations on Sulawesi revealed stone artefacts with minimum age of 194 ka in Walanae Depression, long before earliest known Homo sapiens arrived in area (73-63 ka in Sunda). Stone artefacts at Calio in fossiliferous layers dated to at least 1.04 Ma, possibly up to 1.48 Ma (E Pleistocene), suggesting Sulawesi populated by hominins at about same time as Flores hominins, if not earlier)

Hameau, S., C. Falgueres, J.J. Bahain, F. Semah, A.M. Semah & J.M. Dolo (2007)- ESR dating in Song Terus cave (East Java, Indonesia). Quaternary Geochronology 2, p. 398-402.

(Dating on animal teeth from Song Terus cave, Gunung Sewu, E Java shows cave belongs to karstic system which has been in place since M Pleistocene (216, 392 ka))

Handini, R., S. Noerwidi, H.O. Sofian, M.R. Fauzi, U. Prasetyo, I.M. Geria, M Ririmasse et al. (2023)- New evidence on the early human occupation in Sumba Islands. L'Anthropologie 127, 3, Special Issue Prehistoire de l'Indonesie, 103152, p. 1-13.
(Excavations of archaeological sites in Sumba in 2016-2022 revealed individual remains, burial jars and artifacts suggesting Austronesian settlements with a complex culture inhabiting Sumba around 3000 years ago)

Harbowo, D,G. & L.K. Agustina (2024)- Biogeographical land bridges of Bali-Nusa Penida-Lombok: A possible dispersal pathway for terrestrial fauna during the Pleistocene Glacial periods. Proc. 6th EMBRIO International Symposium (EIS 2023), Bogor 2023, BIO Web of Conferences 112, 10001p. 1-11.
(online at: https://www.bio-conferences.org/articles/bioconf/pdf/2024/31/bioconf_eis2024_10001.pdf)
(Migration of prehistoric humans and terrestrial fauna during Pleistocene glacial periods (across Wallace Line) possibly facilitated by possible biogeographical land bridge between Bali and Lombok, via Nusa Penida)

Hardjasasmita, H.S. (1982)- Kehadiran elemen fauna vertebrata Asia di sebelah timur garis Wallacea pada kala Plestosen. Kumpulan Makalah Pertemuan Ilmiah Arkeologi (PIA) II, Jakarta 1980, Pusat Penelitian Arkeologi Nasional, p. 521-532.
('Presence of elements of Asian vertebrate fauna East of the Wallace Line during the Pleistocene')

Hardjasasmita, H.S. (1983)- Evolution of the genus *Sus* (Suidae, Mammalia) in Indonesia. Taxonomy, phylogeny and paleogeography. Ph.D. Thesis Institut Teknologi Bandung (ITB), p. 1-177. *(Unpublished)*
(In Indonesia fossil suids (pig family) known from Pliocene- Holocene, mainly represented by genus Sus. On Sulawesi also Babyrousa and Celebochoerus. Two fossil species (Sus brachygnathus, S. macrognathus), one subspecies (Sus macrognathus terhaari) and two questionable species (Sus stremmi, S. sangiranensis), nine Recent species and subspecies are recognized)

Hardjasasmita, H.S. (1987)- Taxonomy and phylogeny of the Suidae (Mammalia) in Indonesia. Scripta Geologica 85, p. 1-68.
(online at: https://repository.naturalis.nl/pub/317394)
(Overview of ?Pliocene- Recent wild pigs in Indonesia, mainly of genus Sus, mainly from Java)

Harrison, T. (1978)- Present status and problems for Paleolithic studies in Borneo and adjacent islands. In: F. Ikawa-Smith (ed.) Early Palaeolithic in South and East Asia, Mouton Publishers, The Hague, p. 38-57.
(Few or no Paleolithic fossils found on Borneo (unlike Sulawesi, Java, etc.) and some may be Chinese drugstore imports. Niah cave in Sarawak rel. rich record of human and associated fossils dating to ~35 ka. M Pleistocene tektites of coastal NW Brunei cannot be used for dating of 'Jerudong Terrace', as most or all are reworked into younger gravel terraces)

Harrison, T., J. Krigbaum & J. Manser (2006)- Primate biogeography and ecology on the Sunda Shelf islands: a paleontological and zooarchaeological perspective. In: S.M. Lehman & J.G. Fleagle (eds.) Primate biogeography, Springer, New York, p. 331-372.
(Non-human primates on Sundaland taxonomically diverse (27 species), and relatively high provinciality and endemism. By Late Pliocene main islands of Sunda Shelf had primate fauna that included Pongo pygmaeus, Hylobates spp., Macaca nemestrina etc. on Sumatra, Java, Borneo and Mentawai Islands. Most probably arrived during Pretiglian cold phase, starting at ~2.8 Ma, when sea levels fell by >100m)

Hartatik, Sunarningsih, N.M.E. Fajari & H.O. Sofian (2023)- New discoveries of ancient iron smelting sites in Kalimantan. L'Anthropologie 127, 3, 10316, p. 1-16.
(Evidence of iron making in the Meratus mountains, SE Kalimantan and Lamandau watershed in SW Kalimantan. Radiocarbon dating at Cililin 1 Cave site in Meratus Mountains indicates earliest iron ore smelting occurred during early Christian era (2000-1700 BP))

Harvati, K. & H. Reyes-Centeno (2022)- Evolution of *Homo* in the Middle and Late Pleistocene. J. Human Evolution 173, 103279, p. 1-12.
(Review of complex evolution of 19 named species of M-L Pleistocene hominids, incl. Home floresiensis and H. luzonensis in SE Asia)

Hawkins, S., S. O'Connor, T.R. Maloney, M. Litster, S. Kealy, J.N. Fenner, K. Aplin et al. (2017)- Oldest human occupation of Wallacea at Laili Cave, Timor-Leste, shows broad-spectrum foraging responses to late Pleistocene environments. Quaternary Science Reviews 171, p. 58-72.
(online at: http://www.sciencedirect.com/science/article/pii/S0277379117302470)
(Laili Cave in Laleia, Timor-Leste, preserves oldest occupation in Wallacea (~43-45 ka) of modern humans, earlier than other Pleistocene sites known in Wallacea (E Indonesia). Pleistocene humans used abundant local chert and engaged in mobile broad-spectrum foraging)

Hayes, S., G.D. van den Bergh, I. Sutisna, H. Insani, U.P. Wibowo, R. Setiawan, I. Kurniawan & S.T. Turvey (2025)- Insular Mid-Pleistocene giant rats from the So'a Basin (Flores, Indonesia). Quaternary (MDPI) 8, 3, 44 p.
(online at: https://www.mdpi.com/2571-550X/8/3/44)
(M PleistoceneMata Menge site on Flores yielded > 670 fossil murine (rat) molars of Papagomys group from two intervals. Lower interval (0.7 Ma) smaller size molars similar to Hooijeromys nusatenggara Musser 1981. Upper interval (~70 ka) most similar to mid-Holocene Papagomys theodorverhoeveni)

Heaney, L.R. (1985)- Zoogeographic evidence for Middle and Late Pleistocene land bridges to the Philippine islands. Modern Quaternary Research in Southeast Asia 9, p. 127-143.
(In Sunda shelf region rel. widespread faunal distribution, corresponding stronlgy with M and Late Pleistocene land bridge formation. Number of species corresponds with size of island area. Elephant species rel. widespread in SE Asia Pleistocene, probably because they are strong swimmers and do not necessary need land bridges. Unlikely there was land bridge between Asia and The Philippines, except M Pleistocene connection from NE Borneo to Palawan. Therefore unlikely that Homo erectus reached Philippines)

Heaney, L.R. (1986)- Biogeography of mammals in SE Asia: estimates of rates of colonization, extinction and speciation. Biological J. Linnean Society, London, 28, 1-2, p. 127-165.

Heaney, L.R., D.S. Balete & E.A. Rickart (2016)- The mammals of Luzon Island: biogeography and natural history of a Philippine fauna. Johns Hopkins University Press, Baltimore, p. 1-285.

Hemmer, H. (1971)- Fossil mammals of Java II. Zur Fossilgeschichte des Tigers (*Panthera tigris* L.) in Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 74, 1, p. 35-52.
('Fossil mammals of Java 2: the fossil record of the tiger Panthera tigris in Java'. On presence of extinct tiger 'Trinil Tiger' (Panthera tigris trinilensis) in beds of 1.2 Ma old at Trinil)

Hemmer, H. (1971)- Fossil mammals of Java III. Zur Kenntnis der Evolution javanischer Kleinkatzen: *Prionailurus bengalensis koenigswaldi* ssp. n. und *Felis chaus* ssp. aus dem Neolithikum von Sampung, Mittel-Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 74, p. 365-367.
('On the knowledge of the evolution of javanese small cats Prionailurus, etc. Two new subspecies from Neolithic of Sampung, C Java')

Hemmer, H. & G. Schutt (1972)- Pleistozane Leoparden (*Panthera pardus*) aus Java und Sudchina. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B76, p. 37-49.
('Pleistocene leopards from Java and S China')

Hemmer, H. & G.H.R. von Koenigswald (1964)- Fossile Nebelparder (*Neofelis*) aus dem Pleistozan Sudchinas und Javas. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 67, p. 1-16.
('Fossil leopards (Neofelis) from the Pleistocene of South China and Java')

Henneberg, M., R.B. Eckhardt, S. Chavanaves & K.J. Hsu (2014)- Evolved developmental homeostasis disturbed in LB1 from Flores, Indonesia, denotes Down syndrome and not diagnostic traits of the invalid species *Homo floresiensis*. Proc. National Academy of Sciences USA (PNAS) 111, 33, p. 11967-11972.
(online at: www.pnas.org/content/111/33/11967.full.pdf)
(LB1 type specimen of Homo floresiensis ('hobbit') viewed as anomalous specimen in small-bodied Australomelanesian Homo sapiens population, possibly afflicted by Down syndrome. Conclusion disputed by Westaway et al. 2015)

Henneberg, M. & J. Schofield (2008)- The Hobbit trap. Money, fame, science, and the discovery of a 'new species'. Wakefield Press, Kent Town, S.A., p. 1-159.
(Book critical of interpretations of Homo floresiensis ('hobbit'), claiming it to be much younger than reported and to be a 'normal' dwarfed Homo sapiens island population. Also 2011 2nd Edition)

Hennig, E. (1911)- Die Fischreste. In: E. Selenka & M. Blanckenhorn (eds.) Die Pithecanthropus Schichten auf Java, Engelmann, Leipzig, p. 54-60.
(Description of Pleistocene fish remains from Trinil, excavated by Selenka expedition)

Herman, D.Z. (2011)- Fossilization type of *Elephas hysudrindicus* from Blora on the basis of petrographic and Scanning Electron Microscopic analyses. Jurnal Geologi Indonesia 6, 2, p. 75-84.
(online at: www.bgl.esdm.go.id/publication/index.php/dir/article_detail/304)
(2009 discovery of nearly complete skeleton of Pleistocene elephant Elephas hysudrindicus in Solo River terrace in Sunggun area near Medalem Village (not clear from what part of Pleistocene/ terrace level?; JTvG). Some precipitation of calcite and other authigenic minerals)

Herrera, R.J. & R.G.-Bertrand (2022)- Interwoven evolution of *Homo erectus, Homo floresiensis*, and *Homo sapiens*. In: Sex and cohabitation among early humans, Elsevier, Chapter 16, p, 551-573.
(General review. Homo erectus was first of hominin ancestors to have sapiens-like body proportions (shorter arms and longer legs relative to its torso). H. erectus was also first hominin to migrate out of Africa and survived for more than 1.5 Myrs. Etc.)

Hertler, C., A. Bruch & M. Marker (2013)- The earliest stages of hominin dispersal in Africa and Eurasia. In: I. Ness (ed.) The Encyclopedia of Global Human Migration, Blackwell Publishing, Chapter 2, p. 1-9.
(online at: www.academia.edu/5016352/The_earliest_stages_of_hominin_dispersal_in_Africa_and_Eurasia)
(Brief review of earliest movements of hominins (proto-humans) in Africa, starting from ~7 Ma, and focusing on genesis of Australopithecus and Homo and dispersal into Eurasia soon after 2 Ma. Earliest evidence for hominins in E Asia is from C Java (Lower Pleistocene Homo erectus from Sangiran, with age of >1.5 Ma)

Hertler, C. & Y. Rizal (2005)- Excursion guide to the Pleistocene hominid sites in Central and East Java. JW Goethe University, Frankfurt, and ITB Bandung, p. 1-35.
(at www.palaeo.net/biologie/material/Excursion%20guide.pdf)
(Overview of Pleistocene mammal fauna biostratigraphy, paleoanthropology and fossil hominid sites)

Hertler, C., Y. Rizal & Y. Zaim (2007)- Habitat differentiation in the Pleistocene of Jawa- Introduction of the new Pleistocene fossil locality Majalengka. Courier Forschungsinstitut Senckenberg 259, p. 165-175.
(Pleistocene mammal faunas from Java three successive faunas based proboscidean genera: (1) E Pleistocene Mastodon-Geochelone fauna, (2) early M Pleistocene Stegodon- Homo erectus fauna and (3) late M Pleistocene Elephas- Homo sapiens fauna. Stegodon- Homo erectus fauna contains elements from successive migration waves and different ecological settings. Introduce model for endemic evolution in Java and newly discovered Pleistocene mammal locality in W Java)

Hertler, C., A.A.E. van der Geer, M.R. Puspaningrum, J.O Reschke, I.P. Anwar & E. Holzchen (2025)- *Stegodon* SEA-crossing: Swim, shrink, and disperse. Earth History and Biodiversity 4, 100026, p. 1-15.
(online at: https://www.sciencedirect.com/journal/earth-history-and-biodiversity/vol/4/suppl/C)

(Stegodons, an extinct family of proboscideans, managed to settle on roughly a dozen Wallacean (E Indonesia) islands during Pleistocene (Java, Flores, Sangihe, Sumba, Timor, Sulawesi, Luzon, Mindanao, Japan). Distribution interaction of (1) swimming performances and (2) island dwarfing. Reduction of body mass likely negatively affected swimming performance. No fossil record for Stegodon Maluku islands or in New Guinea/ NW Australia. With dispersal routes scenarios across Sundaland- Wallacea)

Hertler, C. & R. Volmer (2005)- Assessing prey competition in fossil carnivore communities- a scenario for prey competition and its evolutionary consequences for tigers in Pleistocene Java. Palaeogeogr. Palaeoclim. Palaeoecology 257, p. 67-80.
(online at: http://www.roceeh.net/fileadmin/download/Publications/Hertler-volmer-2008.pdf)
(Five carnivore species overlapped in time in Java Pleistocene: two pantherines, a hyaenid and two canid species, each in one or more faunal levels. Significant increase in body mass of tigers in Ngandong faunal level reflects intense competition among carnivores in preceding Kedung Brubus level)

Hidayah, A.R., U.P. Wibowo, A. Purwoarminta, G.J. Price & S. Noerwidi (2021)- Palaeoenvironments and palaeontology of the Atambua Basin, West Timor, Indonesia. Proc. 1st Asia-Pacific Conference on Human Evolution, Quaternary International 603, p. 82-89.
(M Pleistocene or earlier vertebrate faunas long known from Timor, but rarely studied. Dwarf proboscideans such as Stegodon once widespread across Timor, including Atambua Basin, but cause of extinction unknown. No example yet of direct association of lithic artefacts with extinct Pleistocene faunas: vertebrate fossils from M Pleistocene or earlier, predating oldest records of modern humans on Timor since~45,000 years ago)

Hilgen, S.L., F.J. Hilgen, S. Adhityatama, K.F. Kuiper & J.C.A. Joordens (2022)- Towards an astronomical age model for the Lower to Middle Pleistocene hominin-bearing succession of the Sangiran Dome area on Java, Indonesia. Quaternary Science Reviews 297, 7, 107788, p. 1-19.
(online at: https://www.sciencedirect.com/science/article/pii/S027737912200419X?via%3Dihub)
(Milankovitch cycle control applied to grain size cyclicity observed in Pleistocene of Sangiran and Bapang Fms. Onset of Mid-Pleistocene Transition with age of ~1.2 Ma in Upper Sangiran Fm ('even though multiple tuning options are still possible'. Tuned ages mostly consistent with 'short' paleoanthropological chronology, i.e. ~1.2 or 1.27 Ma for first arrival of Homo erectus)

Hilgen, S.L., E. Pop, S. Adhityatama, T.A. Veldkamp, H.W.K. Berghuis, I. Sutisna, D. Yurnaldi, G. Dupont-Nivet et al. (2023)- Revised age and stratigraphy of the classic *Homo erectus*-bearing succession at Trinil (Java, Indonesia). Quaternary Science Reviews 301, 107098, p. 1-18.
(online at: https://www.sciencedirect.com/science/article/pii/S027737912200539X)
(New chronostratigraphic framework for Trinil site from 40Ar/39Ar, paleomagnetic and luminescence (pIRIR290) dating methods. Two distinct Pleistocene fossiliferous channel fills (1) lower Bone-bearing channel 1 dates to 830-773 ka, (2) Bone-Bearing Channel 2 ~430-450 ka. Also significantly younger T2 terrace deposits. Dubois may have excavated fossils from three fossiliferous units with different ages)

Hocknull S.A., P.J. Piper, G.D. van den Bergh, R. Awe Due, M.J. Morwood & I. Kurniawan (2009)- Dragon's Paradise Lost: Palaeobiogeography, evolution and extinction of the largest-ever terrestrial lizards (Varanidae). PLoS One 4, 9, e7241, p. 1-15.
(online at: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0007241)
(Phyletic giantism and westward dispersal from mainland Australia best explanation for palaeodistribution of V. komodoensis and newly identified species of giant varanid from Timor. Pliocene giant varanid fossils from Australia suggest origin for V. komodoensis on mainland Australia (>3.8 Ma))

Holloway, R.L. (1981)- The Indonesian *Homo erectus* brain endocasts revisited. American J. Physical Anthropology 55, p. 503-521.
(online at: http://www.columbia.edu/~rlh2/IndonHerectAJPA1981.pdf)
(New brain endocasts of Homo erectus discoveries from Java. Mean brain volume 930 ml (Homo soloensis 1000-1250 cm3, modern humans ~1250 cm3; JTvG))

Hooijer, D.A. (1946)- Prehistoric and fossil rhinoceroses from the Malay Archipelago and India. Zoologische Mededelingen 26, 1, p. 1-138.
(online at: www.repository.naturalis.nl/document/150703)
(Description of rhinoceros fossils collected by Dubois in caves of C Sumatra in 1888-1890, Trinil, etc. Reprint of doctoral thesis)

Hooijer, D.A. (1947)- On fossil and prehistoric remains of *Tapirus* from Java, Sumatra and China. Zoologische Mededelingen 27, p. 253-299.
(online at: www.repository.naturalis.nl/document/149527)

Hooijer, D.A. (1947)- *Pithecanthropus, Meganthropus* en *Gigantopithecus*. Geologie en Mijnbouw 9, 12, p. 230-239.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0VWdxUXpJQl9fYlU/view)
(Review of Von Koenigswald (1940) and Weidenreich (1945) monographs on Java hominids)

Hooijer, D.A. (1948)- *Rhinoceros sondaicus* Desmarest from kitchen-middens of Bindjai Tamiang, North Sumatra. Geologie en Mijnbouw 10, 5, p. 115-116.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0alp1ZkJOMzhwQkU/view)
(Rhinoceros tooth from Paleolithic (Late Pleistocene?) refuse-heap at Binjai Tamiang, 15km from mouth of Tamiang River in N Sumatra. Mound composed of layers of Meretrix mollusc shells alternating with ashy layers with stone tools and remains of land mammals, fish and crabs. Evidence of cannibalism. Rhinoceros tooth comparable to material from Sibrambang cave (Padang Highlands))

Hooijer, D.A. (1948)- Prehistoric teeth of man and of the orang-utan from Central Sumatra, with notes on the fossil orang-utan from Java and Southern China. Zoologische Mededelingen, Leiden, 29, p. 175-301.
(online at: http://repository.naturalis.nl/document/150691)
(Study of collection of teeth excavated by Dubois in late 1880's from Lida Ajer and other caves in Padang Highlands, C Sumatra. Includes orang-utan skulls and some hominid teeth indistinguishable from modern humans (may be oldest known Homo sapiens in Indonesian region (~70 ka; Westaway et al. 2017). Also extensive study of Pleistocene orang-utan teeth from Java and S China))

Hooijer, D.A. (1948)- Pleistocene vertebrates from Celebes. I. *Celobochoerus heekereni* nov.gen.nov. spec. Proc. Koninklijke Nederlandse Akademie van Wetenschappen 51, 8, p. 1024-1032.
(First of series of descriptions of Pleistocene mammal fossils from around Tjabenge, S Sulawesi, ~100 km NE of Makassar, collected by Van Heekeren)

Hooijer, D.A. (1948)- Pleistocene vertebrates from Celebes. II. *Testudo margae* nov. spec. Proc. Koninklijke Nederlandse Akademie van Wetenschappen 51, 9, p. 1169-1182.
(Pleistocene giant land tortoise fossils from S Sulawesi)

Hooijer, D.A. (1948)- Pleistocene vertebrates from Celebes. III. *Anoa depressicornis* (Smith) subsp. and *Babyrousa babyrussa beruensis* nov. subsp. Proc. Koninklijke Nederlandse Akademie van Wetenschappen 51, 10, p. 1322-1330.

Hooijer, D.A. (1949)- Pleistocene vertebrates from Celebes. IV. *Archidiskodon celebensis* nov. spec. Zoologische Mededelingen, Leiden, 30, 14, p. 205-226.
(online at: www.repository.naturalis.nl/document/150021)
(Pleistocene dwarf elephant fossils from S Sulawesi)

Hooijer, D.A. (1949)- Voorlopige mededeling over vondsten van Pleistocene gewervelde dieren in Zuid-Celebes. Oudheidkundig Verslag 1941-1947, Oudheidkundige Dienst in Indonesia, Jakarta, p. 111-112.
(online at: https://www.delpher.nl/nl/tijdschriften/view?coll=dts&identifier=MMKITLV3:002517002:00126)
('Preliminary communication about discoveries of Pleistocene vertebrate fossils in South Sulawesi'. Brief communication about Pleistocene fossils discovered by H.R. van Heekeren near Tjabenge, 100km NE of

Makassar. Previously unknown from Sulawesi, and questioning the validity of the zoogeographic 'Wallace Line'. Parts already published by Hooijer in 1948-1949. Three most interesting finds are an endemic Babirusa-like pig, dwarf elephant and large land tortoise. Fauna different from age-equivalent faunas of Java)

Hooijer, D.A. (1949)- The Pleistocene vertebrates of southern Celebes. Chronica Naturae 105, 5, p. 148-150.

Hooijer, D.A. (1950)- The fossil hippopotamidae of Asia, with notes on the Recent species. Zoologische Verhandelingen, Leiden, 8, p. 3-124.
(online at: https://repository.naturalis.nl/pub/317594)
(Review of fossil Hippopotamus from Asia. Three species in Pleistocene of Java, formerly recored as Hippopotamus antiquus, H. simplex and H. namadicus by Von Koenigswald here renamed Hippopotamus sivalensis koenigswaldi (E Pleistocene Djetis fauna, rel. small, Trinil, Mojokerto, Kedung Brubus), H. sivalensis sivajavanicus and H. sivalensis soloensis n.ssp (M-U Pleistocene, descended from H. koenigswaldi)

Hooijer, D.A. (1950)- Man and other mammals from Toalian sites in south-western Celebes. Verhandelingen Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, Afd. Natuurkunde, 2, 46, p. 1-164.
(online at: https://dwc.knaw.nl/DL/publications/PU00011890.pdf)
(Descriptions of Holocene ('Toalian'; Neolithic) subfossil human and mammal bones from of Bola Batoe cave near Badjo Barebo district, 20 km SW of Watampone in Central Bone and ~100 km NE of Makassar in and other SW Sulawesi sites)

Hooijer, D.A. (1950)- Fossil evidence of Australomelanesian migrations in Malaysia? Southwestern Journal of Anthropology 6, 4, p. 416-422.
(Presence of big-teeth humans resembling Australian aboriginals and Melanesians may once lived in Malaya (=Indonesian) region, as evidenced by Wajak man of Java (Dubois 1920) and subfossil man from Gua Lawa Sampung and Bojonegoro as described by Mijsberg (1932). View disputed by Von Koenigswald 1952)

Hooijer, D.A. (1951)- The geological age of *Pithecanthropus, Meganthropus and Gigantopithecus*. American J. Physical Anthropology 9, 3, p. 265-282.
(Questions some of the ages assigned to Von Koenigswald (1934) Pleistocene vertebrate zonation of Cisande, Cijulang, Kali Glagah, Jetis, Trinil and Ngandong faunas, and places most of them in Pleistocene. Distinctions between zones may also not be as clear-cut as assumed)

Hooijer, D.A. (1951)- Pygmy elephant and giant tortoise. The Scientific Monthly 72, 1, p. 3-8.
(On migration of earliest, Pleistocene mammal 'island fauna' from S Sulawesi, probably from Java)

Hooijer, D.A. (1952)- *Palaeoloxodon* cf. *namadicus* (Falconer et Cautley) from Borneo. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B55; 4, p. 395-398.
(Molar from Samarinda region, E Kalimantan, is first record of M Pleistocene Stegodon fauna on Borneo. Resembles M Pleistocene elephant Palaeoloxodon namadicus of S and E Asia. Represents link between Stegodon- Ailuropoda fauna of Java and Indochina)

Hooijer, D.A. (1952)- Fossil mammals faunas and the Plio-Pleistocene boundary in Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B55, p. 436-443.
(Tjidjulang and Kali Glagah mammal assemblages of Java, originally considered to be of Middle-Late Pliocene age by Von Koenigswald, etc., should be assigned to Lower Pleistocene)

Hooijer, D.A. (1953)- Pleistocene vertebrates from Celebes. V. Lower molars of *Archidiskodon celebensis* Hooijer. Zoologische Mededelingen, Leiden, 31, 28, p. 311-318.
(On teeth of Pleistocene dwarf elephant from S Sulawesi)

Hooijer, D.A. (1953)- Pleistocene vertebrates from Celebes. VI. *Stegodon* spec. Zoologische Mededelingen, Leiden, 32, 11, p. 107-112.
(online at: www.repository.naturalis.nl/document/149397)

(On Pleistocene small Stegodon sp. elephantoid from Cabenge area, Sopeng District, S Sulawesi, collected by Van Heekeren)

Hooijer, D.A. (1953)- Pleistocene vertebrates from Celebes. VII. Milk molars and premolars of *Archidiskodon celebensis* Hooijer. Zoologische Mededelingen, Leiden, 32, 20, p. 221-232.
(online at: https://www.repository.naturalis.nl/document/150455)

Hooijer, D.A. (1954)- Pleistocene vertebrates from Celebes. VIII. Dentition and skeleton of *Celebochoerus heekereni* Hooijer. Zoologische Verhandelingen, Leiden, 24, p. 1-46.
(online at: www.repository.naturalis.nl/document/149035)
(More on Pleistocene pig-like suid from Cabenge area, S Sulawesi)

Hooijer, D.A. (1954)- Pleistocene vertebrates from Celebes. IX. Elasmobranchii. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B57, 4, p. 475-485.
(On Pleistocene shark teeth from Tjabenge area, S Sulawesi)

Hooijer, D.A. (1954)- Pleistocene vertebrates from Celebes. X. Testudinata. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B57, 4, p. 486-489.
(On Pleistocene large tortoise fossils from S Sulawesi)

Hooijer, D.A. (1954)- A pygmy *Stegodon* from the Middle Pleistocene of Eastern Java. Zoologische Mededelingen 33, 14, p. 91-102.
(online at: www.repository.naturalis.nl/document/149695)
(Dwarf elephantoid from area N of Djetis and Perning in E Java, collected by Cosijn)

Hooijer, D.A. (1954)- Pleistocene vertebrates from Celebes. XI. Molars and a tusked mandible of *Archidiskodon celebensis* Hooijer. Zoologische Mededelingen, Leiden, 33, 15, p. 104-120.
(online at: www.repository.naturalis.nl/document/149831)
(?E Pleistocene elephantoids from near Cabenge, 100km NE of Makassar, SW Sulawesi))

Hooijer, D.A. (1954)- Crocodilian remains from the Pleistocene of Celebes. Copeia 1954, p. 263-266.

Hooijer, D.A. (1955)- Fossil Proboscidea from the Malay Archipelago and the Punjab. Zoologische Verhandelingen 28, p. 1-146.
(online at: www.repository.naturalis.nl/document/149023)
(On Pleistocene elephant fossils from Java, Sumatra, Sulawesi, India, etc.: Stegodon, Stegolophodon, Archidiskodon, Elephas, etc.))

Hooijer, D.A. (1956)- The lower boundary of the Pleistocene in Java and the age of *Pithecanthropus*. Quaternaria 3, p. 5-10.
(Mammal fossils suggest Tjidjoelang and Kali Glagah faunas of Java are of basal Pleistocene age)

Hooijer, D.A. (1957)- The correlations of fossil mammalian faunas and the Plio-Pleistocene boundary in Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 60, 1, p. 125-128.
(Reiteration of 1952 position that Tjidjulang and Kali Glagah mammal faunas of Java should be assigned to Villafranchian, Early Pleistocene)

Hooijer, D.A. (1957)- Report upon a collection of fossil vertebrates from Ola Bula, Central Ngada, Flores. Berita Madjelis Ilmu Pengetahuan Indonesia, Jakarta, November 1957, p. 6-9.
(Vertebrate remains discovered in 1956 at Ola Bula on Soa Plateau of Flores belong to extinct primitive Asian elephantids Stegodon, similar to S. trigonocephalus Martin from Pleistocene of Java)

Hooijer, D.A. (1957)- A *Stegodon* from Flores. Treubia 24, p. 119-129.
(online at: e-journal.biologi.lipi.go.id/index.php/treubia/article/download/2721/2332)

(Stegodon remains discovered by Verhoeven in 1956 at Ola Bula on Soa Plateau of Flores described as Stegodon trigonocephalus floresiensis. Rel. common Stegodon fossils, but remarkable lack of associated fauna)

Hooijer, D.A. (1957)- Three new giant prehistoric rats from Flores, Lesser Sunda Islands. Zoologische Medelingen 35, p. 229-314.
(online at: www.repository.naturalis.nl/document/150080)
(Three new forms of large Pleistocene rat fossils collected by Verhoeven in cave deposits at Liang Toge near Warukia, Manggarai, W Flores: Papagomys armandvillei besar, P. verhoeveni and Spelaeomys florensis. Associated with Mesolithic flake and blade industry)

Hooijer, D.A. (1958)- Fossil Bovidae from the Malay Archipelago and the Punjab. Zoologische Mededelingen 38, p. 1-110.
(online at: https://repository.naturalis.nl/pub/317642/ZV1958038001.pdf)
(Bovidae (buffaloes, cows, bison, etc.) rel. common in Pleistocene of Java, mainly of genera Bibos, Bubalus, Leptobos, Duboisia, etc.)

Hooijer, D.A. (1958)- The Pleistocene vertebrate fauna of Celebes. Archives Neerlandaises Zoologie 13, Supplement 1, p. 89-96. (also in Asian Perspectives 2, 2, p. 71-76)
(Pleistocene mammal faunas discovered in fluvial deposits of Tjabenge area SW Sulawesi by Van Heekeren in 1948 different from any Pleistocene fauna in Indo-Australian region: island fauna with dwarf elephant and buffalo, giant tortoise, freshwater sharks and rays, etc.)

Hooijer, D.A. (1960)- Quaternary gibbons from the Malay Archipelago. Zoologische Verhandelingen 46, 1, p. 1-42.
(online at: http://repository.naturalis.nl/document/148964)
(Apes (Pongidae) from Indonesian region include Late Pleistocene gibbons (Symphalangus and Hylobates) from limestone caves in Padang Highlands, C Sumatra, collected by Dubois. Possibly also incave material from C Java (Pacitan) and Sarawak)

Hooijer, D.A. (1962)- Quaternary langurs and macaques from the Malay Archipelago. Zoologische Verhandelingen 55, 1, p. 1-64.
(online at: www.repository.naturalis.nl/document/148851)
(Pleistocene monkeys (Presbytis, Trachypithecus, Macaca) from limestone caves in Padang Highlands, C Sumatra, and cave deposits on Java)

Hooijer, D.A. (1962)- Paleontology of hominid deposits in Asia. The Advancement of Science, London, 1962, p. 485-489.

Hooijer, D.A. (1962)- Report upon a collection of Pleistocene mammals from tin-bearing deposits in a limestone Cave near Ipoh, Kinta Valley, Perak. Federation Museums Journal 7 (new series), p. 1-5.
(Study of mammal fossils from depth of ~9 m in tin bearing deposits inside a limestone cave near Ipoh,and sent to Museum of Natural History (Leiden) by B.A.V. Peacock in 1957. 51 specimens from 7 species of mammals, interpreted to be of M Plleistocene age. Includes antelope Duboisia santeng and hippopotamus, known from M Pleistocene of Java, but not recorded yet from mainland SE Asia. Also Java rhinoceros and possible buffalo)

Hooijer, D.A. (1964)- Pleistocene vertebrates from Celebes. XII. Notes on pygmy Stegodonts. Zoologische Mededelingen 40, 7, p. 37-44.
(online at: www.repository.naturalis.nl/document/149873)
(Sulawesi Stegodon decidedly smaller than Stegodon trigonocephalus Martin from Java, but probably similar to newly discovered pygmy stegodonts from Flores. Renamed Stegodon sompoensis n. sp.)

Hooijer, D.A. (1964)- Pleistocene vertebrates from Celebes. XIII. *Sus celebensis* Muller & Schlegel, 1845. Beaufortia 222, 16, p. 215-218.
(online at: www.repository.naturalis.nl/document/548530)

Hooijer, D.A. (1964)- On two milk molars of a pigmy stegodont from Ola Bula, Flores. Bull. Geological Survey Indonesia l, 2, p. 49-52.

Hooijer, D.A. (1964)- New records of mammals from the Middle Pleistocene of Sangiran, Central Java. Zoologische Mededelingen 40, 10, p. 73-87.
(online at: www.repository.naturalis.nl/document/149780)
(On mammal fossils from Sangiran donated to Leiden museum by Van Heekeren and Houboldt. Not much new. Questions Von Koenigswald's perfectly valid suggested use of tektites as dating tool in Sangiran)

Hooijer, D.A. (1965)- Note on *Coryphomis buhleri* Schaub, a gigantic murine rodent from Timor. Israel J. Zoology 14, p. 128-133.
(Large Pleistocene rat fossils from Liang Leluat cave, SW Timor, collected by Verhoeven at Maubesi River)

Hooijer, D.A. (1967)- Indo-Australian insular elephants. Genetica 38, 1, p. 143-162.
(Pleistocene dwarfed elephants known from Celebes, Flores and Timor described and relationships considered. Pygmy forms arose independently on each island as result of isolation and genetic drift favouring small size. Wherever we find pygmy elephants we have also giant rodents)

Hooijer, D.A. (1969)- The *Stegodon* from Timor. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B72, 3, p. 201-210.
(More detailed description of Pleistocene dwarf elephant mandible and molars collected by Verhoeven E of Atambua, W Timor N coast, and first described as Stegodon timorensis by Sartono 1969)

Hooijer, D.A. (1970)- Pleistocene South-East Asiatic pygmy stegodonts. Nature 225, 5231, p. 474-475.

Hooijer, D.A. (1971)- A giant land tortoise, *Geochelone atlas* (Faulconer & Cautley) from the Pleistocene of Timor. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B74, 5, p. 504-525.
(Common material of giant land tortoise from Pleistocene gravel deposits at Raebia in Atambua area, W Timor. Originally described as Testudo margae Hooijer (1948), but indistinguishable from Geochelone atlas from E Pleistocene of Java)

Hooijer, D.A. (1972)- *Stegodon trigonocephalus florensis* Hooijer and *Stegodon timorensis* Sartono from the Pleistocene of Flores and Timor. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B75, p. 12-33.
(Flores and Timor Middle-Late Pleistocene Stegodon elephants much smaller than Java Stegodon, from which they probably evolved)

Hooijer, D.A. (1972)- *Varanus* (Reptilia, Sauria) from the Pleistocene of Timor. Zoologische Mededelingen, Museum Leiden 47, p. 445-448.
(online at: www.repository.naturalis.nl/document/149563)
(On Pleistocene 'komodo dragon'-like lizard fossils from gravel deposits, collected by Verhoeven at Raebia in Atambua area, W Timor)

Hooijer, D.A. (1972)- Pleistocene vertebrates from Celebes. XIV. Additions to the *Archidiskodon-Celebochoerus* fauna. Zoologische Mededelingen 46, 1, p. 1-15.
(online at: www.repository.naturalis.nl/document/150627)
(Descriptions of new mammal material collected by 1970 Dutch-Indonesian expedition to Beru area, Sulawesi. Additional material of Archidiskodon celebensis, Stegodon sompoensis, S. cf. trigonocephalus, Celebochoerus heekerei and Anoa depressicorni)

Hooijer, D.A. (1974)- *Elephas celebensis* (Hooijer) from the Pleistocene of Java. Zoologische Mededelingen 48, 11, p. 85-93.
(online at: www.repository.naturalis.nl/document/150495)

Hooijer, D.A. (1975)- Quaternary mammals west and east of Wallace's line. In: G.J. Bartstra & W.A. Casparie (eds.) Modern Quaternary Research in Southeast Asia, A.A. Balkema, Rotterdam, 1, p. 37-51.
(Introduces concept of hypothetical M Pleistocene 'Stegoland' landmass, comprising Flores, Timor and Sulawesi, characterized by pygmy stegodonts S. sompoensis (based on erroneous assumption that elephants are not good swimmers and therefore unlikely to migrate between islands; JTvG))

Hooijer, D.A. (1975)- Quaternary mammals west and east of Wallace's line. Netherlands J. Zoology. 25, p. 46-56.
(Same paper as above)

Hooijer, D.A. (1981)- What, if anything new, is *Stegodon sumbaensis* Sartono? Modern Quaternary Research in Southeast Asia 6, Balkema, Rotterdam, p. 89-90.
(Nature of Sumba Stegodon described by Sartono (1979) still uncertain. Not necessarily pygmy Stegodon)

Hooijer, D.A. (1982)- The extinct giant land tortoise and the pygmy stegodont of Indonesia. Modern Quaternary Research in Southeast Asia 7, Balkema, Rotterdam, p. 171-176.
(Mainly critical discussion of Sondaar (1981) paper on Geochelone faunas. All known Geochelone tortoises in Indonesia can be grouped in G. atlas and made their way to Indonesian islands by overseas dispersal (not land bridges). All pygmy stegodonts are conspecific, Stegodon sompoensis)

Hooijer, D.A. (1982)- Premolars of *Elephas planifrons* Falconer & Cautley from the Pleistocene of Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B85, 3, p. 265-272.

Hooijer, D.A. (1983)- Comment II: Remarks upon the Dubois collection of fossil mammals from Trinil and Kedungbrubus in Java. Geologie en Mijnbouw 62, p. 337-338.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0TXdLZ3dtNVloTHc/view)
(Comments on De Vos and Sondaar (1982) and De Vos et al. (1982) papers on Dubois collection. Part of problem is that the Dubois collections from Trinil may come from two different deposits: M Pleistocene Kabuh Fm and Late Pleistocene terrace sediments. Also Dubois collection from Kedungbrubus- Gunung Butak is mixture of Late Pliocene- E Pleistocene Jetis and M Pleistocene Trinil fauna)

Hooijer, D.A. (1984)- The mammalian faunas of Trinil and Kedungbrubus in Java once more. Modern Quaternary Research in Southeast Asia 8, Balkema, Rotterdam, p. 95-102.
(Comments on De Vos et al. (1982) and Sondaar et al. (1983) papers. Hooijer does not accept that the more diverse Kedungbrubus fauna is younger than Trinil fauna, but are roughly equivalent)

Hooijer, D.A. & B. Kurten (1984)- Trinil and Kedungbrubus: the *Pithecanthropus*-bearing fossil faunas of Java and their relative age. Annales Zoologici Fennici 21, p. 135-141.
(online at: http://www.sekj.org/PDF/anzf21/anzf21-135-141.pdf
(Authors dispute De Vos (1982) interpretation of Trinil mammal fauna as older than Kedungbrubus fauna)

Hope, G.S. & S.G.Haberle (2005)- The history of the human landscapes of New Guinea. In: A. Pawley et al. (eds.) Papuan pasts: cultural, linguistic and biological histories of Papuan-speaking peoples, Australian National University, Canberra, p. 541-554.
(online at: http://palaeoworks.anu.edu.au/pubs/Hope&Haberle05.pdf)
(Humans have been in highland valleys of New Guinea for at least 30,000 years and presumably occupied savannah plains that then connected New Guinea to Australia for 50,000 years or more)

Houbolt, J.H. (1940)- Bijdrage tot de kennis van der verspreiding van Palaeolithische artefacten in Nederlandsch-Indië. Tijdschrift Indische Taal Land en Volkenkunde 80, p. 614-617.
('Contribution to the knowledge of the distribution of Paleolithic artefacts in the Netherlands Indies')

Hrdlicka, A. (1930)- The skeletal remains of early Man. Smithsonian Miscellaneous Collections 83, p. 1-379.

(online at: https://repository.si.edu/handle/10088/24010)
(Review of fossils of early Man worldwide. Book contains a section on Pithecanthropus fossils of Dubois (which H. was allowed by Dubois to inspect at the Teylers Museum in Haarlem in 1923) and Hrdlicka's visit to the Trinil Site in May 1924. Also describes the 'discovery' of second Pithecanthropus skull cap at Trinil by a Dr. C.E.J. Heberlein from Surabaya in August 1926, which turned out to be a kneecap of a Stegodon elephant)

Huffman, O.F. (1998)- An Early Pleistocene way of life- *Homo erectus* of Sangiran Dome, Central Java. In: P. Lunt, R. Netherwood & O F. Huffman (eds.) Guidebook for field trip to Central Java, Oct 1998, Indonesian Petroleum Association (IPA), Jakarta, p. 1-15.
(online at: www.utexas.edu/cola/files/776909)

Huffman, O.F. (1999)- Pleistocene environmental variety in eastern Java and early *Homo erectus* paleoecology- a geological perspective. Buletin Geologi (ITB) 31, 2, p. 93-107
(Late Pliocene-Pleistocene paleogeography of E Java. Homo erectus homeland was volcanic archipelago)

Huffman, O.F. (2001)- Plio-Pleistocene environmental variety in eastern Java and early *Homo erectus* paleoecology- a geological perspective. In: T. Simanjuntak et al. (eds.) Sangiran: man, culture, and environment in Pleistocene times, Proc. International Colloquium on Sangiran, Solo- Indonesia, Solo 1998, National Research Centre of Archaeology, Yayasan Obor Indonesia, Jakarta, p. 231-256.
(online at: www.utexas.edu/cola/depts/anthropology/projects/huffman/6-SoloProc2001.pdf)
(Same paper as above. Homo erectus' homeland was volcanic archipelago with variety of paleoenvironments like Java today. With Late Pliocene-Pleistocene paleogeography; similar to Huffman (1999))

Huffman, O.F. (2001)- Geologic context and age of the Perning/Mojokerto *Homo erectus*, East Java. J. Human Evolution 40, 4, p. 353-362.
(Perning/Mojokerto Homo erectus from Upper Pucangan Fm in E Kendeng hills of E Java likely to be of latest Pliocene age (1.81 Ma radiometric age of hornblende in associated tuff; Swisher et al.) and reflect early H. erectus occupation of Java (but relation of tuff and Mojokerto skull debated in later literature, e.g. Huffman et al. 2006; JTvG)

Huffman, O.F., A.W.J. Berkhout, P.C.H. Albers, J. de Vos & F. Aziz (2022)- Geology and discovery record of the Trinil *Pithecanthropus erectus* site, Java. PaleoAnthropology 2022, 2, p. 266-326.
(online at: https://paleoanthropology.org/ojs/index.php/paleo/article/view/83/773)
(Extensive documentation of Trinil type locality of Homo erectus and excavations by the Dubois team in 1891-1893, the Selenka Expedition 1907-1908, etc. Associated Pleistocene mammal fossils of extinct 'Trinil Fauna' in Pithecanthropus erectus discovery bonebed ('Hauptknochenschicht') include Axis lydekkeri, Duboisia santeng and Stegodon trigonocephalus. Terrestrial-vertebrate elements overwhelmingly disarticulated and frequently broken, with little indication of fluvial abrasion. No indication of hominin or carnivore involvement in deaths reported, suggesting catastrophic mortality along floodplain of a paleo-river, followed by lahar-flood deposition of gravel-size lithic and biotic materials)

Huffman, O.F., J. de Vos, A.W. Berkhout & F. Aziz (2010)- Provenience reassessment of the 1931-1933 Ngandong *Homo erectus* (Java), confirmation of the Bone-bed origin reported by the discoverers. PaleoAnthropology 2010, p. 1-60.
(online at: www.paleoanthro.org/journal/content/PA20100001.pdf)
(Geologists of Geological Survey of Netherlands Indies unearthed 14 Homo erectus fossils in 1931-1933 from Excavation site I Ngandong. Hominin discoveries and other vertebrate remains from thin, gravelly volcaniclastic stratum near base of fluvial terrace remnant ~20m above Solo River)

Huffman, O.F., P. Shipman, C. Hertler, J. de Vos & F. Aziz (2005)- Historical evidence of the 1936 Mojokerto skull discovery, East Java. J. Human Evolution 48, p. 321-363.
(Extensive review of history of discovery of Mojokerto child skull (Perning 1) in E Java in 1936, from hill-slope outcrop of folded conglomeratic sandstone in Duyfjes' Pucangan Fm on Kedungwaru anticline. Now accepted

as Homo erectus (long believed one of oldest hominids from Java. Radiometric age of 1.81 ± 0.04 Ma from pumice near discovery section by Swisher et al., 1994, but may be from deeper beds; Huffman et al. 2006)

Huffman, O.F. & Y. Zaim (2003)- Mojokerto Delta, East Jawa: paleoenvironment of *Homo modjokertensis*-first results. J. Teknologi Mineral (ITB), 10, 2, p.
(manuscript online at: www.utexas.edu/cola/files/793055)
(Homo modjokertensis remains found in 1936 found in situ in Plio-Pleistocene bedrock at Perning site, SW of Surabaya. With Plio-Pleistocene paleogeographic map. Discovery site formed as fluvial channel on delta plain of ancient Mojokerto Delta)

Huffman, O.F. & Y. Zaim (2003)- Geological age estimate for the Mojokerto child's skull Homo erectus. Comment on Berghuis et al. "The eastern Kendeng Hills (Java, Indonesia) and the hominin-bearing beds of Mojokerto, a re-interpretation" [Quat. Sci. Rev. 295 (2022) 107692]). Quaternary Science Reviews 307, 108058, p. 1-4.
(Disagree with Berghuis et al. (2022) analysis of stratigraphic context of Mojokerto child's skull, unearthed near Perning in 1936)

Huffman, O.F., Y. Zaim, J. Kappelman, D.R. Ruez, J. de Vos, Y. Rizal, F. Aziz & C. Hertler (2006)- Relocation of the 1936 Mojokerto skull discovery site near Perning, East Java. J. Human Evolution 50, p. 431-451.
(Mojokerto skull discovery site was probably in beds 20m higher than ash layer dated as 1.81 Ma by Swisher et al. 1994, 2000)

Husain, M., S. Eko, H.A. Mahfi, W. Sunata & P. Sanyoto (1997)- Magnetostratigrafi daerah Patiayam, Kudus, Jawa Tengah. Proc. 22nd Annual Conv. Indonesian Association Geophysicists (HAGI), Bandung, p.
('Magnetostratigraphy of the Patiayam area, Kudus, C Java'. Hominid fossils in Patiayam area near Top Brunhes normal polarity subchron)

Husson, L., T. Salles, A.E. Lebatard, S. Zerathe, R. Braucher, S. Noerwidi, S. Aribowo et al. (2022)- Javanese *Homo erectus* on the move in SE Asia circa 1.8 Ma. Nature Scientific Reports 12, 19012, p. 1-12.
(online at: https://www.nature.com/articles/s41598-022-23206-9)
(Reappraisals of age dates (including new cosmic nucleids technology; TCN), suggest H. erectus reached Java and dwelled at Sangiran, Java at ~1.8 Ma. Hospitable terrestrial conditions of Sundaland facilitated prior dispersal of hominins to edge of Java, where they waited until Java archipelago emerged from sea and connected to Sundaland. Dispersal of H. erectus across Sundaland occurred over 10s to 100s kyr (N.B: TCN age of Grenzbank here suggested to be 1.78 +/- 0.35 Ma; i.e. much older than the widely accepted age of ~1Ma; JTvG)

Hutterer, K.L. (1983)- Absolute dates for the hominid-bearing deposits in Java: an overview. Asian Perspectives 25, 2, p. 53-65.
(online at: https://scholarspace.manoa.hawaii.edu/bitstream/10125/19247/1/AP-v25n2-53-65.pdf)
(Literature review of published radiometric dates of Upper Pucangan Beds (~0.85-1.2 Ma; with 'Djetis Fauna near top at ~0.8 Ma) and Lower Kabuh Beds (~0.5-0.7 Ma; with Trinil Fauna, ~0.5 Ma) and Notopuro Beds (<0.1 Ma; with Ngandong Fauna). Tektite from Sangiran 730,000 ± 50,000 yrs)

Hyodo, M. (2001)- The Sangiran geomagnetic excursion and its chronological contribution to the Quaternary geology of Java. In: T. Simanjuntak et al. (eds.) Sangiran: man, culture, and environment in Pleistocene times, Proc. International Colloquium on Sangiran, Solo- Indonesia 1998, National Research Centre of Archaeology, Yayasan Obor, Jakarta, p. 320-335.
(Sangiran geomagnetic excursion, characterized by westerly declinations, ranges from below T1 Tuff up to just above diatomite layer in Pucangan Fm. Age estimated 1.56-1.48 Ma. Good time marker for tectonic event in M Matuyama chron in C and E Java)

Hyodo, M., S. Matsu'ura, Y. Kamishima, M. Kondo, Y. Takeshita, I. Kitaba, T. Danhara, F. Aziz, I. Kurniawan & H. Kumai (2011)- High-resolution record of the Matuyama-Brunhes transition constrains the age of Javanese

Homo erectus in the Sangiran dome, Indonesia. Proc. National Academy of Sciences USA (PNAS) 108, 49, p. 19563-19568.
(online at: https://www.pnas.org/doi/pdf/10.1073/pnas.1113106108)
(Paleomagnetic study in Sangiran area. A reverse-to-normal polarity transition in 7m thick section across Upper Tuff in Bapang (= Kabuh) Fm, showing 3 short reversal episodes overlain by thick normal polarity magnetozone, a pattern that closely resembles Matuyama-Brunhes transition (~0.78 Ma). Hominid last occurrence and tektite level in Sangiran nearly coincident, just below Upper Middle Tuff, which underlies MB transition. Meteorite impact preceded M-B reversal by ~12 ka)

Hyodo, M., H. Nakaya, A. Urabe, H. Saegusa, S. Xue, J. Yin & X. Ji (2002)- Paleomagnetic dates of hominid remains from Yuanmou, China, and other Asian sites. J. Human Evolution 43, 1, p. 27-41.
(Geomagnetic data suggest Homo erectus-affinity Yuanmou, SW China, hominid remains from early Brunhes chron ~0·7 Ma). Hominid fossils from Sangiran and Mojokerto, Java, do not exceed 1·1 Ma in age)

Hyodo, M., W. Sunata & E.E. Susanto (1990)- A long-term geomagnetic excursion from Plio-Pleistocene sediments in Java. In: M. Kono (ed.) Rock magnetism and paleogeophysics 17, Tokyo, p. 31-37. *(online at: http://peach.center.ous.ac.jp/rprep/Rock%20Magnetism%20and%20Paleogeophysics%20vol17%201990.pdf)*
(Similar to Hyodo, Sunata & Susanto et al., 1992))

Hyodo, M., W. Sunata & E.E. Susanto (1992)- A long-term geomagnetic excursion from Plio-Pleistocene sediments in Java. J. of Geophysical Research 97, B6, p. 9323-9335.
(Paleomagnetic records from Sangiran and Mojokerto suggest large-scale declination swing between Olduvai and Jamarillo events, lasting ~130,000 years)

Hyodo, M., W. Sunata, E.E. Susanto & H. Wahyono (1988)- Paleomagnetism of Plio-Pleistocene sediments in Sangiran, Central Java. In: M. Kono (ed.) Rock magnetism and paleogeophysics 15, DELP Publ. 23, Tokyo, p. 31-34. *(online at: http://peach.center.ous.ac.jp/rprep/Rock%20magnetism%20and%20Paleogeophysics%20vol15%201988.pdf)*
Nine polarity boundaries were located in outcrops of Sangiran area. The stratigraphic levels with Pithecantropus fossils (uppermost Pucangan- Kabuh Fms) range from lower boundary of Jaramillo event at 0.97 Ma to Brunhes-Matuyama boundary at 0.73 Ma)

Hyodo, M., N. Watanabe, W. Sunata & E.E. Susanto (1993)- Magnetostratigraphy of hominid fossil bearing formations in Sangiran and Mojokerto, Java. Anthropological Science 101, 2, p. 157-186.
(online at: https://www.jstage.jst.go.jp/article/ase1993/101/2/101_2_157/_pdf)
(Also in Rock magnetism and paleogeophysics 19 (1992). Paleomagnetic study of Plio-Pleistocene formation at Sangiran and Mojokerto. Levels of hominid fossils in Sangiran range from lower Jamarilo event at 0.97 Ma to Brunhes-Matayama boundary at 0.73 Ma. Homo modjokertensis in Mojokerto lies at lower border of Jamarillo event at 0.97 Ma)

Indonesia-Japan Research Cooperation Programme (CTA-41) (1979)- Progress report of the Indonesia-Japan joint research project on geology of human fossil bearing formations in Java. Bull. Geological Research Development Centre (GRDC) 1, p. 47-60.

Indonesia-Japan Joint Research Team (1979)- Stratigraphy and geological structure in the Central Part of the Sangiran Dome. Bull. Geological Research Development Centre (GRDC) 2, p. 55-61.
(Brief description of Late Pliocene- Pleistocene stratigraphy exposed in Sangiran Dome, C Java. Structure is dome, with radial and concentric faults. With four mud volcanoes with exotic blocks and some natural gas seepage. Described in more detail in Watanabe & Kadar, 1985)

Indonesia-Japan Joint Study Team (1991)- Quaternary geology of the northern foot area of Mount Lawu and along the middle course of the Solo river, Central and East Java. Geological Research Development Centre (GRDC), Bandung, p. 1-84.

Indonesia-Japan Joint Study Team (1990)- Stratigraphical correlation of the Quaternary system in the Sangiran area and its surroundings, Central Java. United Nations, CCOP, Technical Bulletin 21, p. 117-134.

Indriati, E. (2004)- Indonesian fossil hominid discoveries from 1889 to 2003: catalogue and problems. In: S. Akiyama et al. (eds.) Proc. Fifth and Sixth Symposia on collection building and natural history studies in Asia and the Pacific Rim, National Science Museum, Tokyo, Monograph 24, p. 163-177.
(Indonesian hominid fossil discoveries catalogue in 1975 listed 57 hominids, in 2003 list more than doubled, albeit lacking provenance for some discoveries)

Indriati, E. & S.C. Anton (2008)- Earliest Indonesian facial and dental remains from Sangiran, Java: a description of Sangiran 27. Anthropological Science 116, 3, p. 219-229.
(online at: https://www.jstage.jst.go.jp/article/ase/116/3/116_070814/_pdf)
(Sangiran 27 only known facial skeleton of Homo erectus. Probably from earliest Pleistocene Sangiran (Pucangan) Fm. Highly mineralized. Ar/Ar dates on bracketing tuffs suggest age of 1.58-1.66 Ma. Sangiran 27 best considered member of H. erectus, but more robust morphology than contemporaneous fossils from Georgia and Kenya)

Indriati, E., C.C. Swisher, C. Lepre, R.L. Quinn, R.A. Suriyanto, A.T. Hascaryo, R. Grun, C.S. Feibel et al. (2011)- The age of the 20 meter Solo River terrace, Java, Indonesia and the survival of *Homo erectus* in Asia. PloS One 6, 6, e21562, p. 1-10.
(online at: www.plosone.org/article/info:doi%2F10.1371%2Fjournal.pone.0021562)
(Wide range and conflicting results of radiometric ages for hominid-bearing beds of Solo River 20m terrace deposits in E Java. New radiometric ages suggest older age than currently accepted. Ar/Ar ages from '20m terrace' at Ngandong and Jigar 546 ± 12 ka, but ESR/U-series results 143 ka± 20 ka, possibly indicating leaching of uranium (N.B.: Rizal et al. (2019) suggest most likely age of bone bed is ~117-108 ka))

Ingicco, T. (2010)- Les primates quaternaires de Song Terus (Java Est, Indonesie): implications paleobiogeographiques et archeozoologiques pour l'Asie du Sud-Est. Doct. Thesis Museum National Histoire Naturelle, Paris, p. 1-281.
('The Quaternary primates of Song Terus (East Java, Indonesia): paleobiogeographic and archeozoological implications for SE Asia'. Bones of non-human primates 70% of E Holocene mammal remains in caves in Punung region of SE Java. Facial parts of monkeys found with pre-Neolithic human burials, as noted by Van Heekeren. Primate speciation patterns in islands of SE Asia suggest continuous connections between Sunda Islands through forested terrain during Pleistocene. Tooth micro-wear suggests commensalism between monkeys (especially Trachypithecus auratus) and humans at SongTerus from 9000-5000 years ago)

Ingicco, T. (2012)- Les primates quaternaires de Song Terus (Java Est, Indonesie): implications paleobiogeographiques et archeozoologiques pour l'Asie du Sud-Est. Bulletins Memoires Societe Anthropologie de Paris 24, p. 199-203.
(Summary of thesis above)

Ingicco, T., N. Amano, K. Setiagama, A.M. Moigne, Budiman, A.M. Semah, T. Simanjuntak & F. Semah (2020)- From food to grave good, Nonhuman primate exploitation in Early to Mid-Holocene Eastern Java (Indonesia). Current Anthropology 61, 2, p. 264-276.
(online at: https://www.researchgate.net/publication/340390122_From_Food_to_Grave_Good_Non-human_Primate_Exploitation_in_Early_to_Mid-Holocene_Eastern_Java_Indonesia)
(On exploitation of Cercopithecidae (monkeys) by E Holocene humans, from acquisition through processing, in two cave sites in Gunung Sewu region of E Java: Song Terus and Braholo. Trachypithecus auratus accounts for more than 90% of monkey remains, suggesting deliberate targeting of this over other monkey species. Placement of butchery marks allowed reconstruction of how monkeys were processed (for food and materials for tool and ornament manufacture))

Ingicco, T., J. de Vos & O.F. Huffman (2014)- The oldest gibbon fossil (Hylobatidae) from insular Southeast Asia; evidence from Trinil (East Java, Indonesia), lower/middle Pleistocene. PloS One 9, 6, E99531, p. 1-15.

(online at: http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0099531)
(Fossil femur excavated by Dubois in 1891-1900 in E-M Pleistocene bone bed of Trinil and now in Dubois collection at Naturalis museum, Leiden, recognized here as that of Hylobatidae (gibbon), making it oldest insular record in SE Asia. Living Hylobatidae inhabit evergreen rain forests, so paleoenvironment in greater Trinil area included forests of this kind in E-M Pleistocene)

Ingicco, T., A.M. Moigne, K.F. Setiagama, N. Amano, A. Kusno, A. Mirza, F.S. Detroit, A.M. Semah & F. Semah (2014)- The fauna of Song Terus cave (East Java, Indonesia) and LGM impact on the Sunda shelf: is the Keplek fauna an impoverished Wajak fauna? In: N. Amano et al. (eds.) Southeast Asia: human evolution, dispersals and adaptations, 17th Congress International Union for Prehistoric and Protohistoric Sciences (UISPP), Burgos 2014, p. 110-115.

Ingicco, T., M.C. Reyes, J. de Vos, M. Belarmino, P.C.H. Albers, I. Lipardo X. Gallet, N. Amano, G.D. van den Bergh et al. (2020)- Taphonomy and chronosequence of the 709 ka Kalinga site formation (Luzon Island, Philippines). Nature Scientific Reports 10, 11081, p. 1-17.
(online at: https://www.nature.com/articles/s41598-020-68066-3.pdf)
(Early Middle Pleistocene Kalinga site with 2.5m thick fossiliferous 'unit F' with 73 stone artefacts along with complete butchered rhinoceros skeleton and two tektites, embedded in mudflow within a paleochannel (see also Ingicco et al. 2018). Most likely, the rhino carcass was transported along with the hominid stone artefacts used for butchering by the mudflow)

Ingicco, T., F. Semah, Y. Zhou, A.M. Semah & H. Forestier (2022)- The early lithic productions of Island Southeast Asia: Traditions or convergences? L'Anthropologie 126, 1, 102997, p. 1-23.
(manuscript online at: https://hal.science/hal-03659596v1/file/S0003552122000024.pdf)
(Recent and old excavations produced stone artefacts as old as 1 My on Flores, 800,000 years on Java, 700,000 on Luzon and >118,000 years on Sulawesi, Java Pacitanian, etc. Surface collections dating back to Lower Paleolithic)

Ingicco, T., G. van den Bergh, J. de Vos, A. Castro, N. Amano & A. Bautista (2016)- A new species of *Celebochoerus* (Suidae, Mammalia) from the Philippines and the paleobiogeography of the genus *Celebochoerus* Hooijer, 1948. Geobios 49, 4, p. 285-291.
(online at: https://ro.uow.edu.au/cgi/viewcontent.cgi?article=5182&context=smhpapers)
(Celebochoerus is suid (pig familiy) with large upper tusks, previously known only from Plio-Pleistocene of Sulawesi. New discovery of canine fragment of Celebochoerus from Cagayan Valley of Luzon, N Philippines, described as Celebochoerus cagayanensis nov. sp. Propose migration route from Philippines to Sulawesi, possibly out of Taiwan, independent from better known Pleistocene migration route from India into Java)

Ingicco, T., G.D van den Bergh, C. Jago-On, J. Bahain, M. G.Chacon, N. Amano, H. Forestier, C. King, K. Manalo et al. (2018)- Earliest known hominin activity in the Philippines by 709 thousand years ago. Nature 557 (7704), p. 233-237.
(manuscript online at: https://ro.uow.edu.au/cgi/viewcontent.cgi?article=6441&context=smhpapers)
(Excavations at Kalinga in Cagayan Valley of N. Luzon since 2013 yielded 57 stone-tools associated with 75% complete disarticulated skeleton of Rhinoceros philippinensis, with clear signs of butchery, together with other fossil fauna. All finds originate from clayey bone bed dated to 777k- 631k years ago using electron-spin resonance method of tooth enamel and fluvial quartz. This pushes back proven period of hominid colonization of Philippines by 100,000s of years. Basal Unit F bone bed was channel filled up to 3.25m thick mudflow, which covered bones, along with in situ tektite and 57 stone-tools and sparse fossils of other animals (Geoemydidae, Varanus cf. salvator, Stegodon cf. luzonensis, Cervus))

Insani, H., U.P. Wibowo, E. Setiyabudi & I. Kurniawan (2015)- On variation of extinct Java Hippopotamuses: a note from a new finding of Hippopotamidae fossil from Subang, West Java, Indonesia. Proc. Joint Convention HAGI-IAGI-IAFMI-IATMI, Balikpapan, JCB2015-277, p. 1-4.
(online at: www.researchgate.net/publication/285482685_ON_VARIATION_OF_EXTINCT_JAVA_H Etc.)

(New maxilla fossil of small hippopotamus from conglomeratic sandy layer of E Pleistocene Citalang Fm in Pasir Cabe, Wanareja Village, Subang District. Assigned to Hippopotamus sivalensis sivajavanicus)

Itihara, M., Sudijono, D. Kadar, T. Shibasaki, H. Kumai, S. Yoshikawa, F. Aziz et al. (1985)- Geology and stratigraphy of the Sangiran area. In: N. Watanabe & D. Kadar (eds.) Quaternary geology of the hominid fossil bearing formations in Java. Geological Research Development Centre (GRDC), Bandung, Special Publ. 4, p. 11-43.

Itihara, M., Sudijono, Wikarno & D. Kadar (1985)- Mud volcanoes in the Sangiran Dome. In: N. Watanabe & D. Kadar (eds.) Quaternary geology of the hominid fossil bearing formations in Java. Geological Research Development Centre (GRDC), Bandung, Special Publ. 4, p. 117-124.

Itihara, M., N. Watanabe, D. Kadar & H. Kumai (1994)- Quaternary stratigraphy of the hominid fossil bearing formations in the Sangiran area, Central Java. In: J.L. Franzen (ed.) 4th International Conference 100 years of *Pithecanthropus*. The *Homo erectus* problem, Frankfurt 1991, Courier Forschungsinstitut Senckenberg 171, p. 123-128.

Itihara, M., Wikarno & Y. Kagemori (1985)- Tektites from the Sangiran area. In: N. Watanabe & D. Kadar (eds.) Quaternary geology of the hominid fossil bearing formations in Java. Geological Research Development Centre (GRDC), Bandung, Special Publ. 4, p. 125-128.
(Horizon with tektites, glassy material from widespread M Pleistocene Australasian meteorite strewnfield, present between 2nd and 3rd Tuff Horizon in Bapang/ Kabuh Fm of Sangiran Dome)

Irwanto, D. (2019)- Sundaland. Tracing the cradle of civilizations. Indonesia Hydro Media, Bogor, p. 1-386.
(Paleogeography of Sundaland during the Late Pleistocene Last Glacial Maximum, and its role in the dispersal of modern humans across SE Asia)

Jablonski, N.G. & D.E. Tyler (1999)- *Trachypithecus auratus sangiranensis*, a new fossil monkey from Sangiran, Central Java, Indonesia. International J. of Primatology 20, 3, p. 319-326.
(Description of new subspecies of Javan lutung monkey, based on tooth-bearing upper jaw fragment from volcanic breccia between U Kalibeng Fm and Lower Pucangan Fm, 500m S of Sangiran. Geochronological age 1.9 Ma, making it oldest monkey in SE Asia. Morphologic similarities to living leaf monkeys of Java, but larger (age disputed by Larick et al. 2000; probably younger))

Jacob, T. (1964)- A new hominid skull cap from Pleistocene Sangiran. Anthropologica, n.s. 6, p. 97-104.
(After two decades of slowdown in paleoanthropolical research in Java since start of the Pacific War, a new joint Paleoantropological Research Project of Gadjah Mada University/ Archeological Survey and Geological Survey uncovered new Pithecanthropus skull fragments in 1963, in MPleistocene upper Kabuh Fm ('Trinil Beds'). Possibly belog to same individual as fragments found at same locality earlier by Sartono (1963))

Jacob, T. (1966)- The sixth skull cap of *Pithecanthropus erectus*. American J. Physical Anthropology 25, 3, p. 243-259.
(Sixth Pithecanthropus skull (named skull V) from cross-bedded sandstone of upper Trinil beds in Tanjung village, Sangiran, C Java. Pithecanthropine characteristics, with cranial capacity ~975 cm3. Absence of cranial base does not necessarily indicate that specimen was victim of cannibalism)

Jacob, T. (1967)- Some problems pertaining to the racial history of the Indonesian Region. Doct. Thesis, Rijksuniversiteit Utrecht, p. 1-156. *(Unpublished)*

Jacob, T. (1967)- Recent *Pithecanthropus* finds in Indonesia. Current Anthropology 8, 5, p. 501-504.
(New mandible and skull caps ('Skull VI' and 'Skull VII') of Pithecanthropus from Sangiran, C Java)

Jacob, T. (1972)- The absolute date of the Djetis beds at Modjokerto. Antiquity 46, p. 148.

(online at: https://www.cambridge.org/core/journals/antiquity/article/abs/absolute-date-of-the-djetis-beds-at-modjokerto/EBD86F3E21192C35FAEB8291A1E97C0B)
(Brief note on K-Ar radiometric age of pumice tuff from a few m above horizon with 'Mojokerto child' at Perning, by G.H. Curtis from University of California gave age of 1.9 Ma, with range of +/- 0.4 Ma)

Jacob, T. (1972)- The problem of head-hunting and brain-eating among Pleistocene men in Indonesia. Archaeology & Physical Anthropology in Oceania 7, 2, p. 81-91.
(Many indications, but not enough convincing evidence that Pleistocene men in Java were cannibals)

Jacob, T. (1973)- Palaeoanthropological discoveries in Indonesia with special reference to the finds of the last two decades. J. Human Evolution 2, 6, p. 473-485.
(Reviews of paleoanthropological research in Indonesia since 1889. Three periods, with most finds in second one (1931-1941). Most finds are skull fragments of Pithecanthropus erectus, from M Pleistocene Kabuh Fm in Sangiran. K/Ar dating gives age of 1.9 ± 0·4 Million years for Jetis beds at Perning (site of Mojokerto juvenile calvaria) and 0.83 Ma for (upper?) tuff from Trinil beds (Kabuh Fm) at Sangiran)

Jacob, T. (1974)- Studies on human variation In Indonesia. J. National Medical Association 66, 5, p. 389-399.
(online at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2609252/pdf/jnma00489-0061.pdf)

Jacob, T. (1975)- Morphology and paleoecology of early man in Java. In: R.H. Tuttle (ed.) Paleoanthropology morphology and paleoecology, Mouton, The Hague, p. 311-325.
(Between 1889-1941 at least 31 hominid fossils discovered in Indonesia, mainly pithecanthropines. Since 1952 >17 additional individuals found)

Jacob, T. (1975)- The pithecanthropines of Indonesia. Bulletins Memoires Societe Anthropologie de Paris 13, 2, p. 243-256.
(online at: www.persee.fr/web/revues/home/prescript/article/bmsap_0037-8984_1975_num_2_3_1816)
(Bodily remains of Pithecanthropus of Java consist of cranial and lower limb bones of ~50 individuals, from Lower and Middle Pleistocene beds (~1.9 to 0.2 Ma). Hiatus of at least 150,000 years existed between last Pithecanthropus and first Homo remains in Asia)

Jacob, T. (1976)- Man in Indonesia: past, present and future. Modern Quaternary Research in Southeast Asia 2, Balkema, Rotterdam, p. 39-48.
(Brief review of hominids in Indonesian region in last 2 Myrs)

Jacob, T. (1977)- Evolution of man in Southeast Asia. Berkala Ilmu Kedokteran (J. Medical Sciences) 9, 4, p. 175-186.
(online at: https://journal.ugm.ac.id/bik/article/download/4724/3981)

Jacob, T. (1978)- The puzzle of Solo Man. Modern Quaternary Research in Southeast Asia 4, p. 31-40.
(Solo man from Ngandong first discovered in 1931 is M Pleistocene pithecanthropine (advanced Homo erectus). Solo man made stone tools. May have been cannibalistic, but not necessarily so)

Jacob, T. (1978)- New finds of Lower and Middle Pleistocene hominines from Indonesia. In: F. Ikawa-Smith (ed.) Early Palaeolithic in South and East Asia, Mouton Publishers, The Hague, p. 13-22.
(Brief review of occurrences of Pleistocene hominid fossils of Java. Most new finds of M Pleistocene H. erectus fragments from Kabuh Fm of Sangiran area. K/Ar ages of Kabuh Fm hominid-bearing beds average 830,000 years; associated tektites 710,000 years)

Jacob, T. (1979)- Hominine evolution in South East Asia. Archaeology Physical Anthropology Oceania 14, 1, p. 1-10.

Jacob, T. (1980)- The *Pithecanthropus* in Indonesia: phenotype, genetics and ecology. In L.K. Konigsson (ed.) Current argument on Early Man, Proc. Nobel Symposium, Karlskoga 1978, Pergamon Press, Oxford, p. 170-179.

Jacob, T. (1981)- Solo Man and Peking Man. In: B.A. Sigmon & J.S. Cybulski (eds.) *Homo erectus*: papers in honor of Davidson Black, University of Toronto Press, Toronto, p. 87-104.

Jacob, T. (1981)- *Meganthropus, Pithecanthropus* and *Homo sapiens* in Indonesia: Evidence and problems. In: D. Ferembach (ed.) Les processus de l'hominisation, L'evolution humaine, les faits, les modalites, Colloque 599, CNRS Editions, Paris, p 81-84.

Jacob, T. (1983)- Early Man in Indonesia: the "defossilization" of human fossils. In: A. Bryan (ed.) Symposium on Human evolution, University of Alberta, Edmonton, 1982, Canadian J. Anthropology 3, 2, p. 191-195.
(online at: https://tspace.library.utoronto.ca/bitstream/1807/4410/1/CJA%20Vol%203%20%232%201983.pdf)
(Brief review of three genera of fossil hominids from 7 sites on Java)

Jacob, T. (1984)- The fossil skull cap from Sambungmacan and its implication to human evolution. Berkala Bioanthropologi Indonesia (Indonesian Journal of Bioanthropology) 5, p. 19-27.

Jacob, T., E. Indriati, R.P. Soejono, K. Hsu, D.W. Frayer, R.B. Eckhardt, A. J. Kuperavage, A. Thorne & M. Henneberg (2006)- Pygmoid Australomelanesian *Homo sapiens* skeletal remains from Liang Bua, Flores: population affinities and pathological abnormalities. Proc. National Academy of Sciences USA (PNAS) 103, 36, p. 13421-13426.
(online at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1552106/pdf/zpq13421.pdf)
(Liang Bua 1 skull descrined as new species Homo floresiensis by Brown et al. (2004) and Morwood et al. (2005) not new species, but here placed within modern human ranges of variation, resembling Australomelanesian populations. LB1 probably drawn from earlier pygmy H. sapiens population)

Jacob, T. & D. Kadar (1978)- A new pithecanthropine cranial endocast S34 from the Sangiran Dome area, Central Java. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 1, p. 1-7.

Jacob, T., R.P. Soejono, L.G. Freeman & F.H. Brown (1978)- Stone tools from Mid-Pleistocene sediments in Java. Science 202, 4370, p. 885-887.
(Stone chopper and retouched flake from mid-Pleistocene channel fills at Sambungmacan, C Java)

Jaekel, O. (1911)- Die fossilen Schildkrotenreste von Trinil. In: M.Selenka & M. Blanckenhorn (eds.) Die *Pithecanthropus*-Schichten auf Java, Geologische und Palaontologische Ergebnisse der Trinil Expedition (1907 und 1908), Wilhelm Engelmann, Leipzig, p. 75-81.
('The fossil turtle remains from Trinil'. Description of Pleistocene turtle fossils from Trinil beds, collected by Selenka expedition. High degree of fragmentation of thick tortoise shells suggest these are food remains of Pithecanthropus erectus)

Janensch, W. (1911)- Die Reptilienreste (exkl. Schildkroten). In: M. Selenka & M. Blanckenhorn (eds.) Die *Pithecanthropus*-Schichten auf Java, Geologische und Palaontologische Ergebnisse der Trinil Expedition (1907 und 1908), Wilhelm Engelmann, Leipzig, p. 61-74.
('The reptilian remains (excluding turtles) of Trinil, C Java. Descriptions of Gavialis bengawanicus Dubois, Crocodilus ossifragus and varanus vertebrae)

Janssen, R. (2017)- Isotope records in vertebrate fossils: from the Cretaceous seas to Quaternary Sundaland. Ph.D. Thesis Vrije Universiteit, Amsterdam, p. 1-142.
(online at: https://research.vu.nl/en/publications/isotope-records-in-vertebrate-fossils-from-cretaceous-seas-to-qua)
(Includes chapter on Carbon and Sr isotopes of teeth enamel of Pleistocene mammals from Trinil-Sangiran, C Java, and Padang Highlands))

Janssen, R., J.C.A. Joordens, D.S. Koutamanis, M.R. Puspaningrum, J. de Vos, J.H.J.L. van der Lubbe, J.J.G. Reijmer et al. (2016)- Tooth enamel stable isotopes of Holocene and Pleistocene fossil fauna reveal glacial and interglacial paleoenvironments of hominins in Indonesia. Quaternary Science Reviews 144, p. 145-154.
(online at: https://research.vu.nl/ws/portalfiles/portal/41930773)
(Carbon and oxygen isotopes composition of tooth enamel used to investigate diet and habitat of bovids, cervids and suids from Holocene and Pleistocene sites on Java and Sumatra. Data from Homo erectus bone samples possibly contaminated by diagenetic overprint. C4-dominated isotope signal suggests Trinil specimens in Dubois and Selenka collections were excavated from narrow stratigraphical interval representing dry, glacial climate state (similar in Sangiran))

Jatmiko (2023)- Chronology of lithic culture in Prehistoric at Flores: Case study on Liang Bua site and Soa Basin. In: M. Hasyim (Ed.), Proc. 3rd International Conference on Linguistics and Cultural (ICLC 2022), Makassar, Advances in Social Science, Education and Humanities Research 756, Atlantis Press, p. 618-630.
(online at: https://www.atlantis-press.com/article/125988200.pdf)
(Rel. old (M Pleistocene) stone artefacts from Liang Bua and So'a Basin in Flores)

Jatmiko, G. van den Bergh, M.J. Morwood & I. Kurniawan (2009)- Excavations at Kobatuwa, central Flores, Indonesia. In: F. Aziz et al. (eds.) Pleistocene geology, palaeontology and archaeology of the Pleistocene Soa Basin, Central Flores, Indonesia, Pusat Survei Geologi, Bandung, Special Publ. 36, 6, p. 105-117. *(online at: www.researchgate.net/publication/288965361_6_Excavations_at_Kobatuwa_central_Flores_Indonesia)*
(Pleistocene Stegodon elephantoid molar fragments and stone chopper artefacts from excavations at Kobatua)

Joordens, J.C.A., S. Adhityatama, Y. Dida, T. Reimann, D. R.Ekowati, F. Huffman, B. Didit et al. (2017)- Dating Trinil: towards establishing an age framework for the hominin-bearing deposits at the *Homo erectus* site Trinil (Indonesia). 19th EGU General Assembly, EGU2017, Vienna, p. 11546. *(Abstract only)*
(Dating of Javanese hominin sites is notoriously difficult. No new information in abstract)

Joordens, J.C.A., F. d'Errico, F. Wesselingh, S. Munro, J. de Vos, J. Wallinga, C. Ankjaergaard et al. (2015)- *Homo erectus* at Trinil on Java used shells for tool production and engraving. Nature 518, 7538, p. 228-231.
(online at: www.researchgate.net/publication/269102248_Homo_Erectus_at_Trinil_on_Java_Used_She Etc.)
(Fossil freshwater mussel shells from 'Hauptknochenschicht' of Trinil, C Java (= main bone layer; type locality of Homo erectus) with evidence for freshwater shellfish consumption by hominins, one shell tool and shell with geometric engraving. Sediment in shells dated with 40Ar/39Ar and luminescence dating methods as between ~0.54 and 0.43 Ma, i.e. younger than previously estimated from correlation of mammal assemblages to nearby Sangiran section. Engraving presumably made by Homo erectus)

Joordens, J.C.A., F.P. Wesselingh, J. de Vos, H.B. Vonhof & D. Kroon (2009)- Relevance of aquatic environments for hominins: a case study from Trinil (Java, Indonesia). J. Human Evolution 57, p. 656-671.
(online at: http://www.rhinoresourcecenter.com/pdf_files/131/1313207918.pdf)
(Study of ecological context of Java hominids. Homo erectus site of Trinil contained near-coastal rivers, lakes, swamp forests, lagoons, and marshes with minor marine influence, laterally grading into grasslands. Trinil HK environments yielded edible molluscs and fish. Midden-like characteristics of large bivalve shell assemblages from Trinil HK indicate deliberate collection, possibly by hominins)

Jungers, W. (2013)- *Homo floresiensis*. In: D.R. Begun (ed.) A companion to Paleoanthropology, Chapter 30, p. 582-598.

Jungers, W., S.G. Larson, W. Harcourt-Smith, M.J. Morwood, T. Sutikna, Rokhus Due Awe & T. Djubiantono (2009)- Descriptions of the lower limb skeleton of *Homo floresiensis*. J. Human Evolution 57, 5, p. 538-554.

Junghuhn, F. (1857)- Over de fossiele zoogdierbeenderen te Patihajam, in de residentie Djapara, eiland Java. Natuurkundig Tijdschrift voor Nederlandsch-Indie 14, p. 215-219.

('On the fossil mammal bones at Pati Ayam in the Jepara Residency, Java'. Short note on June 1856 visit to reported 'battlefield of giants' at Pati Ayam hills S of Muiah volcano. Area with black clays rich in mammal bones, including Bos sp., Elephas primigenius and Mastodon elephantoides. Latter species also known from Irrawaddy, India. No figures)

Kadar, A.P. (1994)- A review of the Sangiran (Central Java) Plio-Pleistocene environment from marine and non-marine floras and faunas. In: J.L. Rau (ed.) Proc. 29th Session Committee Co-ord. Joint Prospecting Mineral Resources in Asian Offshore Areas (CCOP), Hanoi 1992, 2, p. 51-60.
(Discussion of paleoenvironments and fossils of Late Pliocene- Pleistocene formations of Sangiran Dome, C Java. Mainly brackish water and fluvial deposits)

Kaharudin, H.A.F., Alifah, L. Ramadhan & S. Kealy (2020)- A review of archaeological dating efforts at cave and rockshelter sites in the Indonesian Archipelago. J. Indo-Pacific Archaeology 44, p. 80-112.
(online at: https://journals.lib.washington.edu/index.php/JIPA/article/view/15659)
(Listings of 99 Pleistocene- Holocene archeological cave and rockshelter sites across Indonesia with published absolute dates)

Kaharudin, H.A.F., G.A.R. Ananda, W.H. Prasetya, M.W. Wibisono & J.S.E. Yuwono (2023)- Hunter-gatherers in labyrinth karst: An Early Holocene record from Gunung Sewu, Java. Archaeological Research in Asia 33, 100427, p.
(Evidence of occupation at Gua Mandung site in Southern Mountains (Gunung Sewu) since at least 11,073 cal BP. With abundant terrestrial fauna remains alongside numerous bone tools. "Van Stein Callenfels is the earliest known archaeologist who recognised the potential of the region in 1927. He identified a substantial lithic industry at the site of Punung in Pacitan Regency that was later called the Pacitanian Culture")

Kaharudin, H.A.F., S. O'Connor, S. Kealey & N. Ririmasse (2024)- Islands on the edge: 42,000-year-old occupation of the Tanimbar islands and its implications for the Sunda-Sahul early human migration discourse. Quaternary Science Reviews 338, 108834, p. 4-13.
(online at: https://www.sciencedirect.com/science/article/pii/S0277379124003354)
(Evidence from NE Tanimbar Islands shows human occupation 42,000 years ago. one of oldest records from Wallacea. Early occupants on Tanimbar had access to medium-sized terrestrial mammals, specifically macropods (marsupials- kangaroos), which are now extinct on island)

Kahlke, H.D. (1972)- A review of the Pleistocene history of the Orang-Utan (*Pongo* Lacepede 1799). Asian Perspectives 15, p. 5-14.
(online at: https://scholarspace.manoa.hawaii.edu/server/api/core/bitstreams/d3fd521d-dc49-4940-82ce-9bb0e9fe4bf2/content)
(Review of Pleistocene orang-utan localities in SE Asia (Java, Sumatra, Kalimantan, China, Laos, Vietnam))

Kaifu, Y. (2006)- Advanced dental reduction in Javanese *Homo erectus*. Anthropological Science 114, 1, p. 35-43.
(online at: https://www.jstage.jst.go.jp/article/ase/114/1/114_1_35/_pdf/-char/ja)
(Postcanine tooth crowns of late E Pleistocene Homo erectus from Sangiran smaller than those of older H. erectus remains of same region. Javanese H. erectus still robust root systems, presumably primitive retention)

Kaifu, Y. (2017)- Archaic hominin populations in Asia before the arrival of modern humans, their phylogeny and implications for the "Southern Denisovans". Current Anthropology 58, Suppl. 17, p. S418-S433.
(online at: http://www.journals.uchicago.edu/doi/pdfplus/10.1086/694318)
(Asian hominid fossil record scant, but suggests presence of regionally different evolutionary lineages of archaic Homo in Pleistocene Asia. Javanese Homo erectus may be 'southern Denisovans')

Kaifu, Y., J. Arif, K. Yokoyama, H. Baba, E. Suparka & H. Gunawan (2007)- A new *Homo erectus* molar from Sangiran. J. Human Evolution 52, 2, p. 222-226.
(online at: https://www.researchgate.net/publication/6673783_A_new_Homo_erectus_molar_from_Sangiran)

(Njg 2005.05 molar originally from Bapang-AG levels of Sangiran region)

Kaifu, Y., F. Aziz & H. Baba (2001)- New evidence of the existence of *Pongo* in the Early/Middle Pleistocene of Java. In: Towards ahead: geological museum in a changing world, Int. Symposium on Geological Museum, Geological Research Development Centre (GRDC), Bandung, Special Publ. 27, p. 55-60.

Kaifu, Y., F. Aziz & H. Baba (2005)- Hominid mandibular remains from Sangiran: 1952-1986 collection. American J. Physical Anthropology 128, 3, p. 497-519.
(Descriptions of 8 hominid mandibular and associated dental remains found between 1952 and 1986 in E Pleistocene deposits of Sangiran, C Java. All specimens are surface finds)

Kaifu, Y., F. Aziz & H. Baba (2013)- The origins and early evolution of Indonesian *Homo erectus*: evidence from Sangiran. In: F. Aziz & H. Baba (eds.) *Homo erectus* in Indonesian- Recent progress of the study and current understanding, Centre for Geological Survey, Bandung, Special Publ., Chapter 3, p. 43-64.
(Significant morphological variation and changes within early Homo erectus from Java)

Kaifu, Y., F. Aziz, E. Indriati, T. Jacob, I. Kurniawan & H. Baba (2008)- Cranial morphology of Javanese *Homo erectus*: new evidence for continuous evolution, specialization, and terminal extinction. J. Human Evolution 55, 4, p. 551-580.
(Morphological changes in H. erectus skulls from Java: brain size expansion, anteroposterior lengthening of midcranial base and an anterior shift of posterior temporal muscle, etc. Crania from Sambungmacan transitional between earlier (Bapang Fm above Grenzbank in Sangiran) and later (Ngandong) morphotypes of Java. Development of unique features in later Javanese H. erectus supports hypothesis that this Javanese lineage went extinct without making significant contributions to ancestry of modern humans)

Kaifu, Y., H. Baba & F. Aziz (2006)- Indonesian *Homo erectus* and modern human origins in Australasia; new evidence from the Sambungmacan region, central Java. In: Y. Tomida et al. (eds.) Proc. 7th and 8th Symposium Collection building and natural history studies in Asia and the Pacific Rim, National Science Museum, Tokyo, Monograph 34, p. 289-294.
(online at: https://www.kahaku.go.jp/research/researcher/papers/28135.pdf)
(Fossil and genetic studies generally support African origins of modern humans. Current fossil evidence still insufficient to reject competing claim of continuity between Javanese Homo erectus and modern aboriginal Australians. New H. erectus fossils from Sambungmacan, C Java, intermediate between earlier and later groups of Javanese H. erectus and support discontinuity between H. erectus and H. sapiens in Australasia)

Kaifu, Y., H. Baba & F. Aziz (2013)- Conclusions: current understanding of Indonesian *Homo erectus*. In: F. Aziz & H. Baba (eds.) *Homo erectus* in Indonesian- Recent progress of the study and current understanding, Centre for Geological Survey, Bandung, Special Publ., chapter 6, p. 103-106.
(Earliest Indonesian hominins (Sangiran Grenzbank, Mojokerto, Trinil?) more primimitive than generally thought. Within global hominin context most comparable to Homo ergaster (~1.7-1.5 Ma). Younger E Pleistocene forms (Homo erectus/Bapang-AG) with evolving characteristics towards Late Pleistocene H. erectus/Ngandong (formerly Homo soloensis). Homo erectus unlikely ancestor of Homo sapiens)

Kaifu, Y., H. Baba, F. Aziz, E. Indriati, F. Schrenk & T. Jacob (2003)- Taxonomic affinities and evolutionary history of the early Pleistocene hominids of Java: dentognathic evidence. American J. Physical Anthropology 128, 4, p. 709-726.
(Study of teeth of E Pleistocene Javanese hominids from Sangiran show morphological differences between younger and older groups. Primitive aspects of oldest Javanese hominids suggest hominids dispersed into E Eurasia in earlier Early Pleistocene)

Kaifu, Y., H. Baba, T. Sutikna, M.J. Morwood, D. Kubo, E.W. Saptomo, Jatmiko, R. Due Awe & T. Djubiantono (2011)- Craniofacial morphology of *Homo floresiensis*: description, taxonomic affinities, and evolutionary implication. J. Human Evolution 61, p. 644-682.

(Description of LB1/1 Homo floresiensis cranium. Reductive trend in facial skeleton comparable to H. sapiens, but craniometrically different. LB1 most similar to older Homo erectus from Sangiran and Trinil, consistent with hypothesis that H. floresiensis evolved from early Javanese H. erectus with dramatic island dwarfism)

Kaifu, Y., E. Indriati, F. Aziz, I. Kurniawan & H. Baba (2010)- Cranial morphology and variation of the earliest Indonesian hominids. In: C.J. Norton & D.R. Braun (eds.) Asian Paleoanthropology: from Africa to China and beyond, Springer Science, Chapter 11, p. 143-157.
(Previous arguments suggest oldest Indonesian/ Sangiran hominids characterized by cranial robusticity, but hominids highly variable, with both robust and gracile morphotypes. Cranial size, shape and dentognathic morphology of earliest Indonesian hominids comparable to ~1.7 Ma early Homo erectus from E Africa)

Kaifu, Y., M. Izuho & T. Goebel (2015)- Modern human dispersal and behavior in Palaeolithic Asia: summary and discussion. In: Y. Kaifu et al. (eds.) Emergence and diversity of modern human behavior in Paleolithic Asia, Texas A&M University Press, College Station, p. 535-566.

Kaifu, Y., R.T. Kono, T. Sutikna, E.W. Saptomo, Jatmiko & R. Due Awe (2015)- Unique dental morphology of *Homo floresiensis* and its evolutionary implications. PLoS One 10, 11, e0141614, p. 1-27.
(online at: http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0141614)
(Dental remains of Homo floresiensis both primitive and advanced molar morphologies, a combination unknown in other hominin species. Consistent with alternative hypothesis that H. floresiensis derived from earlier Asian Homo erectus population and experienced size dwarfism in isolated insular setting)

Kaifu, Y., I. Kurniawan, D. Kubo, E. Sudiyabudi, G.P. Putro, E. Prasanti, F. Aziz & H. Baba (2015)- *Homo erectus* calvaria from Ngawi (Java) and its evolutionary implications. Anthropological Science 123, 3, p. 161-176.
(online at: https://www.jstage.jst.go.jp/article/ase/123/3/123_150702/_pdf)
(Endocranial volume of Ngawi 1 skull 959 cm3. Homo erectus crania from Ngawi and Sambungmacan generally similar to those of U Pleistocene Ngandong H. erectus, but also features close to terminal Pleistocene Bapang AG Homo erectus from Sangiran)

Kaifu, Y., I. Kurniawan, S. Mizushima, J. Sawada, M. Lague, R. Setiawan, I. Sutisna, U.P. Wibowo et al. (2024)- Early evolution of small body size in *Homo floresiensis*. Nature Communications 15, 6381, p. 1-13.
(online at: https://www.nature.com/articles/s41467-024-50649-7)
(Additional hominin fossils Mata Menge shorter and thinner than type specimen of H. floresiensis dated to ~60,000 years ago, and smaller than any other Plio-Pleistocene adult hominin hitherto reported. Teeth also exceptionally small and similar to early Javanese H. erectus. H. floresiensis lineage most likely evolved from early Asian H. erectus and was long-lasting lineage on Flores since at least ~700,000 years ago.

Kaifu, Y., I. Kurniawan, D. Yurnaldi, R. Setiawan, E. Setiyabudi, H. Insani, M. Takai, Y. Nishioka et al. (2022)- Modern human teeth unearthed from below the ~128,000-year-old level at Punung, Java: a case highlighting the problem of recent intrusion in cave sediments. J. Human Evolution 163, p. 103-122.
(Late Pleistocene paleontological site of Punung III, East Java, demonstrates how unexpected intrusion of recent human remains into older stratigraphic levels could occur in cave sediments)

Kaifu, Y., E. Setiyabudi, I. Kurniawan, H. Baba & F. Aziz (2013)- Evolution of Indonesian *Homo erectus* in the Early Pleistocene: significance of Sangiran 17. In: F. Aziz & H. Baba (eds.) *Homo erectus* in Indonesian-Recent progress of the study and current understanding, Centre for Geological Survey, Bandung, Special Publ., chapter 4, p. 65-91.
(Sangiran 17, discovered in situ in Bapang Fm in Sangiran in 1969, is best-preserved cranial specimen of Homo erectus. Some of its features transitional to younger Ngandong Homo soloensis, most others similar to Sangiran Bapang AG Homo erectus group)

Kaifu, Y., Y. Zaim, H. Baba, I. Kurniawan, D. Kubo, Y. Rizal, J. Arif & F. Aziz (2011)- New reconstruction and morphological description of a *Homo erectus* cranium: skull IX (Tjg-1993.05) from Sangiran, Central Java. J. Human Evolution 61, 3, p. 270-294.
(New reconstruction of remarkably complete skull of E Pleistocene Homo erectus skull IX (Tjg-1993.05) from Bapang Fm-AG levels in Sangiran (second best preserved cranium of H. erectus ever). Endocranial volume measured as 870 cc. Neurocranium slightly smaller than smallest cranium found earlier, suggesting individual was female. Numerous similarities to other Bapang-AG Homo erectus specimens)

Karl, H.V. (1983)- Die fossilen Schildkroten von Jawa in den Sammlungen des Paläontologischen Museums im Museum fur Naturkunde Berlin. Unpublished Report, Museum fur Naturkunde; Berlin, p.
('The fossil turtles/tortoises from Java in the collections of the Paleontological Museum in the Museum of Natural History Berlin' (presumably Pleistocene fauna from the Selenka collection))

Karl, H.V. (1987)- Revision der fossilen Schildkroten (Reptilia, Testudines) von Jawa. Abhandlungen Berichte des Museums der Natur Gotha 14, p. 37-44.
('Revision of the fossil turtles/tortoises (Reptilia, Testudines) from Java' (from the Selenka Trinil collection, studied earlier by Jaekel (1911)?))

Karl, H.V., A. Safi & H.D. Philippen (2019)- Evidences of cheloniophagy by early hominid (*Homo erectus*) during middle of Pleistocene from beds of Trinil's layers in central Java (Indonesia), with an updated list of Trinil's testudines, and a redescription of *Duboisemys isoclina* (Dubois, 1908). International J. of Zoology Studies 4, 5, p. 73-84.
(online at: www.researchgate.net/profile/Hans-Volker-Karl/publication/338513109_Evidences_of_che Etc.)
(In-situ archaeological assemblage from Trinil Beds in C Java provides unique combination of faunal remains and evidences of butchery of M Pleistocene tortoise remnants, suggesting Homo erectus predated and butchered terrestrial and aquatic testudines (as already suggested by O. Jaekel in 1911). With review of river turtles/ tortoises from Trinil Beds: Chitra chitra javanensis, Pelochelys cantorii, Amyda cartilaginea, Batagur affinis affinis, Orlitia borneensis, Duboisemys (n.gen.) isoclina (Dubois 1908) and Megalochelys sp.)

Kapid, R., J. Arif & D.E. Irawan (2016)- A review on paleoenvironment suitability for hominid fossils and other early vertebrate faunas: a case from Pucangan and Kabuh Formations, Central and East Java, Indonesia. ScienceOpen Research 2016, DOI: 10.14293/S2199-1006.1.SOR-LIFE.AH9PUY.v1, p. 1-7.
(online at: www.scienceopen.com/hosted-document?doi=10.14293/S2199-1006.1.SOR-LIFE.AH9PUY.v1)
(Sangiran, Ngawi and Mojokerto sites in C and E Java. Vertebrate remains and hominid fossils mainly accumulated in continental sediments associated with lacustrine and fluvial systems. Pucangan Fm and lower part of Kabuh Fm in Mojokerto older than same formations from Sangiran and Ngawi B. In Mojokerto those formations were deposited during U Pliocene. Etc.)

Kealy, S., J. Louys & S. O'Connor (2015)- Islands under the sea: a review of early modern human dispersal routes and migration hypotheses through Wallacea. J. Island Coastal Archaeology 11, 3, p. 364-384.
(online at: https://www.researchgate.net/publication/287798057_Islands_Under_the_Sea_A_Review_of_Early_Modern_Human_Dispersal_Routes_and_Migration_Hypotheses_Through_Wallacea)
(Review of possible Late Pleistocene modern humans migration routes across Wallacea, the transitional island biogeographic zone between Sundaland (SE Asia) and Sahul (Australia-New Guinea))

Kealy, S., J. Louys & S. O'Connor (2017)- Reconstructing palaeogeography and inter-island visibility in the Wallacean Archipelago during the likely period of Sahul colonization, 65-45 000 years ago. Archaeological Prospection 24, 3, p. 259-272.
(online at: https://onlinelibrary.wiley.com/doi/full/10.1002/arp. 1570)

Kealy, S., J. Louys & S. O'Connor (2018)- Least-cost pathway models indicate northern human dispersal from Sunda to Sahul. J. Human Evolution 125, p. 59-70.
(online at: https://www.sciencedirect.com/science/article/pii/S0047248418302136)

(Discussion of possible human migration routes from Sundaland to Australia around 65-70 ka. Most likely route believed to be northern path across Sulawesi to Misool to coasts of New Guinea, not Java-Timor route)

Keates, S.G. (1998)- A discussion of the evidence for early hominids on Java and Flores. In: G.J. Bartstra (ed.) Bird's Head approaches; Irian Jaya studies; a programme for interdisciplinary research, Modern Quaternary Research in Southeast Asia, Balkema, Rotterdam, 15, p. 179-191.
(online at: https://www.papuaerfgoed.org/sites/default/files/collectie/files/2006-05/Bartstra_1998_Birds.pdf)
(Discussion of earliest proposed hominid occupation of island SE Asia, specifically of Flores Island)

Keates, S.G. (2004)- Notes on the Palaeolithic finds from the Walanae Valley, southwest Sulawesi, in the context of the Late Pleistocene of island Southeast Asia. In: S.G. Keates & J. Pasveer (eds.) Quaternary Research in Indonesia, Chapter 7, Modern Quaternary Research in Southeast Asia 18, Balkema, Leiden, p. 95-110.
(Stone artefacts from Walanae valley, SW Sulawesi, may represent earliest human (Homo sapiens) activity in early part of Late Pleistocene (first identified as Cabenge flake industry by Van Heekeren in late 1940's). With review of Late Pleistoce stone tool industries in other parts of Indonesia)

Keates, S.G. & G.J. Bartstra (1994)- Island migration of early modern *Homo sapiens* in Southeast Asia: the artifacts from the Walanae Depression, Sulawesi, Indonesia. Palaeohistoria 33/34, p. 19-30.
(online at: http://rjh.ub.rug.nl/Palaeohistoria/article/view/25055/22513)
(Lithic artifacts from Late Pleistocene river terraces in N Walanae depression, S Sulawesi)

Keates, S.G. & G.J. Bartstra (2001)- Observations on Cabengian and Pacitanian artefacts from island Southeast Asia. Quartar 51/52, p. 9-32.
(online at: http://quartaer.eu/pdfs/2001/2001_01_keates.pdf)
(Paleolithic stone artefact collections from Walanae valley near Cabenge in Sulawesi and from Baksoka valley near Pacitan, S Java, date largely to Late Pleistocene)

Kelley, J. (2002)- The hominoid radiation in Asia. In: W.C. Hartwig (ed.) The primate fossil record, Cambridge University Press, p. 369-384.

Kidder, J.H. & A.C. Durband (2004)- A re-evaluation of the metric diversity within *Homo erectus*. J. Human Evolution 46, p. 299-315.

Kirleis, W., J. Muller, C. Kortemeier, H. Behling & S. Soeghondo (2012)- The Megalithic landscape of Central Sulawesi, Indonesia: Combining archaeological and palynological investigations. In: Crossing Borders: Selected papers from 13th Int. Conf. European Association of Southeast Asian Archaeologists, NUS Press, p. 199-218.

Kleiweg de Zwaan, J. (1943)- De oudste mensheid van de Indische Archipel. Servire, Den Haag, p. 1-112.
(online at: https://resolver.kb.nl/resolve?urn=MMKB18B:051970000:pdf)
('The oldest humanity of the Indies Archipelago'. Popular review)

Kloosterman, F.H. (1989)- Groundwater flow systems in the northern coastal lowlands of West- and Central Java, Indonesia. Ph.D. Thesis, Free University, Amsterdam, p. 1-298.
(online at: dare.ubvu.vu.nl/bitstream/1871/12723/1/tekst.pdf)
(Mainly on groundwater systems on N Java coastal areas, with chapters on regional geologic setting, stratigraphy, Pleistocene tectonics and climate, etc. Emergence of Java as island in Pleistocene resulted not only from huge volumes of Pleistocene volcanic rocks but also major uplifts along axis of island)

Koesoemadinata, S. & M. Situmorang (1985)- Quaternary Geologic map of the Bekasi Quadrangle, Jawa, 1:50,000. Systematic Quaternary Geologic Map of Indonesia, Geological Research Development Centre (GRDC), Bandung, p.

Kotaka, T. & F. Hasibuan (1983)- Molluscan fossils from the Sangiran Dome, Central Jawa. Fossils (Palaeontological Association Japan) 33, p. 1-11. *(in Japanese with English abstract)*
(online at: http://ci.nii.ac.jp/naid/110002703460/en)
(Plio-Pleistocene mollusc assemblages from Sangiran Dome area, C Java: 25 species of marine bivalvia, 25 marine gastropoda, 3 non-marine bivalvia and 9 non-marine gastropoda identified from U Kalibeng Formation and Pucangan Fms. Paleoenvironmental conditions changed gradually from warm, shallow marine during U Kalibeng into tidal zone, then to fresh water facies during lower Pucangan Fm. Return of shallow marine environment during lower U Pucangan. Fresh water conditions returned to area during upper U Pucangan)

Koumans, F.P. (1949)- On some fossil fish remains from Java. Zoologische Mededelingen, Leiden, 30, 5, p. 77-82.
(online at: www.repository.naturalis.nl/document/150405)
(Description of rel. large Pleistocene fresh water fish remains from C Java, mainly Trinil, collected by Dubois)

Kramer, A. (1989)- The evolutionary and taxonomic affinities of the Sangiran mandibles of Central Java, Indonesia. Ph.D. Thesis, The University of Michigan, Ann Arbor, p. *(Unpublished)*

Kramer, A., T. Djubiantono, F. Aziz, J.S. Bogard, R.A. Weeks, D.C. Weinand, W.C. Hames et al. (2005)- The first hominid fossil recovered from West Java, Indonesia. J. Human Evolution 48, 6, p. 661-667. *(online at: https://www.academia.edu/102706667/The_first_hominid_fossil_recovered_from_West_Java_Indonesia)*
(M Pleistocene hominid teeth found during 1999 excavation in Cisanca River, Rancah, W Java, SE of Bandung. EPR dating and Ar-Ar dating of associated tuffs suggest age between 516-606 ka.)

Kretzoi, M. (1950)- *Stegeloxodon* nov. gen, a loxodonta elefantok esetleges azsiai ose. Foldtani Kozlony (Bull. Hungarian Geological Society) 80, p. 405-408.
(online at: https://www.biodiversitylibrary.org/item/205705#page/453/mode/1up)
('Stegeloxodon nov. gen, a possible Asiatic ancestor of true loxodonts'.New name Stegoloxodon indonesicus proposed for dwarfed elephantoid from Ci Pangglosoran near Bumiayu, W Java, based on unique loxodont-like M1 molar, as described in Van der Maarel (1932) who referred it to Elephas planifrons (Considered synonym of Elephas celebensis (Hooijer) by Maglio (1973); likely age E Pleistocene)

Kubat, J., A. Nava, L. Bondioli, C. Dean, C. Zanolli, N. Bourgon, A.M.Bacon, F. Demeter, B. Peripoli et al. (2021)- Dietary strategies of Pleistocene *Pongo* sp. and *Homo erectus* on Java (Indonesia). Nature Ecology Evolution 7, p. 279-289.
(manuscript online at: www.researchgate.net/publication/355017366_Dietary_strategies_of_Pleistocene_Pongo_sp_and_Homo_erectus_on_Java_Indonesia)
(Geochemical analyses of orangutan (Pongo sp.), Homo erectus and other Pleistocene mammalian teeth from Sangiran. Pongo sp. diet displays marked seasonal cycles. Omnivorous H. erectus shows less intra-annual Sr/Ca variability and was less dependent on variations in seasonal resource availability)

Kubo, D., R.T. Kono & Y. Kaifu (2013)- Brain size of *Homo floresiensis* and its evolutionary implications. Proc. Royal Society (London), B 280, 20130338, p.
(online at: http://royalsocietypublishing.org/content/royprsb/280/1760/20130338.full.pdf)
(Homo floresiensis from Late Pleistocene of Flores has extremely small endocranial volume (LB1 type specimen ~400cc). Hypotheses discussed: (1) H. floresiensis experienced dramatic brain size reduction from the Homo erectus (~1000 cc) in isolated insular setting; (2) species derived directly from more primitive and smaller-brained form such as Homo habilis (~600 cc) or Australopithecus (~400 cc))

Kumazawa, S. (1994)- Quaternary geology and hydrogeology of the Madiun Basin, Indonesia. J. Geosciences, Osaka City University, 37, 8, p. 213-242.
(online at: http://dlisv03.media.osaka-cu.ac.jp/infolib/user_contents/kiyo/DB00000180.pdf)
(Madiun basin in E Java with >250m of Quaternary sediments deposited above thick argillaceous sediments of Lower Pleistocene Pucangan stage. M-U Pleistocene Kabuh, Notopuro and Setri Fms consist of thick fluvial sediments and form good aquifers)

Kumai, H., M. Itihara, Sudijono, T. Shibasaki, F. Aziz, S. Yoshikawa et al. (1985)- Geology and stratigraphy of the Mojokerto area. In: N. Watanabe & D. Kadar (eds.) Quaternary geology of the hominid fossil bearing formations in Java. Geological Research Development Centre (GRDC), Bandung, Special Publ. 4, p. 55-61.
(Geological survey around the site of Perning near Mojokerto, site of P. mojokertensis discovery by Andoyo. Exact horizon of discovery uncertain)

Kupper, H. (1930)- Paleolitische werktuigen uit Atjeh, Nord Sumatra. Tijdschrift Koninklijk Nederlands Aardrijkskundig Genootschap (2) 47, p. 985-989.
(online at: https://resolver.kb.nl/resolve?urn=MMUBA13:001673001:pdf)
('Paleolithic tools from Aceh, North Sumatra'. Paleolithic stone tools near Lho Seumaweh at N and E coasts of Aceh found during geologic survey work. Mainly made from pebbles of dark quartzite)

Kupper, H. (1948)- Zur Kenntnis des prahistorischen Menschen auf West Java. Annalen Naturhistorischen Museums Wien 56, p. 434-439.
(online at: https://www.zobodat.at/pdf/ANNA_56_0434-0439.pdf)
('About the knowledge of the prehistoric people on West Java'. Report of artifacts found in several areas north of Bandung: obsidian tools, ceramic shards, etc. (at various times also collected and studied by R. von Koenigswald, W. Rothpletz, W. Mohler, etc.; JTvG))

Kurniawan, I., E. Setiyabudi, Y. Kaifu, F. Aziz & H. Baba (2013)- Evolution of Indonesian *Homo erectus* through the Middle Pleistocene: significance of Sambungmacan 4. In: F. Aziz & H. Baba (eds.) *Homo erectus* in Indonesian- Recent progress of the study and current understanding, Centre for Geological Survey, Bandung, Special Publ., Chapter 5, p. 93-102.
(Sambungmacan 4, discovered around 2002 in Solo River in C Java, is exceptionally well preserved Homo skull cap. Original stratigraphic position unclear. Some of its morphology transitional between Homo erectus (Sangiran) and Homo erectus (Ngandong; formerly Homo soloensis))

Kurten, B. (1964)- The sabre-toothed cat *Megantereon* from the Pleistocene of Java. Zoologische Mededelingen, Leiden, 38, 6, p. 101-104.
(online at: www.repository.naturalis.nl/document/149473)
(Description of tooth of sabre-toothed cat from unspecified Java location in Dubois collection, Leiden)

Kusumayuda, S.B., M.T. Zen, S. Notosiswoyo & R.S. Gautama (1999)- Distribution of the Gunung Sewu karstic aquifers based on fractal analysis- case study: Semanu and surrounding area, Yogyakarta, Indonesia. In: G.H. Teh (ed.) Proc. 9th Regional Congress Geology Mineral Energy Resources of SE Asia (GEOSEA '98), Kuala Lumpur 1998, Bull. Geological Society Malaysia 43, p. 345-350.
(online at: https://gsmpubl.files.wordpress.com/2014/09/bgsm1999034.pdf)

Kusumayuda, S.B., M.T. Zen, S. Notosiswoyo & R.S. Gautama (2000)- Fractal analysis of the Oyo River cave systems and topography of the Gunungsewu karst area, Central Java, Indonesia. Hydrogeological Journal 8, p. 271-278.
(Not much geology; wrong age model?)

Laitman, J.T. & I. Tattersall (2001)- *Homo erectus newyorkensis*: an Indonesian fossil rediscovered in Manhattan sheds light on the middle phase of human evolution. Anatomical Record 262, 4, p. 341-343.

Langbroek M. (2015)- Do tektites really date the bifaces from Bose (Baise) Basin, Guangxi, southern China? J. Human Evolution 80, p. 175-178. *(manuscript online at: www.academia.edu/10029842/Do_tektites_really_date_the_bifaces_from_the_Bose_Baise_basin_ Etc.)*
(Discussion of reported co-ocurrences by Chinese authors of Australasian tektes (~0.8 Ma old) and hominid biface artifacts in Bose Basin in S China (see also Michel et al. 2021))

Langbroek, M. & W. Roebroeks (2000)- Extraterrestrial evidence on the age of the hominids from Java. J. Human Evolution 38, p. 595-600.
(Review of uncertainties of numerical ages assigned to Java Homo erectus fossils. Presence of tektites in middle Bapang deposits can be tied to large Australasian strewnfield from asteroid impact near Laos or Cambodia around 700,000-800,000 years BP. This would make age of most Java hominids 1 Ma or younger)

Larena, M., J. McKenna, F. Sanchez-Quinto, C. Bernhardsson, C. Ebeo, R. Reyes, O. Casel et al. (2021)- Philippine Ayta possess the highest level of Denisovan ancestry in the world. Current Biology 31, 19, p. 4219-4230.
(Ayta Magbukon negrito tribe of SW Luzon, Philippines ~30-40% greater Denisovan ancestry than Australopapuans (highest in world). Together with Homo luzonensis, multiple archaic hominid species probably inhabited Philippines prior to arrival of modern humans, including Islander Denisovans)

Larick, R. & R.L. Ciochon (2014)- The Early Palaeolithic of Southeast Asia. In: C. Renfrew & P. Bahn (eds.) The Cambridge World History, Cambridge University Press, 1.21, p. 346-368.
("Digital access for individuals US$868.00- Kind regards from Cambridge University Press)

Larick, R. & R.L. Ciochon (2015)- Early hominin biogeography in island Southeast Asia. Evolutionary Anthropology 24, p. 185-213.
(online at: https://clas.uiowa.edu/sites/default/files/news-events/Larick-Ciochon-2015-Hominin%20Biogeography-ISEA-Evolutionary-Anthropology.pdf)
(Thorough review of hominid distribution in Indonesian region. Early hominin chronology difficult to resolve. 'Long chronology' suggests Homo erectus arrived in Sunda (Sangiran) before 1.5 Ma. An early form arrived E of Wallace Line (Flores) about 1.1 Ma. In Late Pleistocene, probable ultimate survivors of early dispersals found on Java (Ngandong Homo erectus, 125 ka), Flores (Homo floresiensis, 60 ka), and Luzon (Homo sp. indet, 67 ka). Early hominins responded to changing island conditions for 1.5 million years, likely becoming extinct when Homo sapiens colonized region. (first to suggest Homo erectus disappeared from Java for several 100,000 years after 0.8 Ma?; see also Li et al., 2021))

Larick, R., R.L. Ciochon, Y. Zaim, Sudijono, Suminto, Y. Rizal et al. (2001)- Early Pleistocene 40Ar/39Ar ages for Bapang Formation hominins, Central Jawa, Indonesia. Proc. National Academy of Sciences USA (PNAS) 98, 9, p. 4866-4871.
(online at: https://www.pnas.org/content/pnas/98/9/4866.full.pdf)
(Sangiran dome Plio-Pleistocene ~80 Homo erectus fossils. At 5 locations in Bapang (Kabuh) Fm H. erectus fossils associated with epiclastic pumice. 40Ar/39Ar ages from 1.51 Ma at Bapang/ Sangiran Fm contact, to 1.02 Ma above hominid-bearing sequence. Intermediate level with four crania has ~1.25 Ma age)

Larick, R., R.L. Ciochon & Y. Zaim (2004)- *Homo erectus* and the emergence of Sunda in the Tethys Realm. Contributions of potassium-based chronology in the Sangiran dome, Central Java. Athena Review 4, 1, p. 32-39.
(part online at: www.athenapub.com/13sunda.htm)

Larick, R., R.L. Ciochon, Y. Zaim, Sudijono, Suminto, Y. Rizal & F. Aziz (2000)- Lithostratigraphic context for Kln-1993.05-SNJ, a fossil colobine maxilla from Jokotingkir, Sangiran Dome. International J. of Primatology 21, 4, p. 731-759.
(New subspecies of colobine monkey described by Jablonski and Tyler (1999) from near Krikilan, Sangiran dome, C Java, unlikely to be from Late Pliocene Lower Lahar volcanic breccia. Not found in situ, and probably derived from Upper Sangiran (Pucangan) or lower Bapang (Kabuh) Fms)

Larson, S.G., W. Jungers, M.J. Morwood, T. Sutikna, Jatmiko, E.W. Saptomo et al. (2008)- *Homo floresiensis* and the evolution of the hominin shoulder. J. Human Evolution 53, 6, p. 718-731.
(online at: www.researchgate.net/publication/6146487_Homo_floresiensis_and_the_evolution_of_the Etc.)
(Homo floresiensis from Liang Bua probably did not have a modern human shoulder configuration)

Larson, S.G., W. Jungers, M. Tocheri, C. Orr, M.J. Morwood, T. Sutikna, R.A. Awe & T Djubiantono (2009)- Descriptions of the upper limb skeleton of *Homo floresiensis*. J. Human Evolution 57, 5, p. 555-570.

Leakey, R.E. & L.J. Slikkerveer (1993)- Man-ape ape-man: the quest for human's place in nature and Dubois' 'missing link'. Netherlands Foundation for Kenya Widlife Service, p. 1-179.
(Account of life and science of Eugene Dubois and debates on 'Java Man')

Lee, S.H. & D.G. Khorasani (2017)- Spread of hominins in Asia. eLS, p. 1-7.
(Review of migration of hominids in Asia. Earliest hominins in Asia almost as old as first appearance of genus Homo in Africa. Most fossil materials from Asia are without reliable dates)

Leinders, J.J.M., F. Aziz, P.Y. Sondaar & J. de Vos (1985)- The age of hominid-bearing deposits of Java: state of the art. Geologie en Mijnbouw 64, 2, p. 167-173.
(online at: https://drive.google.com/file/d/0B7j8bPm9Cse0ZXh0eVY0UGFkOWc/view)
(New standard biozonation for Plio-Pleistocene continental deposits of Java. Ages of our oldest faunal zones/ mammal assemblages: Satir fauna ~1.5 Ma, Ci Saat/ Satir fauna~1.2 Ma, Trinil HK ~1.0 Ma and Kedung Brubus ~0.8 Ma)

Li, H., M.G. Lotter, K. Kuman, L. Lei & W. Wang (2020)- Population dynamics during the Acheulean at ~0.8 Ma in East and Southeast Asia: considering the influence of two geological cataclysms. Palaeogeogr. Palaeoclim. Palaeoecology 562, 109927, p. 1-10.
(Two sudden penecontemporaneous geological events at ~0.8 Ma (Australasian tektite fall and slightly younger Oldest Toba super-eruption) likely caused severe environmental change (e.g., deforestation), which likely created population bottleneck effect for hominid stone tool makers, who likely occupied the landscape.Oldest Toba supereruption correlates with significant faunal turnover (pygmy stegodon replaced by large-bodied Stegodon, etc,) and Sangiran Homo erectus disappeared for several 100 kyrs after eruption. Study of Acheulean sites in SE and E Asia highlights potential relationship between Acheulean population dynamics and sudden environmental change)

Lilley, I. (1998)- East of Irian: Archaeology in Papua New Guinea. In: G.J. Bartstra (ed.) Bird's Head approaches; Irian Jaya studies; a programme for interdisciplinary research, Modern Quaternary Research in Southeast Asia 15, Balkema, Rotterdam, p. 135-156.
(online at: https://www.papuaerfgoed.org/sites/default/files/collectie/files/2006-05/Bartstra_1998_Birds.pdf)
(Review of archeological reserarch in PNG until 1990))

Lin, S.C., L.T. White, Jatmiko, I.M.A. Julianto, M.W. Tocheri & T. Sutikna (2022)- Characterising the stone artefact raw materials at Liang Bua, Indonesia. J. Paleolithic Archaeology 5, 22, p. 1-32.
(online at: https://link.springer.com/article/10.1007/s41982-022-00133-9)
(At Liang Bua cave (type site of Homo floresiensis) on Flores, stone artefact assemblages dominated by two raw materials:(1) locally available 'silicified tuff' (= actually iron-rich chert) and (2) chert. Both derived from Neogene fossiliferous tuff-bearing limestone with local diagenetic silica nodules. Shift towards increased chert artefacts at Liang Bua after ~46 ka, coincident with arrival of Homo sapiens at site)

Lipson, M., P.R. Loh, N. Patterson, P. Moorjani, Y.C. Ko, M. Stoneking, B. Berger & D. Reich (2014)- Reconstructing Austronesian population history in Island Southeast Asia. Nature Communications 5, 4689, p. 1-7.
(online at: www.nature.com/ncomms/2014/140819/ncomms5689/pdf/ncomms5689.pdf)
(Genetic studies of 31 Austronesian-speaking peoples in SE Asia suggest 'Austronesian expansion,' which began 4000-5000 years ago, likely had roots in Taiwan)

Locatelli, E., R. Awe Due, G.D. van den Bergh & L.W. van den Hoek Ostende (2012)- Pleistocene survivors and Holocene extinctions: the giant rats from Liang Bua (Flores, Indonesia). Quaternary International 281, p. 47-57.
(online at: http://ro.uow.edu.au/cgi/viewcontent.cgi?article=1054&context=smhpapers)

(Liang Bua Cave with common micromammal remains, mainly murids, incl. 3 species of giant rats. Papagomys armandvillei still living on Flores, other (Papagomys theodorverhoeveni, Spelaeomys florensis) went extinct during Holocene. Giant murids examples of insular gigantism. Peak in number of giant rats in cave in Holocene may result from Paleolithic hunting activity

Louys, J. (2007)- Limited effect of the Quaternary's largest super-eruption (Toba) on land mammals from Southeast Asia. Quaternary Science Reviews 26, p. 3108-3117.
(Relatively few species became extinct in SE Asia following the North Sumatra Toba super-eruption at 74 ka)

Louys, J. (2007)- Ecology and extinction of Southeast Asia's megafauna. Ph.D. Thesis University of New South Wales, Sydney, p. 1-290.
(online at: http://unsworks.unsw.edu.au/fapi/datastream/unsworks:1693/SOURCE02)
(On extinction of large Pleistocene mammal species in SE Asia)

Louys, J. (2008)- Quaternary extinctions in Southeast Asia. In: A.M.T. Elewa (ed.) Mass extinction, Springer-Verlag, Heidelberg, p. 159-189.

Louys, J. (2012)- Mammal community structure of Sundanese fossil assemblages from the Late Pleistocene, and a discussion on the ecological effects of the Toba eruption. Quaternary International 258, p. 80-87.
(Mammal community structure of Late Pleistocene W Indonesia fossil sites Punung, Sibrambang and Lida Ajer show community structure more similar to each other than to modern Sundanese communities. Niah Caves exhibits structure consistent with modern communities. Modern community structure in Sundaland may have formed between 128-118 ka and 45-39 ka, possibly result of Toba eruption, although more likely stem from environmental fluctuations accompanying glacial/interglacial cycles)

Louys, J. (2014)- The large terrestrial carnivore guild in Quaternary Southeast Asia. Quaternary Science Reviews 96, p. 86-97.
(SE Asia's Pleistocene large terrestrial carnivores appearances and disappearances from region poorly understood. Two significant extinctions represented by Pachycrocuta and Pliocrocuta. Disappearance of hyenids probably related to climate change. Several large carnivores from Java unique genetic and morphological variations, possibly related to connection between Java and Indochinese mainland in M Pleistocene. At least one endemic large carnivore evolved in Sundaland (Sunda clouded leopard))

Louys, J. (2016)- The giant rats of Timor. Australasian Science 37, 3, p. 24-26.
(Dog-sized giant rats coexisted with humans for 40,000 years on Timor)

Louys, J. (2024)- Quaternary palaeontological and archaeological research in Sumatra. In: J. Louys et al. (eds.) Quaternary Palaeontology and Archaeology of Sumatra, Terra Australis 56, ANU Press, Chapter 1, p. 1-12.
(online at: https://press-files.anu.edu.au/downloads/press/n12084/pdf/ch01.pdf))
(Late Quaternary history of Sumatra experienced relatively little attention. First reports of Pleistocene fossils date to late 1880s through the efforts of E. Dubois in caves of the Padang Highlands. Later finds made by A. Tobler, J. Zwierzycki, P. van Stein Callenfels, Van der Hoop, etc.)

Louys, J , P.C.H. Albers & A.E.A. van der Geer (eds.) (2024)- Quaternary palaeontology and archaeology of Sumatra. Terra Australis 56, ANU Press, p. 1-300.
(online at: https://press-files.anu.edu.au/downloads/press/n12084/pdf/book.pdf)
(Collection of 14 papers)

Louys, J., T.J. Braje, C.H. Chang, R. Cosgrove, S.M. Fitzpatrick M. Fujitah, S. Hawkins, T. Ingicco et al. (2021)- No evidence for widespread island extinctions after Pleistocene hominin arrival. Proc. National Academy of Sciences USA (PNAS) 118, 202023005118, p. 1-8.
(online at: https://www.pnas.org/doi/pdf/10.1073/pnas.2023005118)
(Hominins, including our H.sapiens, have not always had negative impacts on island ecosystems that humans have today. Rather, extinctions accelerated beginning in Holocene. Includes data from Indonesia)

Louys, J., D. Curnoe & H. Tong (2007)- Characteristics of Pleistocene megafauna extinctions in Southeast Asia. Palaeogeogr. Palaeoclim. Palaeoecology 243, p. 152-173.
(online at: http://www.rhinoresourcecenter.com/pdf_files/132/1328059508.pdf)
(Extinction of large-bodied taxa from Pleistocene in mainland SE Asia and Indonesia (incl. proboscideans Stegodon and Palaeloxodon, pygmy hippopotamus Hexaprotodon, hyenas Crocuta and Hyaena, giant panda Ailuropoda, tapirs Tapirus and Megatapirus and giant Asian ape Gigantopithecus, cannot be assigned to single cause. Disappearance likely tied to both climatic and human agents)

Louys, J., M. Duval, G.J. Price, K. Westaway, Y. Zaim, Y. Rizal, Aswan, M. Puspaningrum et al. (2022)- Speleological and environmental history of Lida Ajer cave, western Sumatra. Philosophical Transactions Royal Society (London), B 377, 20200494, p. 1-10.
(online at: https://royalsocietypublishing.org/doi/pdf/10.1098/rstb.2020.0494)
(Lida Ajer in Padang Highlands of W Sumatra believed to contain earliest evidence for presence of modern humans in rainforests in Indonesia. Two human teeth collected here by E. Dubois in 1880s estimated to be 73–63 kyrs old, older than some estimates of modern human migration out of Africa based on genetic data. Associated with abundant orangutan fossils.)

Louys, J. & S. Kealy (2024)- How did *Homo erectus* reach Java? Least-cost pathway models and a consideration of possible Sumatran routes. In: J. Louys et al. (eds.) Quaternary Palaeontology and Archaeology of Sumatra, Terra Australis 56, ANU Press, Canberra, Chapter 9, p. 199-216.
(online at: https://press-files.anu.edu.au/downloads/press/n12084/pdf/ch09.pdf)
(On possible routes that H. erectus could have taken on their journey into Indonesian region. Easiest pathway ran through centre of Sundaland, an area now almost entirely submerged. During periods of higher sea-levels, pathway moved west and could coincide with emergent areas on or just off E coast of Sumatra.Etc.)

Louys, J., S. Kealy, S. O'Connor, G.J. Price, S. Hawkins, K. Aplin, Y. Rizal, J. Zaim et al. (2017)- Differential preservation of vertebrates in Southeast Asian caves. Int. J. Speleology 46, 3, p. 379-408.
(online at: https://digitalcommons.usf.edu/ijs/vol46/iss3/6/)
(Results of three years of cave surveys in Indonesia and Timor-Leste, describing cave systems and their attendant vertebrate accumulations in diverse geological, biogeographical and environmental settings)

Louys, J. & E. Meijaard (2010)- Palaeoecology of Southeast Asian megafauna-bearing sites from the Pleistocene and a review of environmental changes in the region. J. of Biogeography 37, 8, p. 1432-1449.
(Reconstructions of habitat of 25 Pleistocene sites in SE Asia with medium- and large-bodied mammals. Sites classified as closed (continuous tree cover), mixed (heterogeneous tree cover) and open (limited or no tree cover; incl. Trinil)

Louys, J., S. O'Connor, Mahirta, P. Higgins, S. Hawkins & T.Maloney (2018)- New genus and species of giant rat from Alor Island, Indonesia. J. Asia-Pacific Biodiversity 11, 4, p. 503-510.
(online at: https://www.sciencedirect.com/science/article/pii/S2287884X18302152)
(Quaternary giant rat fossils previously reported from Timor, Sumba and Flores. First record of giant rat from Alor island, Makpan Cave: Alormys aplini gen. et sp. nov. Resembles Milimonggamys and Papagomys. Extinction in late Holocene appears coeval with disappearance of giant rats of Timor and Sumba and may be related to increased aridity and deforestation)

Louys, J., G.J. Price & S. O'Connor (2016)- Direct dating of Pleistocene *Stegodon* from Timor Island, East Nusa Tenggara. PeerJ. 2016, 4, e1788, p. 1-16.
(online at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4793331/pdf/peerj-04-1788.pdf)
(U-series dating of Stegodon tusk from Pleistocene Ainaro gravels near Atambua, Timor, indicate presence of stegodons in Timor at or before 130 ka, pre-dating earliest evidence of humans on island)

Louys, J. & P. Roberts (2020)- Environmental drivers of megafauna and hominin extinction in Southeast Asia. Nature 586, 7829, p. 402-406. *(manuscript online at: https://research-repository.griffith.edu.au/bitstream/handle/10072/402368/Louys445678 Etc.)*
(SE Asia home to at least five members of genus Homo. Considerable turnover in Pleistocene megafauna previously linked with hominins or with climate change. Forests of E Pleistocene had given way to savannahs by M Pleistocene, which led to spread of grazers and extinction of browsers. Savannahs retreated by Late Pleistocene and disappeared by Holocene epoch, and replaced by stratified closed-canopy rainforest. Ascendency of rainforest-adapted species as well as Homo sapiens at expense of savannah and woodland specialists, including Homo erectus)

Louys, J. & A. Turner (2012)- Environment, preferred habitats and potential refugia for Pleistocene *Homo* in Southeast Asia. Comptes Rendus Palevol 11, 2-3, p. 203-211.
(online at: https://www.sciencedirect.com/journal/comptes-rendus-palevol/vol/11/issue/2)
(SE Asia dominated by mix of savannah, open woodlands and evergreen forests in much of Pleistocene, conditions ideal for early hominin subsistence. These conditions rare for much of rest of Asia during glacial periods. SE Asia was possible refugium for savannah-adapted species like hominins during these periods)

Louys, J., Y. Zaim, Y. Rizal, G.J. Price, Aswan, M.R. Puspanigrum, H. Smith & A. Trihascaryo (2021)- Palaeontological surveys in Central Sumatra and Bangka. Berita Sedimentologi 47, p. 50-56.
(online at: https://journal.iagi.or.id/index.php/FOSI/article/view/358/327)
(Results of 2018 -2019 surveys by Australian National University and ITB on Quaternary deposits from caves in West Sumatra and attempt to relocate fossil proboscidean remains reported from Bangka Island)

Lovejoy, C.O. (1970)- The taxonomic status of the '*Meganthropus*' mandibular fragments from the Djetis beds of Java. Man, new series, 5, 2, p. 228-236.
(Material described as Meganthropus palaeojavanicus by Von Koenigswald in 1941 from Djetis beds of Java, lies within expected range of variation of Homo erectus)

Lydekker, R. (1885)- Description of a tooth of *Mastodon latidens*, Clift, from Borneo. Proc. Zoological Society of London 53, 4, p. 777-779.
(online at: https://www.biodiversitylibrary.org/page/31015890#page/905/mode/1up)
(Molar of Pleistocene(?) Mastodon from Brunei. Very similar to M. latidens from Siwalik fauna of India and from Burma, possibly also Java as described by Martin)

Lyras, G.A., M.D. Dermitzakis, A.A.E. van der Geer, S.B. van der Geer & J. de Vos (2009)- The origin of *Homo floresiensis* and its relation to evolutionary processes under isolation. Anthropological Science 117, 1, p. 33-43.
(online at: https://www.jstage.jst.go.jp/article/ase/117/1/117_080411/_pdf)
(Morphometric analysis of skulls separates H. floresiensis (LB1) from all H. sapiens, while not possible to separate H. floresiensis from H. erectus. Neolithic skulls from Flores within range of modern humans and not related to LB1)

Mahirta (2005)- Stone technology and the chronology of human occupation on Rote, Sawu and Timor, Nusa Tenggara Timur, Indonesia. Bull. Indo-Pacific Prehistory Association (IPPA) 29, p. 101-108.
(online at: https://journals.lib.washington.edu/index.php/BIPPA/article/view/9483/8470)
(Human occupation in Nusa Tenggara started from at least 30,000-35,000 years ago. Stone technology in region may have changed at start of Last Glacial Maximum (driest period), as indicated by disappearance of large tools with Hoabinhnian-like technologies. Data for earliest period (30-19 ka) mainly from caves Lua Meko on Rote and Lene Hara in E Timor)

Mahmud, M.I. (2018)- The archaeology of Sulawesi: an update, 2016. In: S. O'Connor et al. (eds.) The archaeology of Sulawesi: current research on the Pleistocene to the historic period, Terra Australis 48, ANU Press, Canberra, Chapter 1, p. 1-8.
(online at: https://press-files.anu.edu.au/downloads/press/n4569/pdf/ch01.pdf)

(Review of status of 2016 knowledge of archeology of Sulawesi)

Maloney, T.R., I.E. Dilkes-Hall, P. Setiawan, A.A. Oktaviana, I. Made Geria, M. Effendy, M. Ririmasse et al. (2022)- A late Pleistocene to Holocene archaeological record from East Kalimantan, Borneo. Quaternary Science Reviews 277, 107313, p. 1-21.
(Archaeological excavations at Liang Jon, in karst region of Sangkulirang-Mangkalihat Peninsula, NE Kalimantan, revealed cultural sequence in limestone rock shelter covering period ~16.7 kyr BP- late Holocene)

Mano, K., Sudijono & U.L. Batu (1991)- Stratigraphy along the middle course area of Solo River and its tributaries between Teluk and Ngawi. In: Quaternary geology of the northern foot area of Mount Lawu and along the middle course of the Solo River, central and East Java, Indonesia-Japan Joint Study Team, Geological Research Development Centre, Bandung, p. 14-36.

Maringer, J. & T. Verhoeven (1970)- Die Steinartefakte aus der *Stegodon*-Fossilschicht von Mengeruda auf Flores, Indonesien. Anthropos 65, 1-2, p. 229-247.
('The stone artifacts from the Stegodon fossil beds of Mengeruda on Flores, Indonesia'. Pleistocene volcanoclastic fossiliferous beds on Soa Plateau, W Central Flores (Matamenge, Olabula, etc.). Contains Stegodon trogonocephalus (Hooijer 1957), crocodiles, rodents, etc., and also Pleistocene tektites and rel. crude stone tools similar to 'Sangiran stone tool industry' of C Java and 'Cabenge industry' of Sulawesi (later dated at Mata Menge as ~880 ka; JTvG))

Maringer, J. & T. Verhoeven (1970)- Die Oberflachenfunde aus dem Fossilgebiet von Mengeruda und Olabula auf Flores, Indonesien. Anthropos 65, p. 530-565.
('The surface discoveries from the fossil areas of Mengeruda und Olabula on Flores'. 158 stone artifacts)

Maringer, J. & T. Verhoeven (1970)- Note on somestone artifacts in the National Archaeological Institute of Indonesia at Djakarta, collected from the *Stegodon*-fossil bed at Boaleza in Flores. Anthropos 65, 3-4, p. 638-639.
(Brief report on 12 of many stone artifacts collected by Verhoeven in 1963-1968 from Stegodon beds in Flores)

Maringer, J. & T. Verhoeven (1972)- Steingerate aus dem Waiklau-Trockenbett bei Maumere auf Flores, Indonesien. Eine Patjitanian-artige Industrie auf der Insel Flores. Anthropos 67, 1-2, p. 129-137.
('Stone tools from the Waiklau dry river bed near Maumere, Flores, Indonesia'. 30 artifacts, collected by Verhoeven in dry bed of Waiklau river in 1965 and 1968. Similaries with Pacitanian of Java)

Maringer, J. & T. Verhoeven (1975)- Die Oberflachenfunde von Marokoak auf Flores, Indonesien. Ein weiterer altpalaolithischer Fundkomplex von Flores, Indonesien. Anthropos 70, 1-2, p. 97-104.
('The surface discoveries of Marakoak on Flores- a new Early Paleolithic occurrence on Flores')
(Description of stone tools from Marokoak locality 16km NW of other localities Olabula and Mengeruda)

Maringer, J. & T. Verhoeven (1977)- Ein palaolithischer Hohlenfundplatz auf der Insel Flores, Indonesien. Anthropos 72, 1-2, p. 256-273.
('A Paleolithic cave deposit on the island of Flores'.Stone tools from Liang Michael cave deposits near Dalong village, West Flores (Manggarei), including choppers, adzes, flakes, etc. Associated with Varanus hooijeri skull (described by Brongersma, 1958), bones of giant rats and mollusc fragments. M-U Paleolithic; similarities with 'Sangiran industry'.)

Maringer, J. & T. Verhoeven (1979)- Recent discovery of a Palaeolithic past in Flores, Indonesia, and its contribution to the research of most ancient Southeast Asia. East and West 29, 1-4, p. 247-263.
(Review of work by Dutch/ German Catholic missionaries and amateur paleontologist/archeologists Theodor T. Verhoeven and Johannes Maringer since 1956 on Flores on Pleistocene mammalian faunas (incl. small Stegodon, Varanus and giant rats) and man-made stone artifacts. These discoveries 'destroyed the Wallace Line hypothesis')

Maringer, J. & J. Verschuuren (1981)- Zum Palaolithikum der Insel Timor, Indonesien. Anthropos 76, 3-4, p. 584-588.
('On the Paleolithic of Timor Island'. Stone tools (mainly multi-colored chert, some obsidian) from Talau River area S of Atambua, E part of W Timor, collected by missionary J. Verschuuren in 1964-1965)

Markov, G.N. & H. Saegusa (2008)- On the validity of *Stegoloxodon* Kretzoi, 1950 (Mammalia: Proboscidea). Zootaxa 1861, 1, p. 55-56.
(Discussion of taxonomic validity of Stegoloxodon, based on dwarf elephantoid molar from Bumiaya, Java)

Marks, P. (1953)- Preliminary note on the discovery of a new jaw of *Meganthropus* Von Koenigswald in the lower Middle Pleistocene of Sangiran, Central Java. Indonesian J. Natural Science (Majalah Ilmu Alam untuk Indonesia) 109, p. 26-33.
(On large fragment of hominid mandible collected by villagers N of Glagahombo village in conglomerate layer probably equivalentto 'Grenzbank'. Attributed to Meganthropus palaeojavanicus)

Marquez, S., K. Mowbray, G.J. Sawyer, T. Jacob & A. Silvers (2001)- New fossil hominid calvaria from Indonesia- Sambungmacan 3. The Anatomical Record 262, p. 344-368.
(online at: http://onlinelibrary.wiley.com/doi/10.1002/ar.1046/pdf)
(On skull cap of M Pleistocene Homo cf. erectus from banks of Solo River near Poloyo village, Sambungmacan district, C Java. Absence of some classic characters attributed to Homo erectus can be interpreted as: (1) known cranial variation of H. erectus from Indonesia and China; (2) calvaria shows evidence of evolutionary change within H. erectus; or (3) more than one species of Homo in M Pleistocene of Java (see also Broadfield et al., 2001, Delson et al. 2001) (Sm3 discovered in Java in 1977 and illegally moved to New York)

Martin, K. (1884)- Ueberreste vorweltlicher Proboscidier von Java und Banka. Sammlungen Geologischen Reichs-Museums Leiden, Ser. 1, 4, p. 1-24.
(online at: www.repository.naturalis.nl/document/552394)
('Remnants of prehistoric elephants from Java and Bangka'. Early paper on molars of Pleistocene elephantoids: Stegodon sp. from Pati Ajam (S of Muriah volcano, C Java) and Elephas sumatranus from tin sands on Bangka)

Martin, K. (1884)- Ueberreste vorweltlicher Proboscidier von Java und Bangka. Jaarboek Mijnwezen Nederlandsch-Indie 13, Wetenschappelijk Gedeelte, Batavia, p. 285-308. *(without plates online at: https://ia600704.us.archive.org/0/items/jaarboekvanhetm04gebigoog/jaarboekvanhetm04gebigoog.pdf)*
('Remnants of prehistoric elephants from Java and Bangka'. Reprint of Martin (1884))

Martin, K. (1886)- Fossile Saugethierreste von Java und Japan. Sammlungen Geologischen Reichs-Museums Leiden, Ser. 1, 4, p. 25-69.
(online at: www.repository.naturalis.nl/document/552399)
('Fossil mammal remains from Java and Japan'. Early description of Pleistocene mammal remains collected by Raden Saleh in 1865-1866, mainly from C Java Solo area. Mainly Stegodon (skull and molars), Elephas, whale bones and deer (also in Jaarboek Mijnwezen Nederlandsch Oost-Indie 16, p. 1-45))

Martin, K. (1887)- Fossile Saugethierreste von Java und Japan. Jaarboek Mijnwezen Nederlandsch Oost-Indie 16, Wetenschappelijk Gedeelte, Palaeontologie Nederlandsch-Indie, Verhandelingen 21, p. 1-45.
('Fossil mammal remains from Java and Japan'. Same as Martin (1886))

Martin, K. (1888)- Neue Wirbelthierreste vom Pati-Ajam auf Java. Sammlungen Geologischen Reichs-Museums Leiden, Ser. 1, 4, p. 87-116.
(online at: www.repository.naturalis.nl/document/552407)
('New vertebrate remains from Pati-Ajam on Java'. Early paper on Pleistocene mammals from Pati-Ajam mountains S of Muriah volcano, near Klaling, Jepara Residency, C Java. Mainly molars of Mastodon, Stegodon, Euelephas, Bos sp. and Cervus sp., Material collected by Koorders, but locality first described by Junghuhn 1857. Similar mammal fauna known from Solo region)

Martin, K. (1888)- Neue Wirbeltierreste vom Pati-Ajam auf Java. Jaarboek Mijnwezen Nederlandsch Oost-Indie 17 (1888), Wetenschappelijk Gedeelte, p. 20-48.
('New vertebrate remains from Pati-Ajam on Java'. Reprint of Martin (1888)

Martin, R.D., A.M. MacLarnon, J.L. Phillips & W.B. Dobyns (2006)- Flores hominid: new species or microcephalic dwarf ? The Anatomical Record, A, 288A, 11, p. 1123-1145.
(online at: http://onlinelibrary.wiley.com/doi/10.1002/ar.a.20389/pdf)
(New hominids 'Homo floresiensis' from Flore, dated at ~18,000 yrs, very small but dentally adult. Commonly interpreted as insular dwarf derived from Homo erectus, but far too small to derive from Homo erectus by normal dwarfing. H. floresiensis most likely microcephalic Homo sapiens with advanced stone tools)

Marwick, B., C. Clarkson, S. O'Connor & S. Collins (2016)- Early modern human lithic technology from Jerimalai, East Timor. J. Human Evolution 101, p. 45-64.
(Jerimalai rock shelter in E Timor with large assemblage of Pleistocene stone artefacts and shell fish hooks dated to 42,000 yrs BP, one of oldest known sites of modern human activity in island SE Asia. Little change in lithic technology over 42,000 year sequence until addition of new types and raw materials in M Holocene. Assemblage dominated by small chert cores and implements rather than pebble tools and choppers (common in island SE Asia as opposed to mainland SE Asia). Jerimalai assemblage resembles Liang Bua assemblage of Flores, associated with Homo floresiensis; both possibly created by modern humans)

Matsumura, H. (2000)- Origins of Southeast Asian people as viewed from cranial and dental morphology. Asian Paleoprimatology, Kyoto, 1, p. 149-160.
(online at: https://core.ac.uk/download/pdf/39322992.pdf)
(Human skeletons of Hoabinian (late Pleistocene- E Holocene) from Malaysia, Indonesia and Vietnam demonstrate Australo-Melanesian- affinity cranial and dental morphology. These were members of population that appeared in late Pleistocene Sundaland, and ancestors of modern Australian Aborigines. Subsequent Neolithic to modern populations in SE Asia have dental and cranial characteristics more similar to modern N and E Asians, suggesting migrants from Asia mainland (China?) expanded into its peninsula and island regions of SE Asia since Neolithic, supporting the "dual ancestry hypothesis" for SE Asia population history)

Matsumura, H., M.F. Oxenham, Nguyen K.T., Nguyen L.C. & Nguyen K.D. (2011)- Population history of mainland Southeast Asia: the Two Layer model in the context of Northern Vietnam. In: N.J. Enfield (ed.) Dynamics of human diversity, Pacific Linguistics, Canberra, p. 153-178.

Matsumura, H., K. Shinoda, T. Simanjuntak, A.A. Oktaviana, S. Noerwidi, H.O. Sofian et al. (2018)- Cranio-morphometric and aDNA corroboration of the Austronesian dispersal model in ancient Island Southeast Asia: support from Gua Harimau, Indonesia. PLoS One 13, 6, e0198689, p. 1-25.
(online at: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0198689)
(Gua Harimau cave in Padang Bindu, S Sumatra, with 84 human skeletons dating from pre-Neolithic to Neolithic, Bronze and Iron Ages (~5600- 1800 BP). Cranial and DNA evidence for at least 2-3 distinct populations from two separate time periods)

Matsu'ura, S. (1982)- A chronological framing for the Sangiran hominids. Bull. National Science Museum, Tokyo, D8, p. 1-53.
(online at: http://ci.nii.ac.jp/els/...)
(Fluorine dating of ~250 E Pleistocene vertebrate fossils from Grenzbank and Kabuh Fms of Sangiran area, C Java, suggesting age range of ~0.8-1.1 Ma for Pithecanthropus Beds)

Matsu'ura, S. (1985)- A consideration of the stratigraphic horizons of hominid finds from Sangiran by the fluorine method. In: N. Watanabe & D. Kadar (eds.) Quaternary geology of the hominid fossil bearing formations in Java. Geological Research Development Centre (GRDC), Bandung, Special Publ. 4, p. 359-366.

Matsu'ura, S. (1986)- Fluorine and phosphate analysis of fossil bones from the Kabuh formation of Trinil. Bull. National Science Museum, Tokyo, D12, p. 1-9.

(online at: http://ci.nii.ac.jp/naid/110000008540/en)
(Bone samples from lower Kabuh Fm in Trinil area analysed for fluorine and phosphate for comparison with Sangiran other areas. No obvious conclusions?)

Matsu'ura, S. (2022)- Toward the resolution of controversial chronologies for the World Heritage Site of Sangiran: Dating the first appearance of *Homo erectus* in the island of Java. The Quaternary Research (Daiyonki-Kenkyu, Japan) 61, 1, p. 1-25. *(In Japanese)*
(online at: https://www.jstage.jst.go.jp/article/jaqua/61/1/61_2108/_pdf/-char/en)
(Review of radiometric and other age dating of Pleistocene stratigraphy at Sangiran, C Java. First appearance datum for Sangiran hominin fossils best considered as 1.3 Ma (<1.5 Ma), which is significantly later than dates widely accepted over last two decades (looks thorough; JTvG))

Matsu'ura, S., M. Kondo, F. Aziz, Sudijono, S. Narasaki & N. Watanabe (2000)- First known tibia of an early Javanese hominid. Current Anthropology 41, 2, p. 297-300.
(On modern-shaped hominid tibia (shin bone) recovered in 1977 from Sambungmacan, C Java)

Matsu'ura, S., M. Kondo, T. Danhara, S. Sakata, H. Iwano, T. Hirata, I. Kurniawan, E. Setiyabudi, Y. Takeshita et al. (2020)- Age control of the first appearance datum for Javanese *Homo erectus* in the Sangiran area. Science 367, 6474, p. 210-214.
(online at: https://www.science.org/doi/pdf/10.1126/science.aau8556)
(Combination of zircon, etc. dating techniques for hominin-bearing sediments at Sangiran, C Java, suggests arrival of Homo erectus at ~1.3 Ma, at least 200,000 years younger than previously thought by many. Age of zircons in Lower Bapang Fm tuff just above Grenzbank very close to 0.9 Ma. Tuff below base Upper Sangiran Fm <1.55 Ma (below oldest Sangiran hominins))

Matthew, W.D. (1928)- The ape-man of Java. Does this fossil prove that a creature which was in a halfway stage between Man and Ape lived about a million years ago? Natural History 28, p. 576-588.

McAllister, M.S., M.W. Morley, J.J. Tyler, F.A. McInerne & A.J. Blyth (2022)- Investigating the palaeoenvironmental context of Late Pleistocene human dispersals into Southeast Asia: a review of stable isotope applications. Archaeological Anthropological Sciences 14, 86, p. 1-27.
(online at: https://link.springer.com/content/pdf/10.1007/s12520-022-01540-3.pdf)
(Review of stable isotope work in Late Pleistocene archeological sites in SE Asia, mainly on tooth enamel and speleothems (still in its infancy))

McKenzie, K.G. & Sudijono (1981)- Plio-Pleistocene ostracoda from Sangiran, Jawa. Publ. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 1, p. 29-51.
(Descriptions of 25 ostracode species in Late Pliocene Upper Kalibeng Fm and Pleistocene Pucangan Fm at Sangiran Dome, C Java. Shallow marine Kalibeng Fm fauna with Neonesidea, Cytherelloidea, Thalmannia, Hemicytheridea, etc. Lower Pucangan Fm with brackish water assemblage of Ishizakiella, Hemicytheridea, Ilyocypris, Stenocypris and Cyprinotus. Lacustrine Upper Pucangan Fm with Physocypria and Indocythere)

Medway, Lord (1972)- The Quaternary mammals of Malesia: a review. In: P. & M. Ashton (eds.) The Quaternary era in Malesia, Transactions of the 2nd Aberdeen-Hull Symposium on Malesian Ecology, University of Hull Dept. of Geography Miscellaneous Series 13, p. 63-98.

Meijaard, E. (2003)- Mammals of South-East Asian islands and their Late Pleistocene environments. J. of Biogeography 30, 8, p. 1245-1257.
(During Last Glacial Maximum several areas in Sunda region remained forest-covered: W Sumatra, NW Borneo, Malacca Straits and around Palawan. Other areas possibly more open vegetation types, like tree savanna or open deciduous forest: Malay/Thai Peninsula, Java Sea (incl. Sunda Strait) and E Borneo)

Meijaard, E. (2004)- Solving mammalian riddles: a reconstruction of the Tertiary and Quaternary distribution of mammals and their palaeoenvironments in island South-East Asia. Ph.D. Thesis Australian National University, Canberra, p. 1-347.
(online at: https://digitalcollections.anu.edu.au/handle/1885/47989)
(New Miocene-Quaternary biogeographic models for SE Asia that help explain present-day distribution patterns and evolutionary relationships between mammal species)

Meijaard, E. & C.P. Groves (2006)- The geography of mammals and rivers in mainland Southeast Asia. In: M. Lehman & J.G. Fleagle (eds.) Primate biogeography- progress and prospects, Springer, New York, p. 305-330.
(Late Pliocene- E Pleistocene environmental changes in mainland SE Asia split up many tropical species leading to diversification, maintained during Pleistocene by further glacial periods. During last glacial maximum this may have led to isolation of rainforest-dependent species in several refugia. M Pleistocene catastrophic comet collision around 0.77 Ma, with center of impact in E Thailand or E Cambodia/S Laos, may have caused widespread extinction in mainland SE Asia in area possibly >1 million km^2)

Meijer, H.J.M. (2014)- The avian fossil record in Insular Southeast Asia and its implications for avian biogeography and palaeoecology. PeerJ 2, e295, p. 1-13.
(online at: https://peerj.com/articles/295.pdf)
(Review of bird fossils from Indonesia/ SE Asia islands. At least 63 species in 54 genera and 27 families recorded. Except for Eocene of Sumatra, all bird fossils are Pleistocene in age)

Meijer, H.J.M. & R.A. Due (2010)- A new species of giant marabou stork (Aves: Ciconiiformes) from the Pleistocene of Liang Bua, Flores (Indonesia). Zoological J. Linnean Society, London, 160, 4, p. 707-724.
(online at: https://academic.oup.com/zoolinnean/article/160/4/707/2625584)
(Fossils of new species of giant marabou stork, Leptoptilos robustus, from Pleistocene of Liang Bua, Flores. Giant bird estimated at 1.80 m in length. Large body size and terrestrial lifestyle responses to unbalanced, island environment with abundant prey and lack of mammalian carnivores)

Meijer, H.J.M., F. d'Errico, A. Queffelec, I. Kurniawan, E. Setiabudi, I. Sutisna, A. Brumm & G.D. van den Bergh (2019)- Characterization of bone surface modifications on an Early to Middle Pleistocene bird assemblage from Mata Menge (Flores, Indonesia) using multifocus and confocal microscopy. Palaeogeogr. Palaeoclim. Palaeoecology 529, p. 1-11.
(online at: https://www.sciencedirect.com/science/article/pii/S0031018219300264)

Meijer, H.J.M., I. Kurniawan, E. Setiabudi, A. Brumm, T. Sutikna, R. Setiawan & G.D. van den Bergh (2015)- Avian remains from the Early/Middle Pleistocene of the So'a Basin, central Flores, Indonesia, and their palaeoenvironmental significance. Palaeogeogr. Palaeoclim. Palaeoecology 440, p. 161-171.
(Soa Basin in C Flores with 16 late Early to early M Pleistocene terrestrial fossil localities with insular endemic faunas. Remains of 6 bird species birds from Mata Menge and Bo'a Leza include a swan (Cygnus sp.) and eagle owl (Bubo sp.), indicative of open environment with open freshwater and nearby grasslands)

Meijer, H.J.M., J. Louys & S. O'Connor (2019)- First record of avian extinctions from the Late Pleistocene and Holocene of Timor Leste. Quaternary Science Reviews 203, p. 170–184
(Late Pleistocene and Holocene bird remains from Jerimalai and Matja Kuru sites on Timor Leste with archaeological sequences dating back to >40 ka. All bird species still present on Timor today, except a crane (Grus sp.) from Late Pleistocene of Jerimalai, and large buttonquail (Turnix sp.) from Matja Kuru)

Meijer, H.J.M., T. Sutikna, E.W. Saptomo, R. Due Awe, Jatmiko, S. Wasisto, H.F. James, M.J. Morwood & M.W. Tocheri (2013)- Late Pleistocene-Holocene non-passerine avifauna of Liang Bua (Flores, Indonesia). J. Vertebrate Paleontology 33, 4, p. 877-894.
(online at: https://repository.si.edu/bitstream/handle/10088/28869/Meijer-13-JVP-Ling%20Bua%20non-passerines.pdf?sequence=1&isAllowed=y;View/Open)
Liang Bua cave deposits on Flores span last 95,000 years, with bird fossils throughout. Late Pleistocene assemblage with 23 taxa. Giant marabou Leptoptilos robustus and vulture Trigonoceps sp. now extinct)

Meijer, H.J.M., T. Sutikna, E.W. Saptomo & M.W. Tocheri (2022)- More bones of *Leptoptilos robustus* from Flores reveal new insights into giant marabou stork paleobiology and biogeography. Royal Society (London) Open Science. 9, 220435, p. 1-32.
(online at: https://royalsocietypublishing.org/doi/epdf/10.1098/rsos.220435)
(Bones of giant marabou storks (Leptoptilos robustus) from Liang Bua site, Flores. Well developed wing bones of L. robustus suggest it was capable of active flight. L. robustus bones broadly similar to Leptoptilos falconeri from sites in Africa and Eurasia (incl. Leptoptilos titan from Ngandong, C Java; Wetmore, 1940). Pleistocene dispersal of L. falconeri into Island SE Asia may have given rise to populations of giant marabou storks in this region. Island SE Asia likely acted as refugium for last surviving members of this lineage)

Meijer, H.J.M., M.W. Tocheri, R. Awe Due, T. Sutikna, E.W. Saptomo & H.F. James (2015)- Continental–style avian extinctions on an oceanic island. Palaeogeogr. Palaeoclim. Palaeoecology 429, p. 163-170.
(Comparison of 25 species of modern non-passerine avifauna of Flores with Late Pleistocene assemblage from cave site of Liang Bua cave. Rel. few extinctions on Flores, except large-bodied avian scavengers Leptoptilos robustus (giant stork) and Trigonoceps (vulture). Possibly linked to extinction of pygmy elephant (Stegodon florensis insularis))

Meijer, H.J.M., L.W. van den Hoek Ostende, G.D. van den Bergh & J. de Vos (2010)- The fellowship of the hobbit: the fauna surrounding *Homo floresiensis*. J. of Biogeography 37, 6, p. 995-1006.
(online at: www.academia.edu/1353347/Five_years_of_Homo_floresiensis)
Flores vertebrate fauna low species richness and disharmonic fauna, resulting from isolated position of island. H. floresiensis associated with common pygmy proboscidean Stegodon florensis insularis, giant rats (Papagomys armandvillei, P. theodorverhoeveni, Spelaeomys florensis) and other murids, bats, Komodo dragon (Varanus komodoensis, V. hooijeri), and large number of birds (incl. giant marabou Leptoptilos). Between fossil-bearing localities Ola Bula Fm (~900-800 ka) and Liang Bua (~95-0 ka) gap of ~700 kyr)

Menzies, J.I. & C. Ballard (1994)- Some new records of Pleistocene megafauna from New Guinea. Science in New Guinea 20, p. 113-139.
(Four new sites with Plio-Pleistocene megafauna in PNG, incl. zygomaturine diprodontid Hulitherium tomasetti (marsupial equivalent of giant panda) and Protemnodon)

Tablizo, M.U., G.D. van den Bergh & A.G.S. Fernando (2025)- Island-hopping across the Wallace Line: A new Pleistocene *Stegodon* fossil skull from Luzon (Philippines) reveals dispersal links to Wallacea. Palaeogeography, Palaeoclimatology, Palaeoecology 677, 113186, p.
(E Pleistocene Stegodon skull from Lannig, Solana, Cagayan (NE Luzon), with affinities to S. trigonocephalus and also S. f. florensis from Flores, Indonesia, suggesting N-S faunal connection between the Philippines and Wallacea)

Mijares, A.S., F. Detroit, P. Piper, R. Grun, P. Bellwood, M. Aubert, G. Champion, N. Cuevas, A. De Leon & E. Dizon (2010)- New evidence for a 67,000-year-old human presence at Callao Cave, Luzon, Philippines. J. Human Evolution 59, 1, p. 123-132.
(Human third metatarsal from Callao Cave in N Luzon dated by U-series ablation as ~67 ka, making it oldest known modern human fossil in Philippines (and SE Asia?). Morphometric analysis indicates gracile structure, close to small-bodied Homo sapiens, but also within ranges of Homo habilis and H. floresiensis)

Mijsberg, W.A. (1927)- Over het in 1926 te Trinil gevonden en ten onrechte als rest van het schedeldak van een praehistorisch mensch beschouwde fossiel. Geneeskundig Tijdschrift Nederlandsch-Indie, 1927, p. 159-165.
('On the fossil found in Trinil in 1926 and erroneously considered to be a skull cap of a prehistoric human'. In August 1926 Dr. C.E.J. Heberlein from Surabaya bought a fossil at Trinil that he identified as a Pithecanthropus skull cap. In December 1926 anatomist Mijsberg at S.T.O.V.I.A in Batavia and H.J.T. Bijlmer identified it as proximal end of the humerus of a fossil elephant, which is not uncommon inTrinil Pleistocene deposits)

Mijsberg, W.A. & H.J.T. Bijlmer (1927)- Report on a fossil found at Trinil in 1926, and erroneously considered as the remains of a skull of *Pithecanthropus*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 30, 1, p. 126-133.
(online at: https://https://dwc.knaw.nl/DL/publications/PU00015425.pdf)
(English version of paper above)

Mijsberg, W.A. (1932)- Recherches sur les restes humains trouves dans les fouilles de l'abris-sous-roche de Goewa-Lawa a Sampoeng et des sites prehistoriques a Bodjonegoro (Java). In: Hommage du Service Archeologie des Indes Neerlandaises au Premier Congres des Prehistoriens d'Extreme-Orient a Hanoi, Batavia, p. 39-54.
('Investigations on the human remains found in the excavations of the Gua Lawa rock shelter in Sampung and prehistoric sites in Bojonegoro (Java)'. Report on skeletal remains of 'big-teeth' prehistoric people from East Java, reminescent of Papua-Melanesian racial group)

Mishra, S., C. Gaillard, C. Hertler, A.M. Moigne & T. Simanjuntak (2010)- India and Java: contrasting records, intimate connections. Quaternary International 223-224, p. 265-270.
(Comparison of similarities of archeological, paleontological and hominin records of India and Java)

Mohler, W.A. (1946)- Zur Stratigraphie der Saugetierfuhrenden Schichten von Java. Experientia- Monatsschrift fur das gesamte Gebiet der Naturwissenschaft, Basel, 2, 8, p. 287-292.
('On the stratigraphy of mammal-bearing beds of Java'. Mammalian faunas recognized, from old to young: (1) Tji Djolang (M Pliocene), (2) Kali Glagah (U Pliocene), (3) Djetis with Pithecanthropus (Lower Pleistocene), (4) Trinil with Pithecanthropus (M Pleistocene), (5) Ngandong with Homo neanderthalensis soloensis (U Pleistocene), (6) Sampoeng (Sampung; Subrecent, postdating M Pleistocene folding (U Pleistocene Ngandong-terraces not folded))

Moigne, A.M., R. Due Awe, F. Semah & A.M. Semah (2004)- The cervids from Ngebung site ('Kabuh series', Sangiran dome, Central Java) and their biostratigraphical significance. In: S.G. Keates & J. Pasveer (eds.) Quaternary Research in Indonesia, Chapter 3, Modern Quaternary Research in Southeast Asia 18, Balkema, p. 31-44.
(Excavations in Lower Kabuh Fm at Ngebung site with fossil riverbank horizon formerly occupied by Homo erectus, with numerous stone artifiacts and broken bones, dated at ~0.8 Ma (Ar-dating of overlying tuff by Salei 1997). Dominant animal fossils large bovids, smaller cervids and Stegodon, an association probably linked to human activity. Two cervids, Cervus (Rusa) and Axis lydekkeri ngebungensis)

Moigne, A.M., F. Semah & A.M. Semah, A. Bouteaux & R. Due Awe (2004)- Mammalian fossils from two sites of the Sangiran Dome (Central Jawa, Indonesia), in the biostratigraphical framework of the Jawanese Pleistocene. In: L.C. Maul & R.D. Kahlke (eds.) Late Neogene and Quaternary biodiversity and evolution: Regional developments and interregional correlations, Proc. 18th International Senckenberg Conference, Weimar 2004, Terra Nostra, Stuttgart, p. 176-178. *(Extended Abstract)*
(online at: www.senckenberg.de/fis/doc/abstracts/68_Moigne_etal_2.pdf)
(Brief review of Bukuran and Ngebung 2 sites, Sangiran. Ngebung 2 with 'late Trinil HK' mammalian assemblage and dated as beginning of M Pleistocene (~0.9 Ma?))

Moncel, M.H., M. Arzarello, E. Boeda, T. Bonilauri, B. Chevrier, C. Gaillard, H. Forestier, Y. Li, F. Semah & V. Zeitoun (2018)- Assemblages with bifacial tools in Eurasia (second part). What is going on in the East? Data from India, Eastern Asia and Southeast Asia. Comptes Rendus Palevol 17, 1-2, p. 61-76.
(online at: www.sciencedirect.com/science/article/pii/S1631068315002122)
(Review of Pleistocene stone tools in Asia, incl. Indonesia: Baturaja-S Sumatra, Pacitan-Java, Ngebung/ Sangiran-Java (part 1 of series was on stone tools in Europe))

Moncel, M.H., M. Arzarello, E. Boeda, T. Bonilauri, B. Chevrier, C. Gaillard, H. Forestier, Y. Li, F. Semah & V. Zeitoun (2018)- Assemblages with bifacial tools in Eurasia (third part). Considerations on the bifacial phenomenon throughout Eurasia. Comptes Rendus Palevol 17, 1-2, p. 77-97.

(online at: www.sciencedirect.com/science/article/pii/S163106831630032X)
(Bifacial stone tool technology believed to become widespread from 800-700 ka onwards, probably reaching Levant from Africa before moving toward Asia, then Europe. However, reality may be more complex. In Indonesia lithic pieces compatible with Acheulean traditions found without stratigraphic context in S Sumatra and associated with Homo erectus fossils at base of Kabuh Fm in Sangiran)

Moore, M.W. & A. Brumm (2007)- Stone artifacts and hominins in island Southeast Asia: new insights from Flores, eastern Indonesia. J. Human Evolution 52, p. 85-102.
(online at: www.academia.edu/466208/Stone_artifacts_and_hominins_in_island_Southeast_Asia_new_ Etc.)
(Review of stone tool types in Indonesia. Large-sized 'core tools' commonly believed to be work of Homo erectus while assemblages of small-sized 'flake tools' usually attributed to Homo sapiens, but both are probably part of same sequences)

Moore, M.W., T. Sutikna, Jatmiko, M.J. Morwood & A. Brumm (2009)- Continuities in stone flaking technology at Liang Bua, Flores, Indonesia. J. Human Evolution 57, 5, p. 503-526.
(online at: www.academia.edu/1259545/Biface_distributions_and_the_Movius_Line_A_Southeast_Asia Etc.)
(At Liang Bua, Flores, stratified unchanging artifact sequence spanning 95 kyr, with minor shift to unifacial flaking after 11 ka. Pleistocene pattern associated with Homo floresiensis skeletal remains. Holocene changes correlate with appearance of Homo sapiens)

Morley, M.W. (2017)- The geoarchaeology of hominin dispersals to and from tropical Southeast Asia: a review and prognosis. J. Archaeological Science 77, p. 78-93.
(Review of geoarchaeology of Late Pleistocene modern human dispersals into and out of SE Asia, incl. Indonesian localities Punung/ Wajak (Java) and Liang Bua (Flores))

Morwood, M.J., F. Aziz, G.D. van den Bergh, P.Y. Sondaar & J. De Vos (1997)- Stone artefacts from the 1994 excavation at Mata Menge, West Central Flores, Indonesia. Australian Archaeology 44, p. 26-34.
(online at: https://www.library.uq.edu.au/ojs/index.php/aa/article/download/996/994)
(1994 excavation in fluvial Ola Bula Fm at Mata Menge near Bajawa, C Flores, yielded M Pleistocene stone tool pieces (basalt and chert flakes) and faunal remains (large Stegodon, crocodile, giant rat). Likely Matuyama- Brunhes boundary (and tektites from same site reported by Maringer and Verhoeven 1970 (but below main fossil layers?))

Morwood, M.J., P. Brown, Jatmiko, T. Sutikna, E. Wahyu Saptomo, K.E. Westaway, R. Awe Due et al. (2005)- Further evidence for small-bodied hominins from the late Pleistocene of Flores, Indonesia. Nature 437, 7061, p. 1012-1017.
(Homo floresiensis from Late Pleistocene of Flores has stature, limb proportions and endocranial volume of African Pliocene Australopithecus. Age of population from 95-74 to 12 thousand years ago. Excavation yielded more evidence for behavioural capabilities, including butchery of Stegodon and use of fire)

Morley, R.J., H.P. Morley, Y. Zaim & O.F. Huffman (2020)- Palaeoenvironmental setting of Mojokerto *Homo erectus*, the palynological expressions of Pleistocene marine deltas, open grasslands and volcanic mountains in East Java. J. of Biogeography 47, 3, p. 566-583.
(online at: www.researchgate.net/publication/338691956_Palaeoenvironmental_setting_of_Mojokerto Etc.)
(Facies in Perning and Jetis sections near Mojokerto indicate landscape with four potential Homo habitats: muddy deltas with widespread Nypa swamps; poorly vegetated sandy delta; open savanna grasslands in lowlands up river of delta; and volcanoes in upper reaches of catchment with perhumid montane podocarp and broad-leaf forests and probable open fire-climax Casuarina junghuhniana forest. Palynological data support Early Pleistocene geological age for Mojokerto H. erectus at about 1.43 Ma)

Morwood, M.J. & W.L. Jungers (eds.) (2009)- Paleoanthropological research at Liang Bua, Flores, Indonesia. J. Human Evolution 57, 5, p. 437-648.
(Collection of papers of Flores studies)

Morwood, M.J. & W.L. Jungers (2009)- Conclusions: implications of the Liang Bua excavations for hominin evolution and biogeography. J. Human Evolution 57, 5, p. 640-648.
(Liang Bua excavations on Flores stratified sequence of stone artifacts and faunal remains spanning ~95- 17 ka, and includes skeletal remains of Homo sapiens in Holocene and Homo floresiensis in Pleistocene. Small H. floresiensis not australopithecine and not dwarfing of ancestral H. erectus population, but probably late representative of small-bodied hominid lineage that exited Africa before emergence of Homo erectus)

Morwood, M.J., P.B. O'Sullivan, F. Aziz & A. Raza (1998)- Fission-track ages of stone tools and fossils on the East Indonesian island of Flores. Nature 392, p. 173-176.
(Zircon fission-track dates from two fossil sites on Flores. Tangi Talo, with endemic fauna, dates to 0.90 ± 0.07 Ma, Mata Menge (with stone tools and elements of continental SE Asian fauna) 0.88- 0.80 Ma. Also older, reworked zircons with ages between 7 and 14 Ma (also with Australasian tektites; Von Koenigswald 1957))

Morwood, M.J., P. O'Sullivan, E.E. Susanto & F. Aziz (2003)- Revised age for Mojokerto 1, an early *Homo erectus* cranium from East Java, Indonesia. Australian Archaeology 57, p. 1-4.
(online at: www.library.uq.edu.au/ojs/index.php/aa/article/viewFile/526/1690)
(Field study and redating of two pumice horizons at site of Homo erectus skull found by Duyfjes (1936) at Mojokerto, E Java, indicates age is <1.49 Ma, not much-hyped 1.8 Ma age of Swisher et al. (1984))

Morwood, M.J., R.P. Soejono, R.G. Roberts, T. Sutikna, C.S.M. Turney, K.E. Westaway et al. (2004)- Archaeology and age of a new hominin from Flores in eastern Indonesia. Nature 431, p. 1087-1091.
(Excavations at Liang Bua cave on Flores yielded tiny hominins, assigned to Homo floresiensis n.sp. It existed from before 38 ka until at least 18 ka. Associated deposits contain stone artefacts and animal remains, including Komodo dragon and endemic, dwarfed species of Stegodon. Homo floresiensis originated from early dispersal of Homo erectus and overlapped in time with Homo sapiens in region)

Morwood, M.J., T. Sutikna, E.W. Saptomo, K.E. Westaway, Jatmiko et al. (2008)- Climate, people and faunal succession on Java, Indonesia: evidence from Song Gupuh. J. Archaeological Science 35, 7, p. 1776-1789.
(Song Gupuh cave in Gunung Sewu Limestones, E Java, over 16m of deposits with faunal sequence spanning 70 ka. Terminal Pleistocene- Early Holocene period of maximum biodiversity. Human activity, especially after onset of Neolithic around 2.6 ka, contributed to progressive loss of species from area)

Morwood, M.J. & P. van Oosterzee (2007)- A new human- The startling discovery and strange story of the "hobbits" of Flores, Indonesia. Harper Collins, p. 1-222.
(Popular book on 2004 discovery of Homo floresiensis and its interpretation as a new species. N.B.: There is a school of paleoanthropologists that still dispute this 'startling discovery; see e.g. Jacob et al. (2006), Martin et al. (2006), Bednarik (2008), Obendorf et al. (2008), Oxnard et al. 2010, Henneberg et al. (2011), Eckhardt et al. (2014) and others, who argue Morwood et al. misrepresented facts and misinterpreted significance of H. floresiensis, which they believe represents aberrant specimens of Homo sapiens)

Movius, H.L. (1948)- The Lower Palaeolithic cultures of southern and eastern Asia. Transactions American Philosphical Society, new series, 38, 4, p. 329-420.
(online at: https://catalog.hathitrust.org/Record/001879951)
(Review of Pleistocene stratigraphy and hominids and stone tools in Java, NW India, Burma and N China)

Movius, H.L. (1949)- Lower Palaeolithic archaeology in southern Asia and the Far East. In: W.W. Howell (ed.) Early man in the Far East, Symposium, American Assoc. Physical Anthropologists, Chicago 1946, p. 17-77.
(Review of Pleistocene stratigraphy and hominids and stone tools in NW India, Burma, N China and Java)

Mubroto, B., Suminto & J. Kimura (1995)- Paleomagnetic analysis of sediments of the Kedungbrubus area. In: Sudijono et al. (eds.) Geology of Quaternary environment of the Solo- Madiun area, Central-East Java, Geological Research Development Centre (GRDC), Bandung, Special Publ. 17, p. 100-104.

Mulyaningsih, S., Sampurno, Y. Zaim, D.J. Puradimaja, S. Bronto & D.A. Siregar (2006)- Perkembangan geologi pada Kuarter Awal sampai masa sejarah di dataran Yogyakarta. Jurnal Geologi Indonesia 1, 2, p. 103-113.
(online at: www.bgl.esdm.go.id/publication/index.php/dir/article_detail/170)
('Developments in Quaternary geology until the beginning of history in the Yogyakarta plain'. Uplift of Southern Mts in E Pleistocene formed Yogyakarta Basin. Merapi volcanic activity took place since ±42 ka or 0.67 Ma)

Mulvaney, D.J. (1970)- The Patjitanian industry: some observations. Mankind (Australian J. Anthropology) 7, 3, p. 184-187.
(Pacitanian stone tools from Java S Mountains contain both hand-axes and flake tools and may not be as old as previously suggested)

Mulvaney, D.J. & R.P. Soejono (1970)- The Australian-Indonesian archaeological expedition to Sulawesi. Asian Perspectives 13, p. 163-177.
(online at: https://typeset.io/pdf/the-australian-indonesian-archaeological-expedition-to-59am14atzv.pdf)
(July-August 1969 joint Indonesian-Australian archeological expedition to Neolithic cave sites in Makassar region of S Sulawesi, to investigate remnants of Toalean culture. Earlier excavation work by Van Stein Callenfels in 1937 was deemed to be very sloppy: many interesting archeological objects remained in the vSC-Noone rubbish piles, and little or none of the 1937 work and findings had been properly documented)

Musser, G.G. (1981)- The giant rat of Flores and its relatives east of Borneo and Bali. Bull. American Museum Natural History 169, 2, p. 67-176.
(online at: http://digitallibrary.amnh.org/handle/2246/568)
(Five Pleistocene- Recent murids (rats) known only from Flores: Papagomys armandvillei and new species P. theodorverhoeveni, Hooijeromys nusatenggara, Floresomys naso and Spelaeomys florensis. Komodomys rintjanus occurs on Flores and Komodo Islands of Rintja and Padar)

Musser, G.G. (1982)- The Trinil rats. Modern Quaternary Research in Southeast Asia 7, Balkema, Rotterdam, p. 65-85.
(Review of extant and fossil murid rodents of Java. Rare murid rodents in Pleistocene Trinil fauna represent two species, incl. small Rattus trinilensis n.sp.)

Musser, G.G. & C. Newcomb (1983)- Malaysian murids and the giant rat of Sumatra. Bull. American Museum Natural History 174, 4, p. 327-598.
(online at: https://digitallibrary.amnh.org/handle/2246/572)
(Three Indo-Malayan murid (rodent) genera: Palawanomys, n. gen., known only from Palawan Island; Berylmys, primarily in Indochina and Sundamys n. gen., containing mountain giant rat of Sumatra, Sundamys is indigenous to W Indonesia (called 'Sunda Shelf' here, but it means onshore around Sunda Shelf; JTvG))

Musser, G.G., A. van de Weerd & E. Strasser (1986)- *Paulamys*, a replacement name for *Floresomys* Musser, 1981 (Muridae), and new material of that taxon from Flores, Indonesia. American Museum (Natural History) Novitates 2850, p. 1-10.
(online at: https://digitallibrary.amnh.org/handle/2246/5247)
(On renaming of subfossil endemic rat fossil material collected by T. Verhoeven from caves in Flores)

Naumann, E. (1887)- Fossile Elephantenreste von Mindanao, Sumatra und Malakka. Abhandlungen und Berichte Koniglichen Zoologischen Anthropologisch-Ethnographischen Museums zu Dresden 1, 6, p. 1-11.
(online at: https://www.biodiversitylibrary.org/item/104465#page/217/mode/1up)
('Fossil elephant remains from Mindanao, Sumatra and Malacca'. Stegodon trigonocephalus and S. aff. Insignis from NW Mindanao, Elephas indicus presumably from Malakka, possible Stegodon from Deli, N Sumatra)

Naumann, E. (1890)- *Stegodon mindanensis*, eine neue Art von Uebergangs-Mastodonten. Zeitschrift Deutschen Geologischen Gesellschaft, Berlin, 42, 1, p. 166-169.

(online at: https://www.biodiversitylibrary.org/item/37733#page/176/mode/1up)
('Stegodon mindanensis, a new species of transitional mastodonts'. Brief note on elephantoid tooth from Mindanao, Philippines, in collections of Dresden Museum, Germany. Initially identified as similar to Stegodon trigonocephalus from Java, but it is different. No figures)

Newman, K., B. Hakim, A.A. Oktaviana, B. Burhan, D. McGahan & A. Brumm (2022)- The missing deposits of South Sulawesi: new sources of evidence for the Pleistocene/ Holocene archaeological transition. Archaeological Research in Asia 32, 100408, p.
(Discussion of apparent gap in archeological record of S Sulawesi between ~20,000-10,000 years ago)

Ninkovich, D. & L.H. Burckle (1978)- Absolute age of the base of the hominid-bearing beds in Eastern Java. Nature 275, p. 306-308.
(Analysis of planktonic marine diatoms from marine intercalations in lowermost hominid-bearing beds and from underlying U Kalibeng Fm marine sediments in E Java (~2.1-1.9 Ma age for U Kalibeng assemblage))

Ninkovich, D., L.H. Burckle & N.D. Opdyke (1982)- Palaeogeographic and geologic setting for early Man in Java. In: R.A. Scrutton & M. Talwani (eds.) The ocean floor, Wiley, New York, p. 211-227.

Nishioka, Y. & C. Vidthayanon (2018)- First occurrence of *Duboisia* (Bovidae, Artiodactyla, Mammalia) from Thailand. Fossil Record (Berlin) 21, 2, p. 291-299.
(online at: https://fr.pensoft.net/article/30607/)
(First record from Thailand of ?M Pleistocene antelope-like bovid Duboisia skull with horns from sandpit at Tha Chang, E of Nakhon Ratchasima (Khorat), NE Thailand. Confirms genus not endemic to Java. Assigned to Duboisia aff. D. santeng. From just below tektite-bearing horizon)

Noerwidi, S. (2012)- The significance of the Holocene human skeleton Song Keplek 5 in the history of human colonization of Java: a comprehensive morphological and morphometric study. Masters Erasmus Mundus Thesis, Museum National Histoire Naturelle, Paris, p. 1-13.
(online at: https://www.academia.edu/34744974/The_significance_of_the_Holocene_human_skeleton_Song_Keplek_5_in_the_history_of_human_colonization_of_Java_a_comprehensive_morphological_and_morphometric_study)
(Detailed study of ~3000 yr old skeleton from Song Keplek rock shelter burial site in Gunung Sewu. S Java. Interpreted as Neolithic pioneer of SE Asian ("Southern Mongoloid") affinity, with strong morphological connections with recent Javanese and mainland SE Asian populations, who probably intermixed with indigenous Australo-Melanesian people. Probably multiple Holocene migrations of modern humans into Java)

Noerwidi, S. (2012)- Younger Toba Tephra 74 kya: impact on regional climate, terrestial ecosystem, and Prehistoric human population. AMERTA 30, 1, p. 9-18.
(online at: https://jurnalarkeologi.kemdikbud.go.id/index.php/amerta/article/view/396)
(Review of impact of large Toba eruption in 74 ka, N Sumatra, on vegetation, animals and human environmental change. Some scholars pointed to global consequence of killing most humans)

Noerwidi, S. (2020)- Hominin diversity in the western Indonesian archipelago during the Quaternary: a dental record perspective. Doct. Thesis Museum National Histoire Naturelle, Paris, p. 1-481.
(online at: https://tel.archives-ouvertes.fr/tel-03680512/document)
(Six groups of Pleistocene-Recent hominid teeth types)

Noerwidi, S., Siswanto & H. Widianto (2016)- Primata besar di Jawa: spesimen baru *Gigantopithecus* dari Semedo. Berkala Arkeologi 36, 2, p. 141-160.
(online at: http://berkalaarkeologi.kemdikbud.go.id/index.php/berkalaarkeologi/article/view/96/142)
('Giant primate of Java: a new Gigantopithecus specimen from Semedo'. Two enigmatic mandible specimens found in 2014 at Semedo, SE of Tegal, C Java. Morphologically similar to common primate jaw, but twice size. Semedo specimens close to Gigantopithecus blacki (first occurrence in Java); a species known from China and Himalyas))

Noerwidi, S., A. Vialet, H. Widianto, I. Kurniawan, J. Zaim, R.A. Suriyanto, J. Joordens et al. (2020)- Exploring the diversity of fossil hominin dental patterns in the western Indonesian archipelago during the Quaternary by Geometric Morphometric Analysis. Application on second upper and lower molars. L'Anthropologie 124, 5, 102791, p.
(manuscript online at: https://www.sciencedirect.com/science/article/pii/S0003552120300741)
(At least 4 groups (dental types) of hominins in Indonesia in Pleistocene times. Result suggest overlap among them throughout E and early M Pleistocene. Another group persists until Late Pleistocene, and two other groups of Homo sapiens present during Holocene)

Noerwidi, S., H. Widianto, M. Martinon-Torres, J.M. Bermudez de Castro, A.M. Moigne, A. Vialet & F. Semah (2019)- External dental characters and diversity of *Homo erectus* among the fossil record of Java. UISPP (J. Int Union for Prehistoric and Protohistoric Sciences) 2, Special Issue 2, p. 13-20.
(online at: https://cir.cenieh.es/bitstream/20.500.12136/1556/1/External%20dental%20characters%20 Etc)
(Dental morphological features of Homo erectus from Java (Sangiran, Wajak, etc.) suggest four groups of fossil hominins that supposedly coexisted in Java in lower Middle Pleistocene (see also Noerwidi, 2020 and others))

Noerwidi, S., H. Widianto, A.M. Moigne, A. Vialet, M.M. Ansyori, M.R. Fauzi, Hua Tu, C. Falgueres, A.M. Semah et al. (2023)- Preliminary study of two deciduous human molars from the Late Pleistocene layers of Song Terus (East Java): A window into the last *Homo erectus* and the first *Homo sapiens* in Java. L'Anthropologie 127, 3 (Special Issue Prehistoire de l'Indonesie), 103159, p. 1-22.
(Two previously undescribed human molars from Pleistocene in Song Terus cave near Punung in Southern Mountains. ST06 is dated >80 ka (belongs to Homo erectus s.l. group) and tooth ST04 <60 ka (close to Homo sapiens))

Noerwidi, S., H. Widianto, I.A. Nurani, L. Yondri, T. Setiawan, K. Wiradnyana, T. Simanjuntak et al. (2023)- Variability of the Late Pleistocene-Holocene *Homo sapiens* groups in Western Indonesian archipelago: focus on second molars. L'Anthropologie 127, 3, 103169, p.
(Late Pleistocene-Holocene human populations in W Indonesia two different dental types: (1) "Preneolithic" in E-M Holocene fossils (12- 4.5 ka), and (2) "Neolithic-Paleometallic" in Late Holocene (from 3.2 ka))

Obendorf, P.J., C.E. Oxnard & B.J. Kefford (2008)- Are the small human-like fossils found on Flores human endemic cretins? Proc. Royal Society (London), B, 275, p. 1287-1296.
(online at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2602669/pdf/rspb20071488.pdf)
(Hominid fossils from Liang Bua, Flores, Indonesia, including nearly complete skeleton dated to 18 ka and assigned to new species are probably "myxoedematous endemic (ME)" cretins, part of an inland population of Homo sapiens)

Obradovich, J.D. & C.W. Naeser (1981)- Indonesia- Age of *Homo erectus* from Java. In: Geological Survey Research 1981, U.S. Geological Survey (USGS) Professional Paper 1275, p. 285-286.
(Sample of dacite pumice collected 12m above level of Homo erectus VIII in Kabuh beds dated with K-Ar and fission track methods as 1.05 +/- 0.1 Ma and 1.6 +/- 0.7 Ma respectively)

O'Connell, C.A. & J.M. DeSilva (2013)- Mojokerto revisited: evidence for an intermediate pattern of brain growth in *Homo erectus*. J. Human Evolution 65, p. 156-161.
(online at: https://www.bu.edu/anthrop/files/2013/08/OConnellDeSilvaJHE2013.pdf)

O'Connell, J.F. & J. Allen (2004)- Dating the colonization of Sahul (Pleistocene Australia- New Guinea): a review of recent research. J. Archaeological Science 31, p. 835-853.
(Date for arrival of human colonization of Sahul area (Australia- New Guinea) generally assumed to be at ~40,000 BP or 60,000 BP. Postulated arrival dates before~42-45 ka not well-supported by data)

O'Connell, J.F., J. Allen, M.A.J. Williams, A.N. Williams, C.S.M. Turney, N.A. Spooner, J. Kamminga et al. (2018)- When did *Homo sapiens* first reach Southeast Asia and Sahul? Proc. National Academy of Sciences USA (PNAS) 115, 34, p. 8482-8490.
(online at: www.pnas.org/content/pnas/115/34/8482.full.pdf)
(Anatomically modern humans (Homo sapiens) began spreading across Eurasia from Africa and adjacent SW Asia at ~50-55 ka. Some argued that human genetic, fossil, and archeological data indicate one or more prior dispersals, possibly as early as 120 ka, but remains to be proven)

O'Connor, S. (2007)- Pleistocene Timor: further corrections, a reply to Bednarik. Australian Archaeology 54, p. 46-51.

O'Connor, S. (2007)- New evidence from East Timor contributes to our understanding of earliest modern human colonisation east of the Sunda Shelf. Antiquity 81, p. 523-535.
(New age date of ~38,000-42,000 yrs BP from rock shelter of Jerimalai, E Timor is earliest evidence for migration by modern humans E of Sunda Shelf into Island SE Asia)

O'Connor, S. (2010)- Pleistocene migration and colonization in the Indo-Pacific Region. In: A. Anderson et al. (eds.), The global origins and development of seafaring, McDonald Institute Archaeological Research, Cambridge, p. 41-55.
(Maritime spread was most likely mode of expansion for early modern humans moving out of Africa and along Indian Ocean Rim and reaching Sahul (Pleistocene Australia-New Guinea) at least 60,000 years ago)

O'Connor, S. & K. Aplin (2007)- A matter of balance: an overview of Pleistocene occupation history and the impact of the Last Glacial Phase in East Timor and the Aru Islands, eastern Indonesia. Archaeology in Oceania 42, 3, p. 82-90.
(Late Pleistocene Last Glacial Maximum (~30- 15 ka) increased aridity and lowered sea level had significant impact on human hunter-gatherers in E Timor and Aru Islands)

O'Connor, S., D. Bulbeck & J. Meyer (eds.) (2018)- The archaeology of Sulawesi- Current research on the Pleistocene to the Historic period. Terra Australis 48, Australian National University Press, Canberra, p. 1-357.
(online at: https://press-files.anu.edu.au/downloads/press/n4569/pdf/book.pdf))
(Review of archeology work in Sulawesi: in 30 papers presented at 2016 symposium in Makassar, covering the last ~100,000 years)

O'Connor, S., S. Kealy, A. Black, M. Ririmasse, Mahirta, S. Hawkins, M. Husni, D. Tanudirjo, Alifah et al. (2020)- The rock art of Kisar Island, Indonesia: a small island with a wealth and diversity of artistic expression. J. Indo-Pacific Archaeology 44, p. 19-51.
(online at: https://journals.lib.washington.edu/index.php/JIPA/article/view/15645)
(Kisar island N of Timor with 40 known painted rock art sites, mainly in shelters and caves in uplifted Quaternary reefal limestone terraces along S and E coasts. Some of art likely dates back to Pleistocene period. Many of the figurative motifs similar to those featured in rock art assemblages of Timor-Leste and other sites in E Indonesia)

O'Connor, S., S. Kealy, C. Reepmeyer, S.C. Samper Carro & C. Shipton (2023)- Terminal Pleistocene emergence of maritime interaction networks across Wallacea. World Archaeology 54, 1, p. 1-20.
(online at: https://www.researchgate.net/publication/368405625_Terminal_Pleistocene_emergence_of_maritime_interaction_networks_across_Wallacea)

O'Connor, S., J. Louys, S. Kealy & S.C. Samper Carro (2017)- Hominin dispersal and settlement East of Huxley's Line; the role of sea level changes, island size, and subsistence behavior. Current Anthropology 58, Supplement 17, p. S567-S582.
(online at: https://www.journals.uchicago.edu/doi/pdfplus/10.1086/694252)
(Pleistocene pre-sapiens hominins opportunistic omnivores, probably constrained to environments with plentiful fresh water animals and plants; therefore rel. difficult to migrate across island archipelago. Homo

sapiens probably able to subsist on marine resources and more easily moved through islands East of Huxley Line)

O'Connor, S., C. Shipton & S. Kealy (2024)- The southern route to Sahul: modern human dispersal and adaptation in the Pleistocene. In: R. Ono & A. Pawlik (eds.) The prehistory of human migration- human expansion, resource use, and mortuary practice in Maritime Asia, Intech Open, Chapter 1, p. 1-29.
(online at: https://www.researchgate.net/publication/376445965_The_Southern_Route_to_Sahul_Modern_Human_Dispersal_and_Adaptation_in_the_Pleistocene)
(Evidence for modern human dispersal and early settlement to southern islands of Wallacean Archipelago (Flores, Timor Leste, etc.). Modern humans (Homo sapiens) thought to have arrived in Sahul (Australia) at least 60,000 years ago (60 ka),but no firm evidence for settlement of Wallacea prior to ~50- 40 ka. Remarkable onset of obsidian lithics exchange network across islands of Alor, Atauro, Timor, and Kisar in terminal Pleistocene from ~17 ka, coinciding with new forms of personal ornamentation and rock art)

O'Connor, S., M. Spriggs & P. Veth (2002)- Excavation at Lene Hara Cave establishes occupation in East Timor at least 30,000-35,000 years ago. Antiquity 76, p. 45-50.
(First discovery of Late Pleistocene flake-based stone tools from Timor, in Lene Hara cave, Timor Leste (one of rel. many Late Pleistocene occurrences of small 'flake tool industries' in caves and rock shelters across E Indonesia region)

Oktaviana, A.A., R. Joannes-Boyau, B. Hakim, B. Burhan, R. Sardi, S. Adhityatama, Hamrullah, I. Sumantri, M. Tang et al. (2024)- Narrative cave art in Indonesia by 51,200 years ago. Nature 631, p. 814-818.
(online at: https://www.nature.com/articles/s41586-024-07541-7)
(New laser-ablation U-series imaging dating of earliest prehistoric rock art in limestone caves of Maros-Pangkep, SW Sulawesi, resulted in minimum ages of ~50-54 Ma)

Oktaviana, A.A., R. Joannes-Boyau, B. Hakim, B. Burhan, R. Sardi, S. Adhityatama, A. Jalandoni, Hamrullah, I. Sumantri et al. (2026)- Rock art from at least 67,800 years ago in Sulawesi. Nature , 16p. *(in press)*
(online at: https://www.nature.com/articles/s41586-025-09968-y)
(Hand stencil from Liang Metanduno on Muna Island, W of Buton, SE Sulawesi, yielded U-series age of 71.6 ± 3.8 ka, providing minimum-age constraint of 67.8 ka for underlying motif. Older than known rock art in Maros-Pangkep (S Sulawesi) and Spain (attributed to Neanderthals))

Olsen, J.W. & R.L. Ciochon (1990)- A review of evidence for postulated Middle Pleistocene occupations in Viet Nam. J. Human Evolution 19, 8, p. 761-788.
(Several archeological localities across Vietnam originally interpreted as of M Pleistocene age, but age control of many localities unreliable)

Ono, R., R. Fuentes, A.F. Pawlik, H.O. Sofian, Sriwigati, N. Aziz, N. Alamsyah & M. Yoneda (2020)- Island migration and foraging behaviour by anatomical modern humans during the late Pleistocene to Holocene in Wallacea: New evidence from Central Sulawesi, Indonesia. Quaternary International 554, 7521, p. 90-106.
(On maritime migration and island adaptation in SE Asia by anatomically modern humans, with new archaeological research E coast of Sulawesi)

Ono, R., A. Pawlik & R. Fuentes (2024)- The Pleistocene maritime migration of modern humans in northern Wallacea: The cases of Topogaro in Sulawesi and Bubog in Mindoro. In: R. Ono & A. Pawlik (eds.) The prehistory of human migration- human expansion, resource use, and mortuary practice in Maritime Asia, Intech Open, Chapter 2, p. 1-24.
(online at: https://www.academia.edu/122541589/The_Pleistocene_Maritime_Migration_of_Modern_Humans_in_Northern_Wallacea_The_Cases_of_Topogaro_in_Sulawesi_and_Bubog_in_Mindoro)
(Two important Pleistocene sites in N Wallacea important for role in maritime migration and seafaring of modern humans in Wallacea and Sahul: Goa Topogaro cave complex in C Sulawesi (42 ka), and Bubog rock-shelter sites in Mindoro, Philippines (at least 35 ka))

Ono, R., H.O. Sofian, R. Fuentes, N. Aziz, M. Ririmasse & I.M. Geria (2023)- Early modern human migration into Sulawesi and island adaptation in Wallacea. World Archaeology 54, 2, p. 229-243.
(New archaeological research in Goa Topogaro 2, C Sulawesi, a likely entry location for northern route of early modern humans' maritime migration from Asia into Wallacean islands and New Guinea during Late Pleistocene)

Oppenheimer, S. (2009)- The great arc of dispersal of modern humans: Africa to Australia. Quaternary International 202, p. 2-13.
(Late Pleistocene dispersal of anatomically modern humans out of Africa. Routes obeyed limitations placed by drinking water and climate-permissive corridors. First spread N in Eemian interglacial (~125 ka). Reached Indonesian region by 75-81 ka. Crossed Wallace Line to reach Australia at least by 48 ka (possibly 60 ka)

Oppenoorth, W.F.F. (1932)- De vondst van Palaeolithische menschelijke schedels op Java. De Mijningenieur 13, 6, p. 106-110.
('The discovery of Paleolithic human skulls on Java'. On new hominid skull discoveries in lower part of 20m river terrace, left bank of Solo River at Ngandong, C. Java)

Oppenoorth, W.F.F. (1932)- *Homo (Javanthropus) soloensis*, een Plistoceene mensch van Java. Voorloopige mededeeling. Wetenschappelijke Mededeelingen Dienst Mijnbouw Nederlandsch-Indie 20, p. 49-75.
('Homo (Javanthropus) soloensis, a new Pleistocene hominid from Solo River terrace at Ngandong, E Java'. First description of Ngandong hominid fossils (skull caps, etc.). Associated with rich mammal assemblage (H. soloensis now variously viewed as 'advanced Homo erectus' or as separate species; JTvG))

Oppenoorth, W.F.F. (1932)- Een nieuwe fossiele mensch van Java. Tijdschrift Koninklijk Nederlands Aardrijkskundig Genootschap (2) 49, p. 704-707.
(online at: https://resolver.kb.nl/resolve?urn=MMUBA13:001678001:pdf)
('A new fossil man from Java'. New hominid skull from Solo River terrace deposits near Ngandong, N of Ngawi, C Java. Named Homo (Javanthropus) soloensis)

Oppenoorth, W.F.F. (1932)- Ein neuer diluvialer Urmensch von Java. Natur und Museum, Senckenberg, Frankfurt, 62, 9, p. 269-279.
('A new Pleistocene ancient hominid from Java'. Similar to above papers on Ngandong 'Solo Man' discovery)

Oppenoorth, W.F.F. (1936)- Een prehistorisch cultuurcentrum langs de Solo Rivier. Tijdschrift Koninklijk Nederlands Aardrijkskundig Genootschap (2) 53, p. 399-411.
(online at: https://resolver.kb.nl/resolve?urn=MMUBA13:001680001:pdf)
('A prehistoric culture center along the Solo River'. On mammal and hominid remains in Pleistocene fluvial terraces near Ngandung, Solo River downstream of Ngawi, E Java. In addition to bones, also various man-made tools made from bones)

Oppenoorth, W.F.F. (1937)- The place of *Homo soloensis* among fossil men. In: G.G. Mac-Curdy (ed.) Early Man, Lippincott Co., New York., p. 348-360.
(Homo soloensis from Ngandong, Java, viewed as oldest known representatives of 'Homo sapiens fossillis'. (Weidenreich 1943 viewed it as transition between Pithecanthropus and modern man; Koenigswald (1956) suggested Ngandong specimens too recent and advanced to be Pithecanthropus, but closer to Neandertals))

Orchiston, D.W. & W.G. Siesser (1982)- Chronostratigraphy of the Plio-Pleistocene fossil hominids of Java. Modern Quaternary Research in Southeast Asia 7, Balkema, Rotterdam, p, 131-149.
(Review of chronostratigraphy of hominid-bearing formations of C and E Java. Most radiometric dates without well documented stratigraphic position or analytical uncertainty, and of dubious value. Most reliable date is 0.83 Ma for pumice in Lower Kabuh Fm. Underlying marine U Kalibeng Fm Late Pliocene microfaunas)

O'Sullivan, P.B., M. Morwood, D. Hobbs, F. Aziz, Suminto, M. Situmorang, A. Raza & R. Maas (2001)- Archaeological implications of the geology and chronology of the Soa Basin, Flores, Indonesia. Geology (GSA) 29, 7, p. 607-610.
(Zircon fission-track dates of tuffaceous deposits associated with stone artifacts attributed to Homo erectus in Soa lacustrine basin on Flores indicate early hominids must have begun colonizing E Indonesia by ~840 ka)

Oxnard C., P.J. Obendorf, B.J. Kefford & J. Dennison (2010)- Post-cranial skeletons of hypothyroid cretins show a similar anatomical mosaic as *Homo floresiensis*. PloS One 5, 9, e13018, p. 1-11.
(online at: http://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0013018&type=printable)
(LB1and LB6 skulls of 'Homo floriensis' most likely, endemic cretins from population of unaffected Homo sapiens. Consistent with recent hypothyroid endemic cretinism throughout Indonesia, including nearby Bali)

Oxnard C., P.J. Obendorf, B.J. Kefford & J. Dennison (2012)- More on the Liang Bua finds and modern human cretins. Homo 63, 6, p. 407-412.
(More on recent discoveries of Flores island hominids)

Patole-Edoumba, E., A.F. Pawlik & A.S. Mijares (2012)- Evolution of prehistoric lithic industries of the Philippines during the Pleistocene. Comptes Rendus Palevol 11, 2-3, p. 213-230.
(online at: https://www.sciencedirect.com/science/article/pii/S1631068311001230)
(Review of major Pleistocene- Holocene lithic (stone tool) industries in Philippines)

Pearch, M.J., S. Bumrungsri, J.L. Schwenninger, D.J. Ward & D.L. Harrison (2013)- A review of the Cainozoic small mammal fauna of Thailand with new records (Chiroptera; Scandentia; Eulipotyphla) from the late Pleistocene. Cainozoic Research 10, 1-2, p. 59-98.
(Fifteen Late Pleistocene small mammal species from cave in Ordovician limestone hill in S Thailand)

Perston, Y.L., M. Moore, Suryatman, M. Langley, B. Hakim, A.A. Oktaviana & A. Brumm (2021)- A standardised classification scheme for the Mid-Holocene Toalean artefacts of South Sulawesi, Indonesia. PLoS One 16(5): e0251138, p. 1-37.
(online at: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0251138)

Petraglia, M.D., P. Ditchfield, S. Jones, R. Korisettar & J.N. Pal (2012)- The Toba volcanic super-eruption, environmental change, and hominin occupation history in India over the last 140,000 years. Quaternary International 258, p. 119-134.
(Middle Paleolithic hominins in India appear to have survived the potentially catastrophic effects of the large Toba volcanic eruption in N Sumatra and the resulting climatic fluctuations in Late Pleistocene)

Pickford, M., H. Nakaya, Y. Kunimatsu, H. Saegusa, A. Fukuchi & B. Ratanasthien (2004)- Age and taxonomic status of the Chiang Muan (Thailand) hominoids. Comptes Rendus Palevol 3, 1, p. 65-75. *(online at: www.academia.edu/7814204/Age_and_taxonomic_status_of_the_Chiang_Muan_Thailand_hominoids)*
(Age of Lufengpithecus chiangmuanensis Chaimanee 2003 originally estimated as ~13-13.5 Ma, but previous studies suggest age closer to ~12-11 Ma. L. chiangmuanensis synonym of L. keiyuanensis Wu)

Polanski, J.M., H.E. Marsh & S.D. Maddux (2016)- Dental size reduction in Indonesian *Homo erectus*: implications for the PU-198 premolar and the appearance of *Homo sapiens* on Java. J. Human Evolution 90, p. 49-54.
(Recent recovery of hominin maxillary third premolar, PU-198 in collections from Punung Cave (E Java) was used to suggest Homo sapiens appeared on Java between 143-115, ka. However, PU-198 overlaps in premolar sizes between H. erectus and H. sapiens, and indicate reduction in premolar size between early and late Javan H. erectus. Question appearance of H. sapiens on Java between 143-115 ka)

Polhaupessy, A.A. (1999)- Palynological evidence for a Pleistocene environment in Trinil, East Java. Proc. 35th Session Coord. Comm. Coastal Offshore Geoscience Programmes E and SE Asia (CCOP), Subic Bay 1998, 2, Technical Reports, p. 299-308.

(Two pollen zones in Pleistocene of Trinil (site of first Homo erectus in C Java): (1) Pucangan Fm rel. low diversity grass-dominated vegetation and lacustrine fresh water plants; (2) Lower Kabuh Fm higher diversity mixed freshwater swamp and terrestrial vegetation, perhaps suggesting slightly drier climate than today)

Polhaupessy, A.A. (1999)- Quaternary palynological study of the Trinil area, East Jawa. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 9, p. 1-7.

Polhaupessy, A.A. (2006)- Environment of early man in Java. In: T. Simanjuntak et al. (eds.) Archaeology: Indonesian perspective, R.P. Soejono's Festschrift, LIPI Press, Jakarta, p. 83-101.
(Review of palynological analyses of Quaternary deposits of Java, of sites with hominids or other vertebrate fossils (Bumiayu (Cisaat), Sangiran, Trinil, Solo and Madiun areas, etc.) Many Pleistocene sites with abundant Graminae (grass) pollen, suggesting more seasonal climate with savannah grasslands. Pollen during Last Glacial Maximum with common Dacrycarpus, etc., suggest temperatures several degrees cooler than today)

Pop, E., S. Hilgen, S. Adhityatama, H. Berghuis, T. Veldkamp, H. Vonhof, I. Sutisna, G.Alink et al. (2023)- Reconstructing the provenance of the hominin fossils from Trinil (Java, Indonesia) through an integrated analysis of the historical and recent excavations. J. Human Evolution 176, 103312, p. 1-26.
(online at: https://www.sciencedirect.com/science/article/pii/S0047248422001725)
(Detailed fieldwork of Dubois' original Trinil excavation site suggest Dubois excavated multiple fossiliferous channel deposits of different ages (Bone-Bearing Channel 1, 830-773 ka; Bone-Bearing Channel 2, 560-380 ka), resulting in collection of mixed hominin and other fossil assemblages)

Pope, G.G. (1982)- Hominid evolution in East and Southeast Asia. Ph.D. Thesis University of California, Berkeley, p. 1-375.

Pope, G.G. (1983)- Evidence on the age of the Asian Hominidae. Proc. National Academy of Sciences USA 80, 16, p. 4988-4992.
(online at: https://www.pnas.org/doi/epdf/10.1073/pnas.80.16.4988)
(All known Asian hominids <1 Million years old, and all early Asian hominids can be accommodated in Homo erectus. Entire record of Homo erectus in Asia may only span period of ~0.9-0.3 Ma. Maximal exposure of continental shelves such as Sunda occurred at ~3 Ma, 1.25 Ma and 0.65-0.45 Ma. Exposures at 1.25 Ma and 0.65-0.45 Ma most likely provided opportunities for migration of hominids and Pleistocene mammals to Java)

Pope, G.G. (1995)- The influence of climate and geography on the biocultural evolution of the Far Eastern hominids. In: E.S. Vrba et al. (eds.) Paleoclimate and evolution, with emphasis on human origins, Yale University Press, New Haven, p. 493-506.

Pope, G.G., S. Barr, A. Macdonald & S. Nakabanlang (1986)- Earliest radiometrically dated artifacts from Southeast Asia. Current Anthropology 27, 3, p. 275-
(M. Pleistocene stone artifact horizon at Ban Mae Tha, Lampang, Thailand below basalt flow dated as 0.8 et 0.6 Ma with K-Ar and paleomagnetic analysis)

Pope, G.G. & J.E. Cronin (1984)- The Asian hominidae. J. Human Evolution 13, 5, p. 377-396.
(Majority of known Asian hominids less than 1.0 Ma old, with maximum age 1.3 Ma. All can be incorporated in Homo erectus or Homo sapiens. Asian fossil records suggest gradual change over ~1 Million years)

Pope, K.O. & J.E. Terrell (2008)- Environmental setting of human migrations in the Circum-Pacific region. J. of Biogeography 35, p. 1-21.
(Rapid expansion of modern humans from Africa into SE Asia along coastal routes facilitated by period of stable climate and sea level from ~45,000- 40,000 yr BP, enabling them to reach coasts of NE Russia and Japan by 38,000- 37,000 yr BP)

Powley, M. J., I. Sutisna, K.M. Mikac, U.P. Wibowo & G.D. van den Bergh (2021)- The *Stegodon* bonebed of the Middle Pleistocene archaeological site Mata Menge (Flores, Indonesia): taphonomic agents in site formation. Quaternary (MDPI) 4, 4, p. 1-20.
(online at: https://www.mdpi.com/2571-550X/4/4/31/htm)
(M Pleistocene fluvial channel site of upper fossil-bearing interval at Mata Menge in Soa Basin, Flores, yielded earliest fossil evidence for Homo floresiensis in association with stone artefacts and endemic insular vertebrate fauna (Stegodon, giant rats, Komodo dragons, etc). Taphonomic analysis of Stegodon bones suggest deposition in stream channel)

Prasetyo, B. (2014)- Perkembangan budaya Akhir Pleistosen- Awal Holosen di Nusantara. Kalpataru, Majalah Arkeologi 23, 1, p. 1-15.
(online at: http://jurnalarkeologi.kemdikbud.go.id/index.php/kalpataru/article/view/47)
('The cultural development during Late Pleistocene-Early Holocene in the Indonesian Archipelago'. Review of Late Pleistocene- Early Holocene human cultures in various parts of Indonesia)

Prideaux, G. J., I.A.R. Kerr, J.D. van Zoelen, R. Grun, S. van der Kaars, A. Oertle, K. Douka et al. (2022)- Re-evaluating the evidence for late-surviving megafauna at Nombe rockshelter in the New Guinea highlands. Archaeology in Oceania 57, 3, p. 223-248.
(online at: https://onlinelibrary.wiley.com/doi/10.1002/arco.5274)
(Nombe rock shelter in Porol Range in highlands of PNG with large marsupial fossils (diprotodontid, kangaroo, thylacine) dated at 27-22 ka ago, i.e. well after arrival of humans (and after ~40 ka extinction of Pleistocene megafauna in Australia))

Přikryl, T., A. Castro, A.G. Fernando, J.R.C. Nogot, C. Magtoto, K. Garas, D. Mediodia & C.H. Lin (2025)- Fossil fish assemblage of the Laguna Formation, Philippines: unveiling the uniqueness of Pleistocene freshwater ecosystems in Southeast Asia. Swiss J. of Palaeontology 144, 5, p. 1-16.
(online at: https://sjpp.springeropen.com/articles/10.1186/s13358-024-00347-0)
(First description of well-preserved, diverse, unique Pleistocene fossil freshwater fish assemblage from lacustrine facies of Laguna Fm in Rizal Province, S Luzon Island, Philippines. Associated with Stegodon fossils, etc.)

Purnomo, A. (2007)- La sedimentation du lac de Guyang Warak (Punung-Java Est, Indonesie). Master Thesis, Museum National Histoire Naturelle, Paris, C.E.R.P. de Tautavel, p. 1-83.
(online at: http://hopsea.mnhn.fr/pc/thesis/Purnomo_Andri_2007.pdf)
('Sedimentation of Guyang Warak lake (Punung, East Java, Indonesia'. South Central Java'. Study of 'Grenzbank' and Kabuh Formation)

Purnomo, A. (2008)- Stratigraphie et sedimentation au Sud Est de Dome de Sangiran: l'environnement des *Homo erectus* au debut de Pleistocene Moyen. Thesis Universita degli Studi di Ferrara, Italy, p. 1-252 + 34
(online at: http://eprints.unife.it/831/1/Purnomo_tesi.pdf)
('Statigraphy and sedimentation in the SE of Sangiran Dome: the environment of Homo erectus at the start of the Middle Pleistocene'))

Purnomo, A. (2008)- The sedimentation of Lake Guyang Warak (Punung-East Java, Indonesia). Annali dell'Universita degli Studi di Ferrara, Museologia Scientifica Naturalistica, Special Volume 2008, p. 151-154.
(Lake Guyang Warak, Punung, NW of Pacitan in S Mountains of C Java, close to famous Paleolithic Site Song Terus Cave. 6m long core shows almost same environment from at least 2000 BP)

Purnomo, A. (2013)- Stratigraphie et sedimentation au Sud Est de Dome de Sangiran: l'environnement des *Homo erectus* au debut de Pleistocene moyen. Ph.D. Thesis Universita degli Studi di Ferrare, Italy, p. 1-251.
(online at: http://eprints.unife.it/831/1/Purnomo_tesi.pdf)
(Stratigraphy and sedimentation at the SE Sangiran Dome: the environment of Homo erectus at the start of the Middle Pleistocene'. Depositional environments of Upper Pucangan- Grenzbank- Kabuh Formations, identification of soil horizons with human occupation, frequent volcanic eruptions, etc.))

Purnomo, A., F. Semah, A.M. Semah & T. Simanjuntak (2014)- Geological structure, sedimentation dynamics and prehistory in the Southeastern part of the Sangiran Dome (Java-Indonesia): research and conservation strategies. In: N. Amano et al. (eds.) Southeast Asia: Human evolution, dispersals and adaptations, 17th Congress International Union Prehistoric and Protohistoric Sciences (UISPP), Burgos 2014, p. 94-99.

Purnomo, G.A., S. Kealy, S. O'Connor, A. Schapper, B. Shaw, B. Llamasa, J.C. Teixeira, H. Sudoyo & R. Tobler (2024)- The genetic origins and impacts of historical Papuan migrations into Wallacea. Proc. National Academy of Sciences USA (PNAS) 121, 52, e2412355121, p. 1-10.
(online at: https://www.pnas.org/doi/epdf/10.1073/pnas.2412355121)
(Tropical archipelago of Wallacea (E Indonesia) was first settled by anatomically modern humans (AMH) by 50 ka. Descendent populations thought to have remained genetically isolated prior to arrival of Austronesian seafarers around 3.5 ka. Modern Wallaceans exhibit Papuan- and Asian-related ancestries. New genomes from 12 Wallacean and three West Papuan populations confirm thatmost Papuan-related ancestry in Wallacea comes from prehistoric migrations originating in New Guinea and only aminor fraction is attributable to founding AMH settlers. Genes from Austronesian seafarers now ~40- 85% of modern Wallacean ancestry)

Puspaningrum, M.R. (2016)- Proboscidea as palaeoenvironmental indicators in Southeast Asia. Ph.D. Thesis, University of Wollongong, p. 1-318.
(online at: https://ro.uow.edu.au/theses/4944)
(Stable C and O isotope analyses on elephantoid tooth enamel from localities in mainland Asia (Linxia Basin, Siwalik, Irrawaddy, Tha Chang, Laos), continental islands (Java, Sumatra) and oceanic islands (Sulawesi, Sangihe, Luzon, Flores, Timor, Sumba). Diets in M Miocene- Pliocene and latest M Pleistocene- Recent dominantly (leaf) browsing, while Pliocene- M Pleistocene taxa mainly (grass) grazers)

Puspaningrum, M.R., I.P. Anwar, C. Hertler, E. Holzchen, S. Kruger, A.T. Hascaryo, J.O. Reschke & A.A. Bruch (2024)- Living in Sangiran: A spatial reconstruction of hominin environment in Java at 1 Ma. Earth History and Biodiversity 1, 100001, p. 1-13.
(online at: https://www.sciencedirect.com/science/article/pii/S2950475924000017)
(Reconstruction of paleoenvironmental conditions encountered by Homo erectus in (East) Java at 1 Ma (age of the 'Grenzbank' horizon) (paleotopography, drought map, vegetation map). Mainly lowland monsoonal forests landscape in E Java. Sangiran at 1 Ma possibly slightly cooler and longer dry season)

Puspaningrum, M.R., G.D. van den Bergh, A.R. Chivas, E. Setiabudi & I. Kurniawan (2020)- Isotopic reconstruction of Proboscidean habitats and diets on Java since the Early Pleistocene: implications for adaptation and extinction. Quaternary Science Reviews 228, 106007, p. 1-19.
(Six Proboscidean taxa found in Java extending back to Early Pleistocene, successively: Stegoloxodon indonesicus, Sinomastodon bumiajuensis, pygmy Stegodon, Stegodon trigonocephalus, Elephas hysudrindicus and Elephas maximus. C and O isotope analyses suggest earliest fauna (S. indonesicus) feeding in closed canopy rainforest during earliest Pleistocene. Towards late E Pleistocene (S. bumiajuensis) adapted to increasingly drier grassy habitats. Proboscidea in Ngandong Fauna suggests fragmented dense evergreen forests or woodland vegetation reappeared towards end M Pleistocene)

Puspaningrum, M.R., A.R. Chivas, I. Kurniawan, U.P. Wibowo, Y. Zaim & G.D. van den Bergh (2021)- Isotopic reconstruction of Proboscidean habitats and diets on enigmatic island of Sulawesi. Berita Sedimentologi 47, 3, p. 79-80. *(Abstract only)*
(online at: https://journal.iagi.or.id/index.php/FOSI/article/view/364/333)
(Stegoloxodon celebensis and Stegodon sompoensis flexible feeders and able to adapt to different niches, from closed canopy forest to open vegetation, while diets of Stegodon sp. B, Celebochoerus heekereni and Palaeoloxodon namadicus suggest they were more specialized)

Puymerail, L., C.B. Ruff, L. Bondioli, H. Widianto, E. Trinkaus, R. Macchiarell (2012)- Structural analysis of the Kresna 11 *Homo erectus* femoral shaft (Sangiran, Java). J. Human Evolution 63, p. 741-749.
(online at: www.academia.edu/13028927/Structural_analysis_of_the_Kresna_11_Homo_erectus_femor Etc.)

(Microtomographic anatomical analyses of Kresna 11, a late E Pleistocene adult H. erectus femoral shaft from 'Grenzbank' horizon at Ngebung site in Sangiran Dome, C Java, discovered in 1992)

Rabett, R.J. & P.J. Piper (2012)- The emergence of bone technologies at the end of the Pleistocene in Southeast Asia: Regional and evolutionary implications. Cambridge Archaeological Journal 22, 1, p. 37-56.
(Unlike during the Paleolithic of Europe, in SE Asia there was shift in technological usage from stone to bone (from a minor component to a medium of choice) during second half of Last Glacial Termination and into Holocene. Suggest this was inventive behaviour demanded of foraging communities)

Raden Saleh (1867)- Over fossiele beenderen van den Pandan. Natuurkundig Tijdschrift voor Nederlandsch-Indie 29, p. 423, 428, 434-435, 449-451 and p. 457-458.
('On fossil bones from the Pandan', C Java. Brief reports of telegrams from Raden Saleh about occurrences of large mammal bones in C Java region, but with little detail. Crates with fossil bones sent to 'Bataviaasch Genootschap van Kunsten en Wetenschappen '. (Pleistocene mammals re-described by K. Martin (1886, 1888); localities SW of Mt. Pandan now commonly known as Kedung Brubus; HvG)

Rahardjo, A.T. (1999)- Perubahan iklim dan batas umur Pliosen-Plistosen berdasarkan analisis foraminifera dan palinologi di daerah Mojoroto, Mojokerto- Jawa Timur. Buletin Geologi (ITB) 3, 1, p. 1-13.
('Climate change and Pliocene-Pleistocene age boundary based on foraminifera and palynology analysis in the Mojoroto area, Mojokerto, East Java')

Rahardjo, A.T. & A.M. Semah (1989)- Penelitian palynology daerah Sangiran. Bull. Dept. Geologi Inst Technologi Bandung (ITB) 1983, 9, p. 23-31.
('Palynology research in the Sangiran area')

Rasmussen, M., X. Guo, Y. Wang, K.E. Lohmueller, S. Rasmussen, A. Albrechtsen et al. (2011)- An aboriginal Australian genome reveals separate human dispersals into Asia. Science 334, 6052, p. 94-98.
(DNA genomic sequence of Aboriginal Australians shows they are descendants of early human dispersal into E Asia, ~62,000- 75,000 yrs ago, separate from dispersal of ancestors of modern Asians 25,000- 38,000 yrs ago)

Reepmeyer, C., S. O'Connor & S. Brockwell (2011)- Long-term obsidian use at the Jerimalai rock shelter in East Timor. Archaeology in Oceania 46,2, p. 85-90.
(online at: https://openresearch-repository.anu.edu.au/server/api/core/bitstreams/47cc4f55-47a4-4882-94b6-166c22b3536f/content)
(Obsidian artefacts from Jerimalai and other rock shelters in E Timor demonstrate exploitation of this rock type began in Pleistocene by 42 ka, and continued periodically into M-L Holocene. High-silicate obsidian from unknown location apparently transported over considerable distance to sites in E of Timor)

Reich, D., N. Patterson, M. Kircher, F. Delfin, M.R. Nandineni, I. Pugach, A.M. Ko, Y.C. Ko et al. (2011)- Denisova admixture and the first modern human dispersals into Southeast Asia and Oceania. American J. Human Genetics 89, 4, p. 516-528.
(online at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3188841/pdf/main.pdf)
(Ancestors of New Guineans, Aboriginal Australians, Near Oceanians, Polynesians, Fijians, E Indonesians, and Mamanwa ('negrito' group from Philippines) inherited part of ancestry from Denisovans, an archaic hominin group from Siberia. However, mainland E Asians, W Indonesians, Jehai (negrito group from Malaysia), and Onge (negrito group from Andaman Islands) have not)

Reis, K.R. & A.M. Garong (2001)- Late Quaternary terrestrial vertebrates from Palawan Island, Philippines. Palaeogeogr. Palaeoclim. Palaeoecology 171, p. 409-412.
(online at: http://users.clas.ufl.edu/krigbaum/6930/Reis&Garong_p3_2001.pdf)
(Humans first colonized Palawan at ~40,000 yr BP. Absence of large carnivores and primates suggests more insular affinity than Borneo, but higher diversity than truly oceanic islands in Late Quaternary)

Reynolds, T. & G. Barker (2014)- Reconstructing Late Pleistocene climates and human activities in northern Borneo from excavations in the Niah caves. In: Y. Kaifu et al. (eds.) Emergence and diversity of modern human behavior in Paleolithic Asia, Peopling of the Americas Publications, Texas A&M University Press, p. 140-157.
(Niah Caves in Sarawak home to oldest anatomically modern 'Deep Skull', now confidently dated as ~37.5 ka. First evidence for associated human activity at caves site goes back to ~50 ka)

Rightmire G.P. (1984)- Comparisons of *Homo erectus* from Africa and Southeast Asia. Courier Forschungsinstitut Senckenberg 69, p. 83-98.

Rightmire G.P. (1985)- The tempo of change in the evolution of mid-Pleistocene *Homo*. In: E. Delson (ed.) Ancestors: the hard evidence, New York, p. 255-264.

Rightmire G.P. (1988)- *Homo erectus* and Later Middle Pleistocene humans. Annual Review of Anthropology 17, p. 239-259.
(Review of worldwide occurrences of H. erectus. No ties between H. erectus and modern humans)

Rightmire, G.P. (1990)- The evolution of *Homo erectus*: comparative anatomical studies of an extinct species. Cambridge University Press, Cambridge, p. 1-276.
(Book including reviews of Homo erectus in Indonesia (Trinil, Sangiran, Sambungmacan, Ngandong))

Rightmire G.P. (1992)- *Homo erectus*: ancestor or evolutionary side branch? Evolutionary Anthropology 1, p. 43-49.

Rightmire G.P. (1994)- The relationship of *Homo erectus* to later Middle Pleistocene hominids. Courier Forschungsinstitut Senckenberg 171, p. 319-326.

Rightmire G.P. (1998)- Human evolution in the Middle Pleistocene: the role of *Homo heidelbergensis*. Evolutionary Anthropology 6, p. 218-227.

Rightmire G.P. (2001)- Patterns of hominid evolution and dispersal in the Middle Pleistocene. Quaternary International 75, p. 77-84.
(At onset of Quaternary Homo erectus spread across Old World from Africa. Populations persisted in Far East until late in M Pleistocene, while H. erectus disappeared relatively early in West. Episode of hominid speciation in mid-Quaternary gave rise to anatomically more modern hominids called Homo heidelbergensis. Relationships of H. heidelbergensis to Neanderthals and recent humans still need clarification)

Rizal, Y. (1998)- Die Terrassen entlang des Solo-Flusses in Mittel- und Ost-Java. Thesis Universitat zu Koln, Cologne, Germany, p. 1-189. *(Unpublished)*
('The terraces along the Solo River in Central and East Java')

Rizal, Y. (2005)- The ages of the Solo Terraces at the Ngancar and Ngandong Region, Middle Jawa, Indonesia. *(Abstract only)*
(online at: http://hopsea.mnhn.fr/doc/2005QP11abstracts.pdf)
(Best estimates of ages of Pleistocene terraces along Solo River: High-terraces ~47 ka, Middle-terraces ~20 ka and Lower-terraces 1.65 + 1.5 ka)

Rizal, Y., W.D. Santoso, A. Rudyawan, R.A. Tampubolon & A.A. Nurfahan (2018)- Sedimentary facies and hydrocarbon reservoir potential of sand flat in the upper part of Tapak Formation in Banyumas area, Central Java. J. RISET Geologi dan Pertambangan (LIPI) 28, 2, p. 251-263.
(online at: http://jrisetgeotam.com/index.php/jrisgeotam/article/view/835/pdf)
(Sedimentology and trace fossils of upper Tapak Fm in Kali Cimande in Banyumas Basin, C Java, suggest tidal flat environments)

Rizal, Y., K.E. Westaway, Y. Zaim, G.D. van den Bergh, E.A. Bettis, M.J. Morwood, O.F. Huffman, R. Grun, R. Joannes-Boyau, R.M. Bailey, Sidarto, M.C. Westaway, I. Kurniawan et al. (2020)- Last appearance of *Homo erectus* at Ngandong, Java, 117,000-108,000 years ago. Nature 577, p. 381-385.
(online at: www.researchgate.net/publication/338021539_Last_appearance_of_Homo_erectus_at_Ngandong_Java_117000-108000_years_ago)
(Twelve H. erectus calvaria (skull caps) and two tibiae (lower leg, shn bones) excavated between 1931-1933 from bone bed in terrace 20m above Solo River at Ngandong, and are youngest form of H. erectus (at that time named Homo soloensis, 'Solo Man'). Bayesian modelling of 52 radiometric ages (U-series dating of speleothems to constrain regional landscape evolution; 40Ar/39Ar and U dating to constrain sequence of terrace evolution, etc.) At least by 0.5 Ma Solo River diverted into Kendeng Hills, and formed Solo terrace sequence between 316 and 31 ka, and Ngandong terrace between ~140- 92 ka. Modelled ages of H. erectus bone bed 117-108 ka)

Rizal, Y., Y. Zaim, Y. Aswan, C. Hertler, S.Mulyaningsih & N. Triany (2009)- Geologi dan paleontologi vertebrata daerah Jembarwangi dan sekitarnya, Kabupaten Sumedang, Jawa Barat. Buletin Geologi (ITB) 39, 2, p. 95-116.
('Geology and vertebrate paleontology of the Jembarwangi area and surroundings, Sumedang Dustrict, West Java')

Rizal, Y., Y. Zaim & Y. Iriani (2005)- Late Tertiary fossil whale from Surade, South Sukabumi, West Java. Buletin Geologi (ITB) 37, 1, p. 29-34.
(Late Miocene whale fossils from Surade, ~60km S of Sukabumi. First record of whale fossils in Indonesia)

Roberts, P. & N. Amano (2019)- Plastic pioneers: hominin biogeography east of the Movius Line during the Pleistocene. Archaeological Research in Asia 17, p. 181-192.
(online at: https://os.pennds.org/archaeobib_filestore/pdf_articles/ArchRAsia/2019_RobertsAmano.pdf)
(E-M Pleistocene Homo erectus and its diminutive descendant(?) Homo floresiensis, was able to move East of the (now defunct) 'Movius Line' and the biogeographic Wallace Line. Migration followed alongside woodland- and grassland-adapted fauna, with some evidence for use of lacustrine and riverine tropical forest settings. By contrast, Homo sapiens, appeared in SE Asia during expansion of rainforest fauna (like today) and adapted to a diversity of tropical rainforest settings. It also developed specialized maritime subsistence strategies that aided its rapid expansion beyond the Wallace Line and Sahul since 45 ka or earlier)

Roberts, P., J. Louys, J. Zech, C. Shipton, S. Kealy, S.S. Carro, S. Hawkins, C. Boulanger et al. (2020)- Isotopic evidence for initial coastal colonization and subsequent diversification in the human occupation of Wallacea. Nature Communications 11, 2068, p. 1-11.
(online at: https://www.nature.com/articles/s41467-020-15969-4.pdf)
(Stable carbon and oxygen isotope analysis of human and faunal tooth enamel reflect diets. Earliest humans to arrive in Wallacea (E Indonesia archipelago) seemingly specialized in use of coastal-marine resources. Rapid initial coastal colonization followed by inland settlement. After 20,000 years BP, diversification in human resource use across Wallacea. On Timor, tropical forest environments remained prevalent in past 45,000 years)

Roberts, R.G., K.E. Westaway, J.X. Zhao, C.S.M.Turney, M.I. Bird, W.J. Rink & L.K. Fifield (2009)- Geochronology of cave deposits in Liang Bua and of adjacent river terraces in the Wae Racang valley, western Flores, Indonesia: a synthesis of age estimates for the type locality of *Homo floresiensis*. J. Human Evolution 57, 5, p. 484-502.
(Seven dating methods used for cave and nearby terrace deposits at Liang Bua. Hominin- and artifact-bearing deposits spanning the last ~100,000 years)

Rolland, N. (2002)- The initial hominid colonization of Asia: a survey of anthropic evidence from Biogeographic and ecological perspectives. Bull. Indo-Pacific Prehistory Association 22 (Melaka Papers, vol. 6), p. 3-15.
(online at: https://journals.lib.washington.edu/index.php/BIPPA/article/viewFile/11800/10428)

Rothpletz, W. (1951)- Alte Siedlungsplatze bei Bandung (Java) und die Entdeckung bronzezeitlicher Gussformen. Sudsee-Studien, Museum fur Volkerkunde, Basel, p. 77-126.
('Old settlement places near Bandung and the discovery of Bronze Age moulds'. Report of 1941-1947 fieldwork by Swiss national Rothpletz around former Lake Bandung, with occurrences of Bronze Age obsidian artefacts and ceramic shards, especially near Dago, N of Bandung. Artefacts of obsidian industry studied by Bandi (1951))

Rozzi, R. (2017)- A new extinct dwarfed buffalo from Sulawesi and the evolution of the subgenus Anoa: an interdisciplinary perspective. Quaternary Science Reviews 157, p. 188-205.
(open manuscript online at: https://www.sciencedirect.com/science/article/abs/pii/S0277379116306795)
(New species of dwarf buffalo (anoa) Bubalus grovesi, from Late Pleistocene/Holocene of Lamoncong area, S Sulawesi (based on material collected by P. van Stein Callenfels in 1934 and stored in Naturalis, Leiden)

Rozzi, R. & M.R. Palombo (2013)- The morphology of femur as palaeohabitat predictor in insular bovids. Boll. Societa Paleontologica Italiana 52, 3, p. 177-186.
(online at: https://pdfs.semanticscholar.org/3357/23a5ad5a4c439bd39897170ce3d7a26cdbfd.pdf)
(Morphology of long bones of antelope-like insular fossil bovid Duboisia santeng (Dubois, 1891) from E-M Pleistocene Java suggests it was forest dweller)

Rozzi, R., D.E. Winkler, J. De Vos, E. Schulz & M.R. Palombo (2013)- The enigmatic bovid *Duboisia santeng* (Dubois, 1891) from the Early-Middle Pleistocene of Java: a multiproxy approach to its palaeoecology. Palaeogeogr. Palaeoclim. Palaeoecology, 377, p. 73-85.
(online at: https://www.academia.edu/download/98314429/j.palaeo.2013.03.01220230206-1-pmviie.pdf)
(Faunal elements of Indo-Chinese origin entered Java during Pleistocene. Most Javanese mammalian taxa close or identical to mainland relatives, like fossil bovids, common in E-M Pleistocene 'Stegodon-Homo erectus fauna' (Bubalus palaeokerabau, Bibos palaesondaicus, Epileptobos groeneveldtii). Javanese bovid Duboisia santeng from Sangiran regarded as insular endemic species due to small size, but present in late M Pleistocene of what is now Peninsular Malaysia (also in Thailand; Nishioka et al. 2018; JTvG))

Ruff, C.B., L. Puymerail, R. Macchiarelli, J. Sipla & R.L. Ciochon (2015)- Structure and composition of the Trinil femora: functional and taxonomic implications. J. Human Evolution 80, p. 147-158.
(online at: www.academia.edu/33511287/Structure_and_composition_of_the_Trinil_femora_Functional Etc.)
(Trinil Femur I derives from more recent time period than calotte of Homo erectus 'holotype'. Less famous and less complete Femora II-V may represent H. erectus at Trinil)

Ruff, C.B., A.D. Sylvester, N.T. Rahmawati, R.A. Suriyanto, P. Storm, M. Aubert, R. Joannes-Boyau et al. (2022)- Two Late Pleistocene human femora from Trinil, Indonesia: implications for body size and behavior in Southeast Asia. J. Human Evolution 172, 103252, p. 1-19.
(online at: https://ir.cwi.nl/pub/32156/32156.pdf)
(Late Pleistocene hominin partial femora from Trinil dated to ~37-32 ka. Trinil 9 and 10 femora discovered by Prof. Teuku Jacob in 1978 along right bank of Solo River in 'T1' terrace deposits)

Saegusa, H., Y. Thasod & B. Ratanasthien (2005)- Notes on Asian stegodontids. Quaternary International 126-128, p. 31-48.
(Stegodontids (elephant-like proboscideans), flourished in Neogene and Quaternary of Asia. Significant recent finding of new stegodontid fossils at Nakhon Ratchasima, Thailand and Yuanmou Basin, Yunnan, show Late Miocene transition from stegolophodons to stegodons, suggesting stegodons originated in mainland Asia)

Saleki, H. (1997)- Apport d'une intercomparaison de methodes nucleaires (230Th/234U, ESR et 40Ar/39Ar) a la datation de couches fossiliferes pleistocenes dans le dome de Sangiran (Java, Indonesie). Ph.D. Thesis, Museum National Histoire Naturelle, Paris, p. 1-238. *(Unpublished)*
('Contribution to comparison of nuclear methods (230Th/234U, ESR and 40Ar/39Ar) to dating of Pleistocene fossiliferous beds in Sangiran dome, Java'. Proposed chronology for Sangiran dome: (1) volcanic breccia deposited between ~2.05 and 1.56 Ma, followed by Pucangan Fm; (2) Grenzbank in upper limit of Jaramillo

period at 0.9 Ma; (3) rapid sedimentation with archeological layer with some bones which supposedly burnt at ~0.8-0.9 Ma, before Brunhes/Matuyama magnetic reversal; (4) Notopuro mud-flow unit deposited at 150 ± 10 ka (incl. 0.8 Ma of tuff overlying Ngebung H. erectus occupation site))

Salles, T., C. Mallard, L. Husson, S. Zahirovic, .A.C. Sarr & P. Sepulchre (2021)- Quaternary landscape dynamics boosted species dispersal across Southeast Asia. Nature Communications Earth & Environment 2, 1, 240, p. 1-12.
(online at: https://www.nature.com/articles/s43247-021-00311-7.pdf)
(Study of Quaternary relief of Sunda Shelf and impact on species migrations. Region was predominantly terrestrial during Late Pleistocene. Four phases of drainage reorganisation and river captures)

Santa Luca, A.P. (1980)- The Ngandong fossil hominids: a comparative study of a Far eastern *Homo erectus* group. Yale University Publications in Anthropology 78, New Haven, p. 1-175.
(Morphological study of Ngandong hominid skulls from Solo River 20m terrace at Ngandong, originally described as Homo soloensis Oppenoorth 1932. Here interpreted as advanced forms of Homo erectus ('Homo erectus s.l.', not H. erectus s.s.') (conclusion followed by many subsequent researchers, but validity questioned in older (Dubois 1937, Von Koenigswald 1956) and more recent studies, e.g., Zaitoun et al. 2010?, M. Westaway et al. 2015)

Sarasin, P. (1914)- Neue lithochrone Funde im Innern von Sumatra. Verhandlungen Naturforschenden Gesellschaft Basel 25, p. 97-111.
(online at: www.zobodat.at/pdf/Verh-natforsch-Ges-Basel_25_1914_0097-0111.pdf)
(Obsidian and chert stone implements from Ngalau Ulu Tjangko (Tiangko) cave between the Merangin and Batang Tabir tributaries of the Jambi River, collected by A. Tobler, and donated to Basel Museum. Some of Sarasin's conclusions disputed by Zwierzycki (1926))

Sarasin, P. & F. Sarasin (1905)- Versuch einer Anthropologie der Insel Celebes: Die Toala-Hohlen von Lamontjong. In: Materialien zur Naturgeschichte der Insel Celebes, C.W. Kreidel's Verlag, Wiesbaden, vol. 5, 1, p. 1-62.
(online at: https://archive.org/details/materialienzurna512sara/page/n6/mode/1up)
('Attempt at an anthropology of the Island Celebes, 1, The Toala caves at Lamontjong'. First account of the well-known mid-late Holocene 'Toalean' cave dwelling people with Toala stone artefact culture in Sulawesi)

Sartono, S. (1961)- Notes on a new find of a *Pithecanthropus* mandible. Djawatan Geologi (Geological Survey), Bandung, Publikasi Teknik- Seri Paleontologi No. 2, p. 1-51.

Sartono, S. (1968)- Early man in Java: *Pithecanthropus* skull VII, a male specimen of *Pithecanthropus erectus*. Proc. Koninklijke Nederlandse Akademie van Wetenschappen B 71, 5, p. 396-422.

Sartono, S. (1969)- On the Plio-Pleistocene boundary of Java. Bull. of National Institute Geology and Mining (NIGM), Bandung 2, 2, p. 1-19.
(In Sangiran area of C Java Plio-Pleistocene boundary should be placed between Corbicula beds and Lower Volcanic Breccia bed. Based on mammalian fauna it should be placed between 'Unterer Wirbeltierhorizont' and 'Oberer Wirbeltierhorizont' of Von Koeningswald)

Sartono, S. (1969)- *Stegodon timorensis*: a pygmy species from Timor (Indonesia). Proc. Koninklijke Nederlandse Akademie van Wetenschappen, B72, p. 192-202.
(First description of Pleistocene dwarf elephant from Timor, collected by Verhoeven in 1964, 5 km E of Atambua. Showing potential migration routes from mainland Asia during Pleistocene)

Sartono, S. (1970)- On the stratigraphic position of *Pithecanthropus* mandible-C. Proceedings Institut Teknologi Bandung (ITB) (now J. Mathematical Fundamental Sciences), 4, 4, p. 91-102.
(online at: https://journals.itb.ac.id/index.php/jmfs/article/view/9737/3712)

(Pithecantropus C mandible collected in 1960 in Sangiran area, C Java, is from surface of Lower Pleistocene Putjangan beds. Encrusting matrix of mandible with planktonic (incl. Globorotalia tumida, Gr. crassa, Globigerinoides obliquus, Pulleniatina) and deeper marine smaller benthic foraminifera, presumably suggesting Lower Pleistocene age (NB: most forams in Sangiran fluvial Pleistocene deposits are reworked from underlying Kalibeng Fm deeper marine marls; JTvG) (see also Siesser & Orchiston 1978))

Sartono, S. (1971)- Observations on a new skull of *Pithecanthropus erectus* (*Pithecanthropus* VIII) from Sangiran, Central Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen B 74, 2, p. 185-194.

Sartono, S. (1972)- Discovery of another hominid skull at Sangiran, Central Java. Current Anthropology 13, 1, p. 124-126.
(New fossil Homo erectus skull discovered in 1969 at S flank Sangiran dome, at base of low bluff on S side of Putjung creek)

Sartono, S. (1973)- On an additional *Stegodon timorensis* Sartono. Direktorat Geologi, Bandung, Publikasi Teknik- Seri Paleontologie No. 3, p. 1-13.
(Description of additional Stegodon tooth, collected by Verhoeven from sandy conglomerate overlying marine claystones near Umaklaren (Atambua), W Timor)

Sartono, S. (1973)- On cranial measurements of *Pithecanthropus erectus* (*Pithecanthropus* VIII). Direktorat Geologi, Bandung, Publikasi Teknik- Seri Paleontologie No. 4, p. 1-21.

Sartono, S. (1973)- On Pleistocene migration routes of vertebrate fauna in Southeast Asia. In: B.K. Tan (ed.) Proc. Reg. Conf. Geology of Southeast Asia, Kuala Lumpur 1972, Bull. Geological Society Malaysia 6, p. 273-286.
(online at: https://gsmpubl.files.wordpress.com/2014/09/bgsm1973018.pdf)
(Pleistocene vertebrates in SE Asia originated on Asian continent and migrated during Pleistocene into the island archipelago of Indonesia. Migration followed two routes, a western route across Sundaland area, and an eastern route via Taiwan and Philippines towards Sundaland area and Sulawesi)

Sartono, S. (1974)- Observations on a newly discovered jaw of *Pithecanthropus modjokertensis* from the lower Pleistocene of Sangiran, central Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, B 77, 1, p. 26-31.

Sartono, S. (1975)- Implications arising from *Pithecanthropus* VIII. In: R.H. Tuttle (ed.) Palaeoanthropology, Mouton Publishers, The Hague, p. 327-360.
(Three Pithecanthropus types in C Java: P. erectus (M Pleistocene; Trinil beds), P. modjokertensis (E Pleistocene; Djetis Beds) and P. dubius (lowest Pleistocene of Sangiran). New Pithecanthropus fossil (P VIII) near-complete and rel. large cranium discovered in Kabuh Fm of Sangiran in 1969 and is different from typical Pithecanthropus morphology)

Sartono, S. (1976)- Genesis of the Solo terraces. Modern Quaternary Research in Southeast Asia 2, Balkema, Rotterdam, p. 1-21.
(Three well-known river terrace levels along transverse valley of Solo River, N of Ngawi, C-E Java: Upper Pleistocene High terrace (Ngandong; possibly coinciding with Riss glaciation), Low Terrace and Flood Terrace. Also three older, less well-known terraces identified, mainly from aerial photos. Highest terraces oldest (Early Pleistocene?) and 97m above Solo River. (see also Bartstra 1977))

Sartono, S. (1978)- The site of *Homo erectus* mandible F. Modern Quaternary Research in Southeast Asia 4, Balkema, Rotterdam, p. 19-24.
(Rel. complete Homo erectus trinilensis lower jaw from black claystones of Pucangan Fm in central part of Sangiran dome)

Sartono, S. (1979)- The discovery of a pygmy *Stegodon* from Sumba, East Indonesia: an announcement. Modern Quaternary Research in Southeast Asia 5, Balkema, Rotterdam, p. 57-63.
(First report of Pleistocene Stegodon mandible from Sumba (Watu Mbaka). Described as Stegodon sumbaensis)

Sartono, S. (1979)- The stratigraphy of the Sambungmacan site in Central Java. Modern Quaternary Research in Southeast Asia 5, Balkema, Rotterdam, p. 83-88.
(Gravels, sands and silts at small Sambungmacan site, off Solo River, C Java, in which Homo erectus cranium was found in 1973, now regarded as of part of Ngandong Fm of Late Pleistocene age. not E-M Pleistocene 'Grenzbank'/ Kabuh Fm- equivalent)

Sartono, S. (1979)- The age of the vertebrate fossils and artifacts from Cabenge in South Sulawesi, Indonesia. Modern Quaternary Research in Southeast Asia 5, Balkema, Rotterdam, p. 65-81.
(Fossiliferous terrace deposits of Walanae River most likely Pleistocene, possibly Holocene)

Sartono, S. (1980)- *Homo erectus ngandongensis*: the possible maker of the 'Sangiran flakes'. Anthropologie 18, 2/3, p. 121-131.
(Primitive stone artefacts named 'Sangiran flakes' limited to Top of Kabuh/ Base Notopuro Formations at Sangiran, none found in lower levels with hominid remains. Appear to be associated with more advanced Homo erectus ngandongensis)

Sartono, S. (1981)- Pre-*Homo erectus* population in Java, Indonesia. Proc. 10th Congreso Union Internacional de Ciencias Prehistoricas y Protohistoricas, Mexico 1981, p. 47-86.

Sartono, S. (1982)- Characteristics and chronology of early man in Java. In: H. and M. De Lumley (eds.) Compte Rendu du 1st Congres International de Paleontologie Humaine, Nice (France), CNRS, 2, p. 495-541.

Sartono, S. (1984)- Notes on the Pleistocene stratigraphy of Java, Indonesia. Modern Quaternary Research in Southeast Asia 8, Balkema, Rotterdam, p. 129-135.

Sartono, S. (1985)- Datings of Pleistocene man of Java, Indonesia. Modern Quaternary Research in Southeast Asia 9, Balkema, Rotterdam, p. 115-125.

Sartono, S. (1985)- Pleistocene peopling of the Southeast Asian Archipelago. Proc. 12th Indo-Pacific Prehistory Association (IPPA) Congress, Penablanca, Cagayan, Phillipines 1985, p. 1-12.

Sartono, S. (1986)- New lights on human evolution in Southeast Asia. In: G.H. Teh & S. Paramananthan (eds.) Proc. 5th Regional Congress Geology Mineral Energy Resources of SE Asia (GEOSEA V), Kuala Lumpur 1984, 2, Bull. Geological Society Malaysia 20, p. 269-288.
(online at: www.gsm.org.my/products/702001-101423-PDF.pdf)
(Until 1980 subdivision of Pleistocene human fossils as proposed by Von Koenigswald (1968) was used. New discoveries in last 5 years necessitate re-assessment. Earliest wave of human migration into SE Asia (Java) at ~1.8 Ma, coincided with onset of Gunz glacial. Before this period most of SE Asia region still inundated by sea, hampering movement of early humans from Asia)

Sartono, S. (1987)- Migrasi manusia Plistocen Indonesia: kaitannya dengan tektonik lempeng. In: Geologi Kuarter dan lingkungan hidup, Geological Research Development Centre (GRDC), Bandung, Special Publ. 7, p. 7-20.
('Pleistocene human migration in Indonesia: relation to plate tectonics')

Sartono, S. (1987)- The long trek to the South. In: N. Thiramongkol (ed.) Proc. Workshop on Economic geology, tectonics, sedimentary processes and environment of the Quaternary in Southeast Asia, Haid Yai, Thailand 1986, IGCP 218/ Chulalongkorn University, Bangkok, p. 193-212.
(online at: http://library.dmr.go.th/Document/Proceedings-Yearbooks/M_1/1986/5083...)

(Review of hominid material from Pleistocene of Java and latest Pliocene- Pleistocene migration routes of hominids into Indonesian region from Asia. Migrations into Java probably aided by Pleistocene sealevel lowstands and exposed Sunda Shelf. Migrations into East Indonesia and Philippines probably not until Late Pleistocene)

Sartono, S. (1987)- Influence of plate tectonics on dispersals of Quaternary faunas in Southeast Asia. Proc. Seminar on International developments in science, Deutscher Akademie Austauschedienst, Bandung, p. 1-13.

Sartono, S. (1990)- Short guide to Sangiran and Trinil, Java. Guidebook 14th Congress Indo-Pacific Prehistory Association (IPPA), Yogyakarta, p. 1-34.

Sartono, S. (1991)- A new *Homo erectus* skull from Ngawi, East Java. Bull. Indo-Pacific Prehistory Association (IPPA) 11, p. 23-35.
(online at: http://ejournal.anu.edu.au/index.php/bippa/article/view/596/585)

Sartono, S. (1991)- *Meganthropus paleojavanicus* v.K.: its place in human evolution. Fourth International Senckenberg Conference, Frankfurt 1991, p.

Sartono, S. (1991)- *Homo (Pithecanthropus) erectus*: le debat sans fin. L'Anthropologie 95, 1, p. 123-136.
('Homo erectus: the debate without end'. On taxonomic status of Homo (Pithecanthropus) erectus. Two groups of early men are found in Java: Homo (H. robustus and H. erectus) and Australopithecus (A. (Meganthropus) palaeojavanicus). Pleistocene hominids from SE Asia probably subjected to evolution while they migrated from Asia toward Australia. Two migration routes: (1) western (Sunda Shelf) and (2) northern (Philippines), both arriving in Wallacea, possibly also in Sahul (Australia-New Guinea)

Sartono, S. (1993)- Insularity by plate tectonics in Quaternary Indonesia. Buletin Jurusan Geologi ITB, Bandung, 23, 2, p. 1-20.

Sartono, S. (1996)- Java: diversity of Upper Pliocene- Pleistocene hominids. Buletin Geologi (ITB) 26, 1, p. 1-12.
(Final paper by Sartono; overview of hominids distribution, evolution, migration on Java. Age of youngest Pithecanthropus in Sangiran, C Java, ~0.78 Ma, oldest (from 'Grenzbank' ~1.0Ma. Two main Pleistocene migration routes proposed.)

Sartono, S. & D. Grimaud-Herve (1983)- Les parietaux de l'hominide Sangiran 31. L'Anthropologie 87, p. 465-468.
('The parietals of the hominid Sangiran 31'. Recent discovery of Sangiran 31 skull in lower Pucangan Fm allowed us to compare biparietal vault with that of all known hominids from this deposit. Its extraordinary thickness and marked robustness similar to Sangiran 4 fossil man)

Sartono, S. & S. Hadiwisastra (1983)- Fosil vertebrata Plistosen di Busur Banda: implikasi struktural. Proc. 12th Annual Conv. Indonesian Association Geologists (IAGI), p. 153-161.
(online at: https://www.iagi.or.id/web/digital/37/PIT-1983-Paper-15.pdf)
('Pliocene fossil vertebrates in the Banda Arc: structural implications'. On potential migration routes of mammals and hominids. No English abstract)

Sartono, S. & R. Marino (1978)- A mandibula and a maxilla of *Stegodon timorensis*. Modern Quaternary Research in Southeast Asia 4, Balkema, Rotterdam, p. 41-50.
(Additional Pleistocene elephant teeth from Weaiwe region, Atambua, NE part of West Timor)

Sartono, S., D.W. Orchiston, W.G. Siesser & T. Djubiantono (1981)- Upper Pliocene sediments in Sangiran, Central Java (Indonesia). Buletin Geologi (ITB) 5, p. 1-25.
(Base Sangiran Fm between nannofossil zones NN16 and NN18 (1.65- 3.25 Ma); Bettis et al. 2004)

Sartono, S., F. Semah, K.A.S. Astadiredja & T. Djubiantono (1981)- The age of *Homo modjokertensis*. Modern Quaternary Research in Southeast Asia 6, Balkema, Rotterdam, p. 91-102.
(On age of fossil child skull cap originally described by Von Koenigswald in 1936, from locality near Perning, 14 km ENE of Mojokerto. Supposedly from Lower Pleistocene U Pucangan Fm/ volcanic facies, but rocks more likely equivalent of M Pleistocene Kabuh Fm of Sangiran. Underlying clay facies of Lower Pucangan/ Lidah Fm with zone N21 planktonic foraminifera (Globorotalia tosaensis, Globigerinoides obliquus extremus; also Asterorotalia trispinosa). New finds of associated Hippopotamus, Sus, Buffelus bubalus, etc., suggest M-U Pleistocene Trinil or Ngandong fauna)

Sartono, S., F. Semah & T. Djubiantono (1984)- Paleomagnetic results from the Bringinan and Gemolong domes- correlations with the Sangiran section. Bull. Jurusan Teknik Geologi I.T.B. 13, p. 17-21.

Sartono S., D.E. Tyler & G.S. Krantz (1995)- A new *Homo erectus* skull from Sangiran, Java: an announcement. In: R.F. Bower & S. Sartono (eds.), Human evolution in its ecological context, 1: Evolution and ecology of *Homo erectus*, Pithecanthropus Centennial Foundation, Leiden, p. 207-210.
(Discovery in ~1993 of one of more complete Homo erectus skulls in Sangiran, C Java (also known as Skull IX or Tjg-1993-05))

Satyana, A.H. (2007)- Geological disaster in the demise of Jenggala and Majapahit empires: a hypothesis of historical mud volcanoes eruptions based on historical chronicles of Kitab Pararaton, etc. folklore of Timun Mas; analogue to present LUSI eruption, and geologic analysis of the Kendeng depression- Brantas Delta. Proc. Joint Convention 36th IAGI, 32nd HAGI, Bali 2007, p. 1-38.
(Jenggala and Majapahit are two empires of 11th to early 16th centuries at Brantas delta, E Java. Rise and fall related to geological processes in Brantas delta. Large mud volcano eruptions may have caused or contributed to demise)

Satyana, A.H. (2008)- Roles of mud volcanoes eruptions in the decline of the Jenggala and Majapahit Empires, East Java, Indonesia: constraints from the historical chronicles, folklore, and geological analysis of the Brantas Delta-Kendeng Depression. Majalah Geologi Indonesia (IAGI) 23, 1-2, p. 1-10.
(online at: https://www.academia.edu/12238239/ROLES_OF_MUD_VOLCANOES_ERUPTIONS_IN_THE_DECLINE_OF_THE_JENGGALA_AND_MAJAPAHIT_EMPIRES_EAST_JAVA_INDONESIA Etc.)
(Mud volcanoes in Kendeng zone may have lead to demise of 11th- 16th century Jenggala and Majapahit empires. (Also re-published in ISPG Memoir 1, p. 6-15 (2019))

Satyana, A.H. (2009)- Sangiran dome, Central Java: mud volcanoes eruption, demise of *Homo erectus erectus* and migration of later hominid. Proc. 37th Annual Conv. Indonesian Association Geologists (IAGI), Bandung 2008, p. 1-12.
(online at: https://www.iagi.or.id/web/digital/14/2008_IAGI_Bandung_Sangiran-Dome.pdf)
(Diapyric deformation and eruption of Sangiran Dome between 0.7- 0.5 Ma, possibly also at 0.12 Ma. Homo erectus fossils at Sangiran from 1.3- 0.7 Ma; its demise possibly triggered by mud volcano eruption)

Schurmann, H. (1928)- Kjokkenmoddinger en Paleolithicum in Noord Sumatra. De Mijningenieur 12, p. 1-19.
(Report on 1928 excavation of Early Holocene man-made shell midden Binjai Tamieng, along lower course of the Tamiang River in Aceh, NE Sumatra coastal plain, by BPM geologist H.M.E. Schurmann. Mound mainly composed of of shells of the brackish bivalve mollusc Meretrix meretrix. Also contain 'Hoabinhian' stone artifacts and 'kitchen debris from E Holocene humans, like vertebrate bones of marine turtles, fish, artiodactyls (deer, pig) and some crocodiles, rhinocerus and small whales. Schurmann collection of vertebrates deposited in Naturalis Museum, Leiden, and briefly examined by Hooijer (1948) and Gruwier (2017) (similar excavation conducted in same area by P.V. van Stein Callenfels in 1925) (see also Hooijer 1948, Gruwier(2017))

Schurmann, H.M.E. (1931)- Kjokkenmoddinger und Palaolithicum in Nord-Sumatra. Tijdschrift Koninklijk Nederlands Aardrijkskundig Genootschap Ser. 2, 48, p. 905-923.
(online at: https://resolver.kb.nl/resolve?urn=MMUBA13:001675001:pdf)

(Same as Schurmann (1928), above. On Paleolithic human-made shell mounds ('shell middens'; prehistoric 'kitchen debris'), containing Paleolithic stone tools in coastal NE Sumatra)

Schutt, G. (1972)- Fossil mammals of Java IV. Zur Kenntnis der pleistozanen Hyanen Javas. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam B 75, p. 261-287.
('On the knowledge of Pleistocene hyenas from Java')

Schutt, G. (1973)- Fossil mammals of Java V. Pleistozane Caniden (Carnivora, mammalia) aus Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 76, 5, p. 446-471.
('Pleistocene canids from Java')

Schwartz, J. (2016)- Beyond *Homo erectus*: Sangiran is key to deciphering the Asian human fossil record and re-evaluating *Homo*. In: F. Ribot Trafi (ed.) Homenaje al Dr. Jose Gibert Clols, una vida dedicada a la ciencia y al conocimiento de los primeros europeos. Publ. Diputacion de Granada, Granada, p. 93-110.
(online at: www.academia.edu/31517054/BEYOND_HOMO_ERECTUS_SANGIRAN_IS_KEY_TO_DE Etc.)
(Homo erectus widespread and highly variable species from which later hominids emerged. Review of relevant specimens from Sangiran, other Javanese sites and Zhoukoudian Lower Cave (China) shows concept 'erectus' masks evidence of taxic diversity in Asian hominid fossil record and raises questions about non-Asian specimens belonging to single species)

Selenka, M.L. & M. Blanckenhorn (eds.) (1911)- Die *Pithecanthropus*-Schichten auf Java. Geologische und palaontologische Ergebnisse der Trinil-Expedition (1907 und 1908). Verlag W. Engelmann, Leipzig, p. 1-342.
(online at: https://ia800303.us.archive.org/23/items/diepithecanthrop00sele/diepithecanthrop00sele.pdf)
(Extensive report of 1907-1908 excavations of Pleistocene beds near Trinil, C Java, by Selenka Trinil Expedition No new hominid fossils found, but extensive documention of localities and good descriptions of Pleistocene vertebrate and invertebrate faunas and flora)

Semah, A.M. (1982)- Etude palynologique de sites a hominides de l'ile de Java. These 3me Cycle, Universite de Provence, Marseille, p. 1-127.
('Palynological study of hominid sites of Java island')

Semah, A.M. (1982)- A preliminary report on a Sangiran pollen diagram. Modern Quaternary Research in Southeast Asia 7, Balkema, Rotterdam, p. 165-170.
(Summary of palynology work on U Pliocene- Pliocene of Sangiran Dome, C Java. Rel. common mangrove pollen (Rhizophora, Sonneratia, Nypa) in Upper Kalibeng Fm Blue Clays (absent in Pucangan Black Clay and in younger section). In Pucangan/ Kabuh interval mainly Graminae (grasses))

Semah, A.M. (1984)- Remarks on the pollen analysis of the Sambungmacan section (Central Java). Modern Quaternary Research in Southeast Asia 8, Balkema, Rotterdam, p. 29-34.
(Pollen diagram of Pleistocene deposits of Sambungmacan, E of Solo (site of 1973 discovery of Homo erectus skull). <10m thick poorly-dated fluvial, sandy-tuffaceous section, unconformable over Pliocene limestone)

Semah, A.M. (1984)- Palynology and Javanese *Pithecanthropus* environment. In: P. Andrews & J.L. Franzen (eds.) The early evolution of man with special emphasis on Southeast Asia and Africa, Courier Forschungsinstitut Senckenberg 69, p. 237-243.
(Vegetation changes in the Solo area, CJava, during Early-M Pleistocene. Littoral mangrove environment at end of Pliocene followed by swampy and more continental vegetation. Open character during Lower Pleistocene Pucangan Fm, resembling rainforest. Near transition to Kabuh Fm less humid vegetation and in Kabuh Fm mixed forest environment)

Semah, A.M. (1986)- Le milieu naturel lors du premier peuplement de Java, Resultats de l'analyse pollinique. Doct. Thesis, Universite de Provence, Marseille, 3 vols., p. 1-322.
('The natural environment of the first human seetlement of Java; results of pollen analysis')

Semah, A.M. & F. Detroit (2006)- Sur les premiers peuplements du Pacifique sud. Comptes Rendus Palevol 5, 1, p. 381-393
('About the first human settlements in the South Pacific'. First arrivals of Homo sapiens in Australia >40ka, possibly 50-60 ka. Debate whether these anatomically modern H. sapiens came from recent 'out of Africa' migration (Out-of-Africa hypothesis) or evolved locally from last Indonesian H. erectus (multiregional hypothesis). Morphometric differences between most recent Indonesian H. erectus and 'robust' Australian fossil H. sapiens from Kow Swamp and Cohuna clearly distinct, questioning local direct evolution)

Semah, A.M. & T. Djubiantono (2007)- Outline of climate and vegetation changes in Java during the Quaternary. In: A.M. Semah & K. Setiagama (eds.) First Islanders; human origins patrimony in Southeast Asia, p. 85-91.
(online at: http://hopsea.mnhn.fr/pc/brochures/2007HOPseaFI.pdf)
(Brief review of Pleistocene climate trends on Java. Between ~1 and 0.2 Ma climate rel. cooler and drier and yielding majority of hominid fossils)

Semah, A.M. & F. Semah (2001)- La signification paleoecologique des couches a hominides de l'ile de Java. In: F. Semah et al. (eds.) Origine des peuplements et chronologie des cultures paleolithiques dans le sud-est asiatique, Semenanjung/Artcom, Paris, p. 251-278.
('The paleoecological significance of the hominid beds of Java island')

Semah, A.M. & F. Semah (2012)- The rain forest in Java through the Quaternary and its relationships with humans (adaptation, exploitation and impact on the forest). Quaternary International 249, p. 120-128.
(online at: https://www.academia.edu/download/49792644/j.quaint.2011.06.01320161022-6152-vftwci.pdf)
(Landscape change in Java over last 2.5 million years highly complex, with repeated expansion and fragmentation of rain forest. Evidence of intensive human impact on rain forest observed late, ~1500 years ago)

Semah, A.M., F. Semah, T. Djubiantono & B. Brasseur (2010)- Landscapes and hominids' environments: changes between the Lower and the Early Middle Pleistocene in Java (Indonesia). Quaternary International 223, p. 451-455.
(Change in paleoenvironments in Lower and early M Pleistocene, based on sediment and pollen records in C Java. Late Lower Pleistocene 'Grenzbank' calcareous conglomerate layer at ~0.9 Ma at Sangiran marks tectonic event (folding of Kendeng zone). Second event is major climate change at E-M Pleistocene boundary, with severe fragmentation of rainforest cover during glacials (~0.8 Ma; close to in-situ 0.793 Ma Australasian tektites and Brunhes- Matuyama boundary in lowermost Kabuh Fm). Faunal turnover in Lower Kabuh Fm tied O-isotope zones MIS19-MIS22 interval. Little detail)

Semah, A.M., F. Semah, A.M. Moigne, T. Ingicco, A. Purnomo, T. Simanjuntak & H. Widianto (2016)- The palaeoenvironmental context of the Palaeolithic of Java: a brief review. In: A. Borel et al. (eds.) Southeast Asia: Human evolution, dispersals and adaptation, Quaternary International 416, p. 38-45.
(online at: https://os.pennds.org/archaeobib_filestore/pdf_articles/QI/2016_416_Semahetal.pdf)
(Earliest Paleolithic implements in Java Island >1 Ma old, postdating oldest Homo erectus fossils. Acheulean-like tools in early M Pleistocene ('Sangiran flakes'; 1.0-0.8 Ma), flake tools assemblages in late M/early U Pleistocene sites and development of cave occupations at end Pleistocene and E Holocene. Environment, mostly forested in E Pleistocene, changing climate during M Pleistocene, then at start of Late Pleistocene. Tectonic and volcanic activities affected local climate, paleogeography and floras. Associated vertebrate faunas reflect periods of contact with mainland (increased biodiversity) and periods of isolation (endemism))

Semah, A.M., F. Semah, R. Moudrikah, F. Frohlich & T. Djubiantono (2004)- A Late Pleistocene and Holocene sedimentary record in Central Java and it palaeoclimatic significance. In: S.G. Keates & J. Pasveer (eds.) Quaternary Research in Indonesia, Chapter 5, Modern Quaternary Research in Southeast Asia 18, Balkema, Leiden, p. 63-88.
(40m long cores from Ambarawa Basin (Solo Depression), C Java, provides sedimentary/ palynological record of last 21,000 years, from Last Glacial Maximum to present. Pollen from ~21-15 ka reflect cooler conditions

and severe dryness, with grassland and lowland forest. From ~15-10.5 ka increased precipitation with swamp conditions. From 10.5 ka onward more forest vegetation. First evidence of human activity at ~1500 yrs BP)

Semah, F. (1982)- Pliocene and Pleistocene geomagnetic reversals recorded in the Gemolong and Sangiran Domes (Central Java). Modern Quaternary Research in Southeast Asia 7, Balkema, Rotterdam, p. 151-164.
(Notopuro and Kabuh Fms of Sangiran and Gemolong all with normal geomagnetic polarity and tied to Brunhes Epoch. Reversal events in underlying Pucangan Fm, tied to Matuyama Event. Top of U Kalibeng Fm normal polarity, probably Olduvai Event (1.67-1.87Ma))

Semah, F. (1984)- The Sangiran Dome in the Javanese Plio-Pleistocene chronology. In: P. Andrews and J.L. Franzen (eds.) The early evolution of man with special emphasis on Southeast Asia and Africa. Courier Forschungsinstitut Senckenberg 69, p. 245-252.
(Review of Late Pliocene- M Pleistocene stratigraphy of Sangiran, C Java (~2.0- 0.5 Ma))

Semah, F. (1986)- Le peuplement ancien de Java: ebauche d'un cadre chronologique. L'Anthropologie 90, 3, p. 359-400.
(Chronology of human population of Java.Incl. radiometric and paleomagnetic dating of Kali Glagah and Kali Biuk site of oldest Java mammal assemblages (Bumiayu) as 2.2-1.67 Ma))

Semah, F. (1997)- Plio-Pleistocene reference sections in Indonesia. In: J.A. van Couvering (ed.) The Pleistocene boundary and the beginning of the Quaternary, World and Regional Geology 9, Cambridge University Press, p. 264-272.
(Review of Plio-Pleistocene stratigraphy in W and E Central Java areas known for vertebrate fossils (Bumiayu, Sangiran dome. Homo erectus mainly in lower and middle Kabuh Fm of Sangiran Dome. Australasian tektite event, radiometrically dated as 0.7- 0.8 Ma, in middle Kabuh Fm, close to Matuyama-Brunhes paleomagnetic boundary)

Semah, F. (2001)- La position stratigraphique du site de Ngebung 2 (dome de Sangiran, Java Central, Indonesie). In: F. Semah et al. (eds.) Origine des peuplements et chronologie des cultures paleolithiques dans le sud-est asiatique, Paris, p. 299-329.
(The stratigraphic position of Ngebung 2 excavation site, Sangiran Dome)

Semah, F. (2014)- Island Southeast Asia and human evolution heritage. In: N. Sanz (ed.) Human origin sites and the World Heritage convention in Asia, UNESCO World Heritage Papers 39, p. 184-210.
(online at: http://unesdoc.unesco.org/images/0022/002291/229174e.pdf)
(Review of 'journey' of Homo erectus and Homo sapiens across Indonesian archipelago since 1.5 Ma)

Semah, F., C. Falgueres, Y. Yokoyama, G. Feraud, H. Saleki & T. Djubiantono (1997)- Arrivee et disparition des *Homo erectus* a Java, les donnees actuelles. Abstracts 3rd Meeting European Association of Archaeologists, p. 11-12. *(Abstract only)*
('Arrival and migration of Homo erectus on Java, the actual data'. Critique of Swisher et al. (1994), suggesting their 1.66 Ma age assigned to H. erectus skulls from Sangiran is based on radiometric age of volcanic tuff that underlies these skulls and is therefore too old)

Semah, F., H. Saleki, C. Falgueres, G. Feraud & T. Djubiantono (2000)- Did early man reach Java during the Late Pliocene? J. Archaeological Science 27, 9, p. 763-769. *(online at: www.researchgate.net/publication/248578878_Did_Early_Man_reach_Java_during_the_Late_Pliocene)*
(Homo erectus (Pithecanthropus) reached Java from Asian continent and became one of oldest islanders in world. Combined 40Ar/39Ar and paleomagnetic data of 'Lower lahar' at base of fossil-bearing series of Sangiran dome show that emergence of first dry land at Sangiran, took place at end and just after Olduvai subchron. Therefore ~1.7 Ma is max. age for arrival of first hominids at Sangiran)

Semah, F. & A.M. Semah (2006)- Palaeolithic settlements in the Southeast Asian archipelagos: an Indonesian perspective. In: T. Simanjuntak et al. (eds.) Archaeology: Indonesian perspective, R.P. Soejono's Festschrift, LIPI Press, Jakarta, p. 148-161.

Semah, F. & A.M. Semah (2015)- Pleistocene migrations in the Southeast Asian archipelagoes. In: P. Bellwood (ed.) The global prehistory of human migration, Wiley-Blackwell, p. 49-54.
(Describes migrations of early hominins in SE Asia, especially to Java. Flores was also reached at least one million years ago. Early modern human migrations beyond Sunda Shelf also discussed)

Semah, F., A.M. Semah & T. Djubiantono (1998)- From the shoreline to the slopes of the volcanoes: the long *Pithecanthropus* trek. In: T. Simanjuntak et al. (eds.) Sangiran: man, culture, and environment in Pleistocene times, Proc. International Colloquium on Sangiran, Solo- Indonesia, Solo 1998, Jakarta, National Research Centre of Archaeology, Yayasan Obor Indonesia, Jakarta, p. 195-218.

Semah, F., A.M. Semah, T. Djubiantono & H.T. Simanjuntak (1992)- Did they also make stone tools? J. Human Evolution 23, p. 439-446.
(online at: https://www.academia.edu/4556315/Did_they_also_make_stone_tools? Etc.)
(Recent excavations in Ngebung, NW part of Sangiran dome, C Java, found archeological layers in M Pleistocene Kabuh beds with several stone tools (flake artefacts, bolas made of andesite and quartz; older than other stone implements known from Java))

Semah, F., A.M. Semah, C. Falgueres, F. Detroit, X. Gallet, S. Hameau, A.M. Moigne & T. Simanjuntak (2004)- The significance of the Punung karstic area (eastern Java) for the chronology of the Javanese Palaeolithic, with special reference to the Song Terus cave. Modern Quaternary Research in Southeast Asia 18, p. 45-62.
(online at: https://www.academia.edu/37335833/The_significance_of_the_Punung_karstic_area_eastern_Java_for_the_chronology_of_the_Javanese_Palaeolithic_with_special_reference_to_the_Song_Terus_cave)
(Caves in Punung and Wonosari areas in Southern Mountains (Gunung Sewu) of Central Java known for its many prehistoric workshops of Neolithic adzes from local silicified rocks, as well as more archaic Paleolithic artefacts in river beds. The karst deposits in caves existed at least since middle M Pleistocene. Human remains, including numerous stone artifacts, date back to ~200 ka)

Semah, F., A.M. Semah, S. Noerwidi, T. Ingicco, T. Simanjuntak & H. Widianto (2023)- Quelques perspectives concernant le peuplement et la Prehistoire ancienne de l'Indonesie dans le contexte de l'Asie du Sud-Est: des premiers hominides jusqu'a l'arrivee de notre espece. L'Anthropologie 127, 3 (Special Issue Prehistoire de l'Indonesie), 103147, p. 1-11.
('Some perspectives on human settlements and ancient prehistory of Indonesia in the context of Southeast Asia: from the first hominids to the arrival of our species'. Another modern review of the highlights of the early hominid story on Java)

Semah, F., A.M. Semah & T. Simanjuntak (2002)- More than a million years of human occupation in insular Southeast Asia. In: J. Mercader (ed.) Under the canopy- The archaeology of tropical rain forests, Rutgers University Press, New Brunswick, N.J., p. 161-190.

Semah, F., A.M. Semah, T. Simanjuntak & H. Widianto (2022)- Humans in Island Southeast Asia prior to *Homo sapiens* settlement, with special reference to Java Island. In: C.F.W. Higham & N.C. Kim (eds.) The Oxford Handbook of Early Southeast Asia, Chapter 1, p. 16-37.

Semah, F., A.M. Semah & H. Widianto H. (2014)- Insular Southeast Asia at the interface of continent-archipelago: geography and chronology. In: C. Smith (ed.) Encyclopedia of Global Archaeology, Springer, New York, p. 3890-3904. *(also in 2nd edition 2020, p. 5781-5795)*

Semah, F., T. Simanjuntak, E. Dizon, C. Gaillard & A.M. Semah (2014)- Insular Southeast Asia in the Lower Palaeolithic. In: C. Smith (ed.) Encyclopaedia of Global Archaeology, Springer, New York, p.

Setiyabudi, E. (2006)- Paleontological study on fossil giant tortoises from the Indonesian islands. M.Sc. Thesis Kagoshima University, Japan, p. 1-325. *(Unpublished)*

Setiyabudi, E. (2009)- Systematic study on the Early Pleistocene tortoises from the Indonesian islands and their paleozoogeographic implication. Dissertation Kagoshima University, Div. Nature System Science, p. 1-74.

Setiyabudi, E. (2009)- An early Pleistocene giant tortoise (Reptilia; Testudines; Testudinidae) from the Bumiayu area, Central Java, Indonesia. J. Fossil Research 42, 1, p. 1-11.
(online at: http://www.kasekiken.jp/kaishi/kaishi_42(1)/kasekiken_42(1)_1-11.pdf)
(Well-preserved ~1.75m long extinct testidinid tortoise from E Pleistocene lower Kali Glagah Fm, N of Bumiayu, originally studied by Van der Maarel (1932) and part of Java Satir Fauna of ~1.5Ma. Here identified as Megalochelys cf. sivalensis. Giant tortoises also known from Myanmar, Flores, etc.)

Setiyabudi, E. (2016)- Pleistocene reptiles of the Soa Basin (Flores, Indonesia): adaptation and implication for environment. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 17, 2, p. 107-124.
(online at: https://jgsm.geologi.esdm.go.id/index.php/JGSM/article/view/23/23)
(Presence of fossil reptiles known from mainland Asia in Pleistocene of Soa Basin, Flores: giant tortoise (Megalochelys sp.), fresh water turtle (Geoemydidae), crocodiles and komodo dragon (Varanus komodoensis). After crossing 'Wallace Line' at ~1 Ma lived in isolated conditions and adapted to savannah environment. Vertebrate faunas of Soe basin dominated by extinct proboscideans (Stegodon))

Setiyabudi, E. (2019)- Konservasi geologi lokasi fosil vertebrata dan manusia purba, daerah Tulungagung Selatan, Jawa Timur. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 20, 4, p. 187-197.
(online at: https://jgsm.geologi.esdm.go.id/index.php/JGSM/article/view/453/415)
(Geoconservation of vertebrate and human ancient fossils site, the South Tulungagung area, East Java'. Late Pleistocene- E Holocene vertebrate/hominid fossil sites associated with 'Wajak Man' of E. Dubois (1890), in marble mining area S of Tulungagung, deserve conservation protection status: Wajak Cave, Song Gentong, and Tenggar Cave)

Setiyabudi, E., I. Kurniawan & G.D. van den Bergh (2012)- Fossils of *Stegodon* and *Varanus komodoensis* Sumba and Flores: a Pleistocene landbridge? Proc. 41st Annual Conv. Indonesian Association Geologists (IAGI), Yogyakarta, 2012-SS-31, p. 618-621.
(online at: www.academia.edu/19327889/FOSSILS_OF_STEGODON_AND_VARANUS_KOMODO Etc.)
(Recent field survey in Sumba relocated original locality of Stegodon sumbaensis mandible described by Sartono (1979). At Lewapaku dwarf Stegodon found with tooth of Varanus komodoensis and giant murine rodent. Lewapaku fauna similar to 900 ka old Tangi Talo fauna from Flores. Similarities between Pleistocene fossils from Flores and Sumba make it tempting to assume land connections between two islands, but all taxa belong to groups that are excellent overseas island colonizers)

Setiyabudi, E., B. Prasthisto, I. Kurniawan & T. Jatmiko (2018)- The Early Holocene vertebrate faunas from Seropan Cave, Gunung Sewu, Yogyakarta, Indonesia. Indonesian J. on Geoscience (IJOG) 5, 1, p. 33-45.
(online at: https://ijog.geologi.esdm.go.id/index.php/IJOG/article/view/404/253)
(Open woodland mammal fauna from Seropan Cave in Gunung Sewu karst area, Wonosari, with Cervus, Sus verrucosus, Bubalus, Panthera cf. pardus, etc. C14 radiocarbon age dating gave date of 9,450 ± 400 yrs. BP)

Setiyabudi, E., A. Takahashi & Y. Kaifu (2016)- First certain fossil record of *Orlitia borneensis* (Testudines: Geoemydidae) from the Pleistocene of Central Java, Indonesia. Current Herpetology 35, 2, p. 75-82.
(online at: https://www.jstage.jst.go.jp/article/hsj/35/2/35_75/_article/-char/en)
(Plleistocene turtle fossil identified as Orlitia borneensis from Solo River bottom in Sambungmacan, eastern C Java, presumably eroded from M Pleistocene fluvial deposits on river bank. O. borneensis had wider distribution in past, but Java population would have become extinct by end of M Pleistocene)

Setyanta, B., H.P. Siagian & H. Wahyono (2014)- Penentuan umur fosil manusia purba di Jawa berdasarkan magnetostratigrafi. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 15, 1, p. 11-24.
(online at: http://jgsm.geologi.esdm.go.id/index.php/JGSM/article/view/67/69)
('Determination of the ages of the ancient hominid fossils in Java based on magnetostratigraphy'. Review of magnetostratigraphic studies of hominid sites of C Java: Sangiran (hominids between base Jamarillo subchron and Brunhes-Matuyama chron boundary in Upper Kabuh Fm (0.78- 1.07 Ma), Trinil (near top Jaramillo (0.99 Ma), Kedungbrubus, Mojokerto (near lower limit Jaramillo (1.07 Ma) and Patiayam (hominids in Brunhes or younger (<0.78 Ma))

Shipton, C., M.W. Morley, S. Kealy, K. Norman, C. Boulanger, S. Hawkins, M. Litster et al. (2024)- Abrupt onset of intensive human occupation 44,000 years ago on the threshold of Sahul. Nature Communications 15, 4193, p. 1-15.
(online at: https://www.nature.com/articles/s41467-024-48395-x)
(Laili rock shelter in Carboniferous crinoid limestone in northern Timor-Leste, wich is on southern dispersal route of Homo sapiens between Eurasia and Australasia and has earliest record of human occupation in E Wallacean archipelago. Abrupt onset of intensive human habitation in shelter started at ~44 ka (~MIS3). Initial occupation shown by burnt fish bones and damaged chert lithics. Intensive early occupation at Laili represents colonisation phase, which may have overwhelmed previous human dispersals in this part of world.)

Shipton, S. O'Connor & S. Kealy (2021)- The biogeographic threshold of Wallacea in human evolution. Quaternary International 574, p. 1-12.
(Wallacean archipelago (E Indonesia) is critical biogeographic boundary for all kinds of animals. Three stage model for how genus overcame this boundary: (1) Lower Paleolithic hominins (H. erectus) were able to colonize larger islands of W Wallacea through incidental seagoing and subsisting on megaherbivores; (2) arrival of early Homo sapiens around ~70 ka; and (3) during terminal Pleistocene and E Holocene greater connectivity in form obsidian exchange networks, inter-island bead and fish hook traditions, colonization of small islands without terrestrial fauna. Etc.)

Shutler, R. & F. Broches (1987)- The paleoanthropology of Pleistocene island Southeast Asia: a review. In: N. Thiramongkol (ed.) Proc. Workshop on Economic geology, tectonics, sedimentary processes and environment of the Quaternary in Southeast Asia, Haid Yai, Thailand 1986, IGCP 218/ Chulalongkorn University, Bangkok, p. 185-191.
(online at: http://library.dmr.go.th/Document/Proceedings-Yearbooks/M_1/1986/5083...)
(Brief review of Pleistocene Java mammal assemblages and issues regarding dating and interpretation)

Shutler, R., J.M. Head, D.J. Donahue, A.J.T. Jull, M.F. Barbetti, S. Matsu'ura, J. de Vos & P.Storm (2004)- AMS radiocarbon dates on bone from cave sites in Southeast Java, Indonesia, including Wajak. Modern Quaternary Research in Southeast Asia 18, p. 89-93.
(14C apatite dates for bones from Wajak Cave, SE Java. Wajak fauna calibrated age ~12.1- 12.9 ka, Wajak femur ~7.5 ka (older ages obtained by Storm et al. 2013))

Siesser, W.G. & D.W. Orchiston (1978)- Micropalaeontological re-assessment of the age of *Pithecanthropus* mandible C from Sangiran, Indonesia. Modern Quaternary Research in Southeast Asia 4, p. 25-30.
(Pithecanthropus mandible C discovered in 1960 in Pucangan Fm near Mandingan, Sangiran Dome. Sartono (1970) reported 11 planktonic and 15 benthic species of foraminifera in claystone adhered to mandible. Co-occurrence of Globorotalia crassaformis and Globigerinoides obliquus suggests Late Pliocene age, between 4.2-1.6 Ma (N.B.: Sangiran Pleistocene fluvial-lacustrine Pucangan Fm contains locally abundant Pliocene planktonic foraminifera, but all reworked from underlying Kalibeng Fm; Van Gorsel and Troelstra 1981))

Sighinolfi, G.P., S. Sartono & G. Artioli (1993)- Chemical and mineralogical studies on hominid remains from Sangiran, Central Java (Indonesia). J. Human Evolution 24, p. 57-68.

Simanjuntak, T. (1995)- Mesolithique de l'Indonesie: une heterogeneite culturelle. L'Anthropologie 99, 4, p. 626-636.

('The Mesolithic of Indonesia: a cultural heterogeneity')

Simanjuntak, T. (2001)- New light on the prehistory of the Southern Mountains of Java. Bull. Indo-Pacific Prehistory Association (IPPA) 21, p. 152-156.
(online at: http://ejournal.anu.edu.au/index.php/bippa/article/view/272/262)
(Many limestone caves in Gunung Sewu, S Java inhabited in Prehistoric times. Excavations revealed chronology from Early Holocene (possibly Late Pleistocene) to 4500 BP. by 'Australomelanesian' people)

Simanjuntak T. (2001)- New insights on the tools of *Pithecanthropus*. In: T. Simanjuntak et al. (eds.) Sangiran: man, culture, and environment in Pleistocene times, Proc. International Colloquium on Sangiran, Solo-Indonesia, Solo 1998, Jakarta, National Research Centre of Archaeology, Yayasan Obor Indonesia, Jakarta, p. 154-170.

Simanjuntak, T. (2002)- Gunung Sewu in prehistoric times. Gadjah Mada University Press, Yogyakarta p.

Simanjuntak, T. (2004)- New insight on the prehistoric chronology of Gunung Sewu, Java, Indonesia. In: S.G. Keates & J. Pasveer (eds.) Quaternary Research in Indonesia, Chapter 2, Modern Quaternary Research in Southeast Asia 18, Balkema, Leiden, p. 9-30.
(C Java Southern Mountains numerous Late Pleistocene- Holocene, Paleolithic- Neolithic prehistoric sites)

Simanjuntak, T. (2006)- Indonesia-Southeast Asia: climates, settlements, and cultures in Late Pleistocene. Comptes Rendus Palevol 5, p. 371-379.
(Late Pleistocene period between Paleolithic culture and E Holocene Preneolithic culture, marked by climate and sea level fluctuations and appearance of modern human (oldest Homo sapiens), replacing H. erectus)

Simanjuntak, T. (2017)- The western route migration: a second probable Neolithic diffusion to Indonesia. In: P.J. Piper et al. (eds.) New perspectives in Southeast Asian and Pacific prehistory, Terra Australis 45, ANU Press, p. 201-211.
(online at: https://www.jstor.org/stable/j.ctt1pwtd26)
(Neolithic in Indonesia generally traced to culture of Austronesian-speaking people who migrated from Taiwan at ~4000 BP. Another earlier probable Neolithic diffusion from mainland SE Asia, probably by Austroasiatic-speaking people. Subsequent dispersal of Austronesian-speaking people into W parts of Indonesia influenced Neolithic cultures there and resulted in replacement of local Austroasiatic languages)

Simanjuntak, T. (2023)- La prehistoire d'Indonesie: une breve synthese. L'Anthropologie 127, 3 (Special Issue Prehistoire de l'Indonesie), 103148, p. 1-19.
('The Prehistory of Indonesia, a brief synthesis'. Prehistory (starting 1.5 Ma with arrival of Homo erectus; seven stages), Protohistoric (cultures linked to metal and megaliths), Historical Period (introduction of Hindu-Buddhist influence in 4th-5th centuries A.D.)

Simanjuntak, T. & I.N. Asikin (2004)- Early Holocene human settlement in Eastern Java. Bull. Indo-Pacific Prehistory Association (IPPA) 24 (Taipei Papers, volume 2), p. 13-19.
(online at: https://journals.lib.washington.edu/index.php/BIPPA/article/view/11867/10494)
(Finds of modern human remains dated back to 45 ka (Late Pleistocene). E Holocene human settlements in E Java called Keplek Phase. Characteristics: exploitations of caves and rock shelters (Gunung Sewu, Sampung, Tulung Agung, Dander, Tuban. Puger, Situbondo), lithic flake industry, shell tool technology, etc. First excavations of E Java caves at Gua Lawah by Van Es (1926) and P. van Stein Callenfels (1928-1931). Etc.)

Simanjuntak, T., B. Prasetyo & R. Handini (eds.) (2001)- Sangiran: man, culture and environment in prehistoric times, Yayasan Obor Indonesia, Jakarta, p. 1-443.
(Collection of papers on Sangiran hominid site, C Java, presented at 1st Int. Colloquium on Sangiran, Solo 1998)

Simanjuntak, T. & F. Semah (1996)- A new insight into the Sangiran flake industry. In: The Chiang Mai papers, 1, Bull. Indo-Pacific Prehistory Association (IPPA) 14, p. 22-26.
(online at: http://journals.lib.washington.edu/index.php/BIPPA/article/view/11584/10215)
(Flake tools made of chalcedony, jasper, etc., and supposedly made by Homo erectus, first discovered by Von Koenigswald in 1936. Flakes can be found throughout Kabuh Fm at Ngebung, NW Sangiran Dome, C. Java)

Simanjuntak, T. & F. Semah (2006)- Indonesia-Southeast Asia: Climates, settlements, and culture in Late Pleistocene. Comptes Rendus Palevol 5, 1-2, p. 371-379.
(online at: www.sciencedirect.com/science/article/pii/S1631068305001314)
(Late Pleistocene period between Paleolithic culture, and Early Holocene Pre-neolithic culture (~45ka-11.8 ka). Climate and sea level fluctuations and appearance of oldest Homo sapiens, replacing Homo erectus)

Simanjuntak, T., F. Semah & C. Gaillard (2010)- The Palaeolithic in Indonesia: nature and chronology. Quaternary International 223-224, p. 418-421.
(Brief review of Indonesian 'older' Paleolithic stone tool assemblages. Human presence on Java dates back to ~1.5 Ma, but no stone tools known older than ~1.0 Ma)

Simanjuntak, T., F. Semah & A.M. Semah (2014)- Tracking evidence for modern human behavior in Paleolithic Indonesia. In: Y. Kaifu et al. (eds.) Emergence and diversity of modern human behavior in Paleolithic Asia, Peopling of the Americas Publications, Texas A&M University Press, p. 158-170.
(Homo sapiens may have colonized Indonesian Archipelago in early U Pleistocene. Most reliable evidence only since ~45 ka (Paleolithic sites with artifacts in Gunung Sewu, S Java))

Simanjuntak, T., F. Semah, A.M. Semah & H. Widianto (2023)- La prehistoire d'Indonesie: une breve synthese. L'Anthropologie 127, 3, 103148, p.
('The Prehistory of Indonesia, a brief synthesis'. Prehistory of Indonesia covers 1.5 million years, punctuated by major changes in evolution, adaptation and culture of human groups. Seven major stages: (1) arrival of Homo erectus; (2) spreading of the Acheulean tradition; (3) arrival of Homo sapiens; (4)) Pleistocene-Holocene transition; (5) arrival of Austro-asiatic populations; (6) arrival of Austronesiens, and (7) influence of megalithic traditions with use of metal, a few centuries before our era. Historical period starts with introduction of Hindu-Buddhist influence at ~4th/5th centuries AD.)

Sipola, M.E. (2018)- Formation of the Ngandong paleoanthropological site and Solo River terrace sequence, Central Java, Indonesia. Ph.D. Thesis University of Iowa, p. 1-246.
(online at: https://ir.uiowa.edu/etd/6286)
(Paleoanthropological excavations in Late Pleistocene fluvial terrace along Solo River at Ngandong uncovered 14 Homo erectus fossils. Preliminary Red Thermo Luminescence analyses of Ngandong terrace deposits did not provide reliable ages. Position at low terrace level and U-series dating suggest these may be the most recent known H. erectus in world (?; Oppenoorth (1932) and many late authors view Ngandong 'Solo Man' as separate species, H. soloensis, and relationship to H. erectus s.s. remains to be determined; JTvG))

Siswanto & S. Noerwidi (2014)- Fosil Proboscidea dari situs Semedo: hubungannya dengan biostratigrafi dan kehadiran manusia di Jawa. Jurnal Berkala Arkeologi 34, 2, p. 115-130.
(online at: http://balaiarkeologiyogya.com/berkalaarkeologi/article/view/20/37)
('Proboscidea fossils from Semedo site: its correlation with biostratigraphy and human arrival in Java'. Semedo site in Tegal district of C Java rich in vertebrate fossils, with high percentage and several species of Proboscidea (elephantoids): Sinomastodon bumiayuensis, Stegodon trigonocephalus, Stegodon 'pygmy' semedoensis, Stegodon hypsilophus, Elephas (Archidiskodon) planifrons and Elephas hysudrindicus)

Siswanto, S. & S. Noerwidi (2017)- Posisi fauna situs Patiayam dalam biostratigrafi Jawa. Berkala Arkeologi Sangkhakala 19, 2, p. 149-166.
(online at: https://sangkhakala.kemdikbud.go.id/index.php/SBA/article/view/31/pdf_1)
('The faunal position of Patiayam site in the biostratigraphy of Java'. Patiayam fauna fits between Cisaat group and Kedungbrubus group, recording environmental change and faunal inhabitants from 1.2-0.8 Ma)

Skeels, A., L.M. Boschman, I.R. McFadden, E.M. Joyce, O. Hagen, O. Jimenez Robles, W. Bach et al. (2023)- Paleoenvironments shaped the exchange of terrestrial vertebrates across Wallace's Line. Science 381, 6653, p. 86-92.
(online at: https://dspace.library.uu.nl/bitstream/handle/1874/429755/science.adf7122.pdf)
(Precipitation tolerance and dispersal ability were key for faunal exchange across the precipitation gradient spanning Indo-Australian region. 'Sundanian' (SE Asian) taxa evolved in humid climate similar to "stepping stones" of Wallacea, facilitating colonization of 'Sahulian' (Australian) continental shelf. Sahulian taxa evolved in more arid conditions, hampering migration into humid Sunda, causing asymmetrical migration)

Smith, H.E., G.J. Price, M. Duval, K. Westaway, J. Zaim, Y. Rizal, Aswan et al. (2021)- Taxonomy, taphonomy and chronology of the Pleistocene faunal assemblage at Ngalau Gupin cave, Sumatra. Quaternary International 603, p. 40-63.
(online at: www.sciencedirect.com/journal/quaternary-international/vol/603/suppl/C)
(Ngalau Gupin caves in Padang Highlands ~90 km from Sawahlunto rich in fossils, mainly isolated animal teeth. Diverse Pleistocene faunal assemblage (Proboscidea, Primates, Rodentia, Artiodactyla, Perissodactyla, Carnivora). Analogous to extant fauna in modern rainforests of Sumatra. Assemblage dated at ~160- ~115 ka)

Smith, T. M., A. Houssaye, O. Kullmer, A. Le Cabec, A.J. Olejniczak, F. Schrenk, J. de Vos & P. Tafforeau (2018)- Disentangling isolated dental remains of Asian Pleistocene hominins and pongines. PLoS One 13, 11, e0204737, p. 1-23.
(online at: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0204737)
(Examination of enamel thickness, enamel-dentine junction shape, and crown development in 33 molars from Von Koenigswald's Chinese Apothecary collection in Frankfurt (11 Sinanthropus officinalis [= Homo erectus], 21 Hemanthropus peii, and one Hemanthropus peii or Pongo) and 7 molars from Sangiran dome, Java (Homo erectus or Pongo). Homo and Pongo molars overlap in enamel thickness, etc. Need for additional studies)

Smith, T.M., A.J. Olejniczac, K.Kupczik, V. Lazzari, J. de Vos, O. Kullmer, F. Schrenk et al. (2009)- Taxonomic assessment of the Trinil molars using non-destructive 3D structural and development analysis. PaleoAnthropology: p. 117-129.
(online at: https://projects.iq.harvard.edu/files/tanyasmith/files/15.pdf)
(Two molars recovered by E. Dubois in 1891-1892 at Trinil, C Java, have been debate and attributed to several ape (Pongo) and human taxa. X-ray imaging suggests the molars are indeed Homo erectus teeth)

Soares, P., J.A. Trejaut, J.H. Loo, C. Hill, M. Mormina et al. (2008)- Climate change and postglacial human dispersals in Southeast Asia. Molecular Biology and Evolution 25, 6, p. 1209-1218.
(online at: https://academic.oup.com/mbe/article/25/6/1209/1134230)
(Modern humans in Island SE Asia since at least 50,000 years, commonly thought to be from Neolithic dispersal from China. Genome sequencing of modern humans suggest migration of humans from Sundaland across region since start of Holocene, at time of Sundaland breaking up into archipelago by rising sea levels)

Soejono, R.P. (1961)- Kebudatjaan Patjitan. Madjelis Ilmu Pengetahuan Indonesia, Penerbitan 2, p. 243-241.
('The Patjitan Culture')

Soejono, R.P. (1961)- Preliminary notes on new finds of Lower-Palaeolithic implements from Indonesia. Asian Perspectives 5, 2, p. 217-232.
(Review of Lower Paleolithic stone tools from Java, Sumatra, Kalimantan)

Soejono, R.P. (1969)- The history of prehistoric research in Indonesia to 1950. Asian Perspectives 12, p. 69-91.
(online at: https://scholarspace.manoa.hawaii.edu/server/api/core/bitstreams/8c05c190-65fc-4fd9-89a0-bcb9a0863292/content)
(Review of archeological work in the Indonesian region before 1950, incl. Paleolithic-Neolithic stone tools, bronze drums and other object, megalithic remains, fossilized human remains)

Soejono, R.P. (1970)- The study of Prehistory in Indonesia: retrospect and prospect. Asian Perspectives 13, p. 11-15.
(Archeological research in Indonesia in 1800s by naturalists, missionaries, military officers, civil servants, painters and photographers. Among founders of Indonesian archaeology are T.S. Raffles, R.H.T. Friederich and J. W. Ijzerman. Early work focused on Hindu-Javanese history. More systematic prehistoric research by Archaeological Service after 1921, with researchers P. V. van Stein Callenfels, R. von Heine-Geldern and A. N. J. Thomassen a Thuessink van der Hoop. After WW II, H.R. van Heekeren became prehistorian at Archaeological Service in Jakarta and made important contributions on Palaeolithic, Mesolithilithic and Bronze ages. Etc.)

Soejono, R.P. (1982)- New data on the Palaeolithic industry in Indonesia. In: M.A. de Lumley (ed.) Colloque Int. CNRS L'*Homo erectus* et la place de l'Homme de Tautavel parmi les hominides fossiles, Nice p. 578-592.

Soeradi, T., T. Shibasaki, D. Kadar, Sudijono, M. Itihari, H. Kumai et al. (1985)- Geology and stratigraphy of the Trinil Area. In: N. Watanabe & D. Kadar (eds.) Quaternary geology of the hominid fossil bearing formations in Java. Geological Research Development Centre (GRDC), Bandung, Special Publ. 4, p. 49-53.
(Re-examination of Dubois and Selenka mammal locality at Trinil, by an Indonesian-Japanese team in 1976-1977. The main bone bed with Dubois' Pithecanthropus erectus at the base of the 45-53m thick Kabuh Fm at Trinil (Hauptknochenschicht') is unconformably over the Pucangan Fm, and is remarakably similar to Pithecanthropus-bearing 'Grenzbank' in Sangiran)

Soerastopo Hadisoemarno (1972)- Geomorphology of the Sangiran dome, Java, Indonesia. Ilmu Alam, 1, p. 57-65. *(in Indonesian)*

Soergel, W. (1914)- Stegodonten aus den Kendengschichten auf Java. Palaeontographica, Supplement 4, 3, 1, p. 1-24.
(online at: http://sammlungen.ub.uni-frankfurt.de/botanik/periodical/pageview/4499569)
('Stegodont elephants from the Kendeng beds on Java'. Descriptions of Pleistocene Stegodont elephant teeth collected by J. Elbert, described by Prof. Wolfgang Soergel of the University of Freiburg)

Sondaar, P.Y. (1981)- The *Geochelone* faunas of the Indonesian Archipelago and their paleogeographical and biostratigraphical significance. Modern Quaternary Research in Southeast Asia 6, Balkema, Rotterdam, p. 111-120.
(E Pleistocene giant tortoise Geochelone atlas known from Java (Sangiran, Kali Klagah), Sulawesi and Timor. Probably part of island faunas, which became extinct with arrival of Asiatic mammals via land bridge in E Pleistocene. (see also critical discussion by Hooijer 1982))

Sondaar, P.Y. (1984)- Faunal evolution and the mammalian biostratigraphy of Java. In: P. Andrews & J.L. Franzen (eds.) The early evolution of man with special emphasis on Southeast Asia and Africa. Courier Forschungsinstitut Senckenberg 69, p. 219-235.
(Re-interpretation of fossil mammal successions on Java, linked to changes in paleobiogeography. Seven Late Pliocene- Pleistocene vertebrate faunas, from old to young: Satir (early island fauna), Ci Saat, Trinil HK (with arrival of Homo erectus), Kedung Brubus, Ngandong, Punung and Wajak faunas)

Sondaar, P.Y. (1987)- Pleistocene man and extinctions of islands endemics. Memoires Societe Geologique France, N.S., 150, p. 159-165.

Sondaar, P.Y. (1989)- Did man reach Australia via the giant rat and Dingo route? Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 5, p. 76-83.
(Discussion of possible routes followed by Pleistocene man from China to Australia)

Sondaar, P. (1993)- De *Stegodon* van Flores. Cranium 10, 1, p. 47-48.
(online at: http://natuurtijdschriften.nl/download?type=document&docid=523307)

(Elephantoid Stegodon rel. common in Plio-Pleistocene of SE Asia. Dwarf species known from Sulawesi, Flores, Sumba and Timor (many of these first found by missionary Verhoeven))

Sondaar, P. (1994)- De *Homo erectus* fauna's van Java. Cranium 11, 2, p. 92-96.
(online at: http://natuurtijdschriften.nl/download?type=document&docid=523346)
('The Homo erectus faunas of Java'. Changes in faunal succession in Pleistocene of Java explained by: (1) geographical position of Java at periphery of SE Asia; 2) succession of glacials, causing sealevel lowering, connecting Java, Borneo and Sumatra with continent; 3) reduction of tropical rainforest during glacials (dry periods, favoring migration of mammals living in open habitats). Two major turnovers in faunal succession of Java: (1) ~1.2 Ma change from unbalanced island fauna (Satir fauna) to continental fauna (Ci-Saat fauna; with Homo erectus and disappearance of endemic island forms); (2) Late Pleistocene (~80.000 yrs ago) extinction of M Pleistocene forms (Stegodon, Hexaprotodon, Homo erectus), replaced with new fauna, with extant Elephas maximus and probably Homo sapiens)

Sondaar, P.Y., F. Aziz, G.D. van den Bergh & J. de Vos (1996)- Faunal change and hominid evolution during Quaternary of Jawa. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 8, p. 1-10.
(Quaternary vertebrate assemblages, from old to young: (1) Satir (lower Kali Glagah, Bumiayu; unbalanced island fauna; ~1.5 Ma), (2) Cisaat (upper Kali Glagah, Bumiayu;still unbalanced fauna, ~1.2 Ma), (3) Trinil H.K ('Hauptknochenschicht'; with Homo erectus, ~1.0 Ma), (4) Kedung Brubus (upper Bapanmin Sangiran), (5) Ngandong (Solo River terrace deposits), (6) Punung (Late Pleistocene (humid tropical rainforest assemblage; last interglacial?, fissure deposits with Homo sapiens) and (7) Wajak (Wajak cave in E Java; Holocene, with Homo sapiens). Homo erectus in several different stratigraphic levels)

Sondaar, P.Y., J. de Vos & J.J.M. Leinders (1983)- Reply. Facts and fiction around the fossil mammals of Java. Geologie en Mijnbouw 62, p. 339-343.
(Discussion of Bartstra (1983) critique of De Vos, Sartono et al. (1982) reinterpretation of relative ages of mammalian faunas of Trinil and Kedungbrubus of Java. See also Hooijer 1983, Hooijer and Kurten 1984)

Sondaar, P.Y., G.D. van den Bergh, J. de Vos & F. Aziz (1995)- *Homo erectus* in S.E. Asia: time space and migration routes, IV. Overseas traveling of *Homo erectu*s and faunal turnovers. In: J. Gibert et al. (eds.) Proc. International Conference The hominids and their environment during the Lower and Middle Pleistocene of Eurasia, Orce 1995, p. 383-388.
(Homo sapiens generally believed to be first human to cross water barriers (since ~50 ka). Stone tools of 0.7 Ma in age found on Flores suggest the much older Homo erectus probably also had this capability)

Sondaar, P.Y., G.D. van den Bergh, J. de Vos & F. Aziz (2001)- The Flores case: the earliest island colonizers. Geological Research Development Centre (GRDC), Bandung, Special Publ. 27, p. 15-19.

Sondaar, P.Y., G.D. van den Bergh, B. Mubroto, F. Aziz, J. de Vos & U.L. Batu (1994)- Middle Pleistocene faunal turnover and colonization of Flores (Indonesia) by *Homo erectus*. Comptes Rendus Academie Sciences, Paris, II, 319, p. 1255-1262.
(Stone artifacts made of basalt, associated with fossil Stegodon trigonocephalus florensis Hooijer and fresh water molluscs in fluvial sands and tuffs of Ola Bula Fm near Mata Menge, W Central Flores. Tools probably made by Homo erectus and dated as slightly older than 0.73 Ma (based on possibly erroneous Matuyama-Brunhes paleomag interpretation and actually closer to 0.8-0.9 Ma: Morwood et al. 1998). Older endemic island fauna at base Ola Bula Fm at Tangi Talo, with pygmy Stegodon, Geochelone (large tortoise) and Varanus komodoensis)

Spriggs, M., C. Reepmeyer, Anggraeni, P. Lape, L. Neri, W.P. Ronquillo, T. Simanjuntak, G. Summerhayes et al. (2011)- Obsidian sources and distribution systems in Island Southeast Asia: a review of previous research. J. Archaeological Science 38, p. 2873-2881.
(online at: https://faculty.washington.edu/plape/pubs/JAS%202011.pdf)
(Review of distribution and origin of stone age obsidian artifacts in Philippines, Sulawesi, Flores, W and E Java, S Sumatra, Borneo and E Timor. Many probably sourced from islands on which they were found)

Stehlin, H.G (1925)- Fossile Saugetiere aus der Gegend von Limbangan (Java). Dienst Mijnbouw Nederlandsch-Indie, Wetenschappelijke Mededeelingen 3, p. 1-12.
('Fossil mammals from the Limbangan area, Java'. Diverse Pleistocene mammal assemblage in conglomeratic sandstone near Limbangan on Pamali River, Brebes District, C Java (16 km W of Tegal). Incl. Stegodon, Elephas, cervids. By Swiss vertebrate paleontologist Hans Georg Stehlin)

Storm, P. (1992)- Two microliths from Javanese Wadjak Man. J. Anthropological Society of Nippon 100, 2, p. 191-203.
(online at: https://www.jstage.jst.go.jp/article/ase1911/100/2/100_2_191/_pdf)
(Two hominid stone tools made from limestone, from rock shelter known as Wajak site on mountain slope in S Java, S of Mt Willis near village of Wajak, site of human skull first found by Van Rietschoten in 1888)

Storm, P. (1994)- De morfologie van *Homo modjokertensis*. Cranium 11, 2, p. 97-102.
(online at: http://natuurtijdschriften.nl/download?type=document&docid=523347)
(Mojokerto skull from E Java, described by Von Koenigswald (1936) as Homo modjokertensis, is juvenile skull, but not possible to determine if early (Homo erectus erectus) or late (Homo erectus soloensis) Javanese form (Dubois 1940 considered this to be rel. young Homo wadjakensis= H. soloensis))

Storm, P. (1995)- The evolutionary significance of the Wajak skulls. Scripta Geologica 110, p. 1-247.
(online at: https://repository.naturalis.nl/pub/317406)
Two Late Pleistocene- Holocene robust hominid skulls from fissure fill breccia in limestone of the Southern Mountains, Gua Wajak, SE Java, first described by Dubois (1922). Most likely interpretation as Mesolithic robust representatives of present Homo sapiens of Java)

Storm, P. (1997)- Fossielen uit de hete hel van Java. Cranium 14, 2, p. 91-98.
(online at: https://natuurtijdschriften.nl/pub/523415/CRAN1997014002006.pdf)
('Fossils from the hot hell of Java'. Report on a 1996 trip to hominid fossil sites of Java)

Storm, P. (2000)- The evolutionary history of humans in Australasia from an environmental perspective. Anthropological Science 108, 3, p. 225-260.
(online at: https://www.jstage.jst.go.jp/article/ase1993/108/3/108_3_225/_pdf)

Storm, P. (2001)- The evolution of humans in Australasia from an environmental perspective. In: R.A.C. Dam & S. van der Kaars (eds.) Quaternary environmental change in the Indonesian region, Palaeogeogr. Palaeoclim. Palaeoecology 171, p. 363-383.
(Incl. climatic sequence for Java: before and around 135 ka climate considerably drier and hot; between 126-81 ka mainly humid-warm (with Punung fauna), becoming drier and cooler (with Wajak fauna) before returning to more interglacial conditions in Holocene)

Storm, P. (2006)- De "Hobbit" van Flores: een mensachtige eilandvorm. Cranium 23, 2, p. 3-14.
(online at: http://natuurtijdschriften.nl/download?type=document&docid=523527)
(Prehistoric "hobbit" of Flores with small size (1m tall) and endocranial volume, cannot be compared with anything else in human family tree. Most likely new species, Homo floresiensis, that fits with island evolution)

Storm, P. (2012)- A carnivorous niche for Java Man? A preliminary consideration of the abundance of fossils in Middle Pleistocene Java. Comptes Rendus Palevol 11, 2-3, p. 191-202.
(online at: https://www.sciencedirect.com/journal/comptes-rendus-palevol/vol/11/issue/2)
(Anatomical and archeological aspects of Homo erectus sites Kedung Brubus and Trinil suggest vertebrates meat was important part of hominid diet)

Storm, P., F. Aziz, J. de Vos, D. Kosasih, S. Baskoro, Ngaliman & L.W. van den Hoek Ostende (2005)- Late Pleistocene *Homo sapiens* in a tropical rainforest fauna in East Java. J. Human Evolution 49, 4, p. 536-545.

(online at: www.academia.edu/ 18341302/Late_Pleistocene_Homo_sapiens_in_a_tropical_rainforest_ fauna_in_East_Java)
(Redescription of two original sites of 'Punung fauna', as first described by Von Koenigswald (1939) and Badoux (1959) from karst hills in Southern Mountains of E Java. Punung and new nearby mammal site Gunung Dawung reflect tropical rainforest environment with common Pongo (orangutan) fossils. Punung fauna younger than Ngandong, possibly around 100 ka)

Storm, P. & J. de Vos (2006)- Rediscovery of the Late Pleistocene Punung hominin sites and the rediscovery of a new site Gunung Dawung in East Java. Senckenbergiana Lethaea 86, 2, p. 121-131.
(On re-location of Von Koenigswalds's Pleistocene Punung sites where in 1930s he collected hominin remains and mammals indicative of tropical rainforest like orang-utans (Pongo) and gibbons (Hylobates))

Storm, P. & A.J. Nelson (1992)- The many faces of Wajak Man. Archaeology in Oceania 27, p. 37-46.
(online at: www.oermens.nl/many_faces_wadjak.pdf)
('Wadjak Man' first fossil hominid found in SE Asia (S Java in 1888 by Dubois). Initially believed to be ancestral to Australian Aborigines, but material sub-recent and first occupation of Australia well into Pleistocene (50ka?))

Storm, P., R. Wood, C. Stringer, A. Bartsiokas, J. de Vos, M. Aubert, L. Kinsley & R. Grun (2013)- U-series and radiocarbon analyses of human and faunal remains from Wajak, Indonesia. J. Human Evolution 64, p. 356-365.
(Radiometric dating of human and faunal bone fragments from Wajak, Java, indicate minimum age of 37.4-28.5 ka, older than previously published radiocarbon estimates, probably due to secondary carbonatisation. Requires reassessment of evolutionary relationships of human remains in SE Asia- Oceania)

Stremme, H. (1911)- Die Saugetiere mit Ausnahme der Proboscidier. In: M.L. Selenka & M. Blanckenhorn (eds.) Die Pithecanthropus-Schichten auf Java, geologische und palontologische Ergebnisse der Trinil Expedition (1907 und 1908), W. Engelmann, Leipzig, p. 82-150.
('The mammals, not including elephants', from Pleistocene of Trinil, from material collected by Selenka expedition, 1907-1908. Descriptions of rodents, Rhinoceros, Sus brachygnathus, Hippopotamus, Cervus (Axis) lydekkeri, Duboisia, Buffelus, Bibos spp., etc.))

Sudijono, K. Mano & R. Wikarno (eds.) (1995)- Geology of the Quaternary environment of the Solo-Madiun area, Central-East Java. Geological Research Development Centre (GRDC), Bandung, Special Publ. 17, p. 1-128.

Suharyogi, I.Y.P., U.P. Wibowo, H. Insani & E. Setiyabudi (2019)- *Duboisia santeng* (Bovidae, Artiodactyla) dari Bumiayu. Bull. Scientific Contribution (UNPAD): Geology 17, 1, p. 1-7.
(online at: http://jurnal.unpad.ac.id/bsc/article/view/20639/pdf)
('Duboisia santeng from Bumiayu', Java. Left bovidae horncore found in 2016 at Santanaya River, Cisaat (Bumiayu). Assigned to Duboisia santeng (Dubois, 1891). Probably from M Pleistocene Gintung Fm, possibly Kedung Brubus Fauna. Environment open woodland)

Suminto, J. Kimura & T. Hirayama (1995)- Subsurface geology in Sangiran, Kedungbrubus and Ponorogo area. In: Sudijono et al. (eds.) Geology of Quaternary environment of the Solo-Madiun area, Central-East Java, Geological Research Development Centre (GRDC), Bandung, Special Publ. 17, p. 81-86.

Suminto, M.J. Morwood, I. Kurniawan, F. Aziz, G.D. van den Bergh & D.R. Hobbs (2009)- Geology and fossil sites of the Soa Basin, Flores, Indonesia. In: F. Aziz et al. (eds.) Pleistocene geology, palaeontology and archaeology of the Pleistocene Soa Basin, Central Flores, Indonesia, Chapter 2, Pusat Survei Geologi, Bandung, Special Publ. 36, p. 19-40.
(online at: https://www.researchgate.net/publication/313226301_Pleistocene_geology_palaeontology_and_ archaeology_of_the_Soa_Basin_Central_Flores_Indonesia)

(Ages from Flores classic Mata Menge site: tuff sealing top of fossil layer (with Stegodon florensis and in-situ stone artefacts) with fission track age of 800 ± 70 Ma, while pink tuffaceous silt immediately below main fossil deposit dated as 880 + 70 ka. Tangi Talo site with pygmy Stegodon sondaari and giant tortoise and FT age 900 ± 70 ka. Around 680 ka lake increased in size one leading to deposition of thin-bedded freshwater limestones of Upper limestone. Tektite at surface of Dozu Dhalu site with in-situ artefact and Stegodon florensis)

Suminto, G.D. van den Bergh, I. Saefudin & K. Mano (1996)- The stratigraphy and sedimentology of the hominid skull find site, Grogolan Wetan, Sangiran area, Central Jawa. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 8, p. 51-57.
(New H. erectus skull from upper Bapang Fm, between Upper Tuff and Uppermost Lahar. This is youngest level of hominid fossil occurrence in Sangiran)

Sunardi, E. (2010)- Penelitian magnetostratigrafi dan penerapan satuan stratigrafi polaritas magnet sebagai satuan kronostratigrafi: studi kasus di cekungan Bandung serta daerah Mojokerto dan Sangiran, Jawa. Jurnal Geologi Indonesia 5, 2, p. 137-150.
(online at: http://ijog.bgl.esdm.go.id/index.php/IJOG/article/view/99/99)
('Magnetostratigraphy study and application of magnetic polarity stratigraphy units as chronostratigraphy units: a case study in the Bandung Basin and the Mojokerto and Sangiran regions, Java'. Magnetic polarity reversals of rocks in Sangiran Area (C Java), Mojokerto (E Java; 3 Ma) and Bandung Basin (W Java; 4 Ma) correlated to magnetostratigraphy of Pleistocene formations at Mojokerto)

Suryatman, R.R. Triwurjani, S. Adhityatama, D. Bulbeck, D. Yurnaldi, A. Murdihastomo, J. Wahyudiono, Abdullah & M.D.K. Dhony (2021)- Flaking stone activity in the tradition of iron smelting from the 8th to 17th centuries AD in the Matano region, South Sulawesi, Indonesia. J. Archaeological Science: Reports 36, 102817, p. 1-12.
(online at: https://archaeobib.iseaarchaeology.org/files/original/4a3b3b4241d9db83aaa35e4fc6f8ccc0.pdf)
(New iron smelting sites in Lake Matano region, E Sulawesi (NE of S Sulawesi provicnce), including submerged sites. Belonging to iron-smelting industry developed here in 8th-17th centuries AD. Thousands of chert artifacts excavated from Rahumpu'u lakeshore site were used as strike-a-lights for combustion of fuel for iron smelting. Flaking chert was one of skills possessed by Matano indigenous iron smelters,to produce high quality iron)

Sutikna, T. (2016)- New archaeological research at Liang Bua on the island of Flores: implications for the extinction of *Homo floresiensis* and the arrival of *Homo sapiens* in eastern Indonesia. Ph.D. Thesis, School of Earth and Environmental Sciences, University of Wollongong, p. 1-206.
(online at: https://ro.uow.edu.au/theses/4745/)
(Series of new excavations at Liang Bua, Flores, between 2007 and 2014 led to new interpretation of time of H. floresiensis extinction (~50 ka) and H. sapiens arrival (~46 ka) (no longer necessarily overlap in age))

Sutikna, T., M.W. Tocheri, J.T. Faith, R.D. Awe, H.J. Meijer, E.W. Saptomo & R.G. Roberts (2018)- The spatio-temporal distribution of archaeological and faunal finds at Liang Bua (Flores, Indonesia) in light of the revised chronology for *Homo floresiensis*. J. Human Evolution 124, 52e74, p. 52-74.
(online at: https://www.sciencedirect.com/science/article/pii/S0047248417303160)
(Liang Bua limestone cave in Flores is, the type site of Homo floresiensis. Sdimentary deposits range from ~190 ka to present. Skeletal remains of H. floresiensis ~100-60 ka old; cultural evidence until ~50 ka. Major shifts in faunal and stone artifacts reflect marked changes in paleoecology and hominin behavior. H. floresiensis and Stegodon florensis insularis, along with giant marabou stork and vulture, likely extinct by ~50 ka. Increased use of chert occurs at ~46 ka may characterize Homo sapiens assemblages at Liang Bua)

Sutikna, T., M.W. Tocheri, M.J. Morwood, E.W. Saptomo, Jatmiko, R.D. Awe, Sri Wasisto, K.E. Westaway, M. Aubert et al. (2016)- Revised stratigraphy and chronology for *Homo floresiensis* at Liang Bua in Indonesia. Nature 532, 7599, p. 366-369.
(online at: www.academia.edu/23978420/Revised_stratigraphy_and_chronology_for_Homo_floresiensis_Etc.)

(Skeletal remains and deposits with H. floresiensis dated as 100- 60 ka. Stone artefacts range from ~190-50 ka. Not clear if H. floresiensis survived after 50 ka and potentially encountered modern humans on Flores or other hominins dispersing through SE Asia)

Suyanto A. & C.H.S. Watts (2002)- Verhoeven's giant rat of Flores, Indonesia (*Papagomys theodorverhoeveni* Musser, 1981; Muridae) is a modern species. Treubia 32, 1, p. 87-93.
(online at: https://pdfs.semanticscholar.org/fee8/f23bfe0979528dbe13e57c4a222a8afbf16c.pdf)
(Report of recent specimen in collections of Museum Zoologicum Bogoriense of P. theodorverhoevei, which was generally believed to be extinct. See also Zijlstra et al. 2008)

Suyono (2009)- The study of fossil faunas in the Walanae Basin, Indonesia. Ph.D. Thesis University of Wollongong, Australia, p. 1-115.
(online at: http://ro.uow.edu.au/theses/3058/)
(Mainly on morphology and phylogenetic history of Celebochoerus heekereni, an endemic pig species from Pliocene in Walanae Basin, SW Sulawesi)

Suzuki, M., Wikarno, Budisantoso, I. Saefudin & M. Itihara (1985)- Fission track ages of pumice tuff, tuff layers, and javites of fossil hominid fossil-bearing formations in Java. In: N. Watanabe & D. Kadar (eds.) Quaternary geology of the hominid fossil bearing formations in Java, Geological Research Development Centre (GRDC), Bandung, Special Publ. 4, p. 309-357.
(Eight radiometric ages from 21 pumice tuff layers and 2 javites of Pleistocene of Sangiran. Tuffs from Pucangan Fm 1.16 Ma and Kabuh Fm 0.71- 0.78 Ma). Javites (tektites) 0.71 Ma)

Swisher, C.C., G.H. Curtis, T. Jacob, A.G. Getty, A. Suprijo & Widiasmoro (1994)- Age of the earliest known hominids in Java, Indonesia. Science 263, p. 1118-1121.
(40Ar/39Ar ages from pumice from Mojokerto hominid sites 1.81 and 1.66 Ma, 0.6 million years older than Homo erectus fossils from Olduvai Gorge, and comparable to age of H cf. erectus (H. ergaster) in Kenya. Ages would suggest Homo erectus may have evolved in Asia instead of Africa (NB: these Java age dating results widely disputed in subsequent literature due to erroneous locality information (Huffman et al. 2006, etc.))

Swisher, C.C., G.H. Curtis & R. Lewin (2000)- Java Man- how two geologists' dramatic discoveries changed our understanding of the evolutionary path to modern humans. Scribner, New York, p. 1-244.
(Popular account of events leading to new conclusions on human evolution. (This 'dramatic discovery' is highly controversial; the unusually old age date of Mojokerto hominids probably based on erroneous location information; JTvG)

Swisher, C.C., W.J. Rink, S.C. Anton, H.P. Schwartz, G.H. Curtis & A. Suprijo (1996)- Latest *Homo erectus* of Java; potential contemporaneity with *Homo sapiens* in Southeast Asia. Science 274, p. 1870-1874.
(Hominid fossils from Ngandong and Sambungmacan considered the most morphologically advanced Homo erectus. Dating of fossil bovid teeth collected from hominid-bearing levels gave mean ages of 27 to 53 ka, much younger than previous age estimates for these hominids (results unrealistically young?; see also Grun and Thorne 1997, Indriati et al. 2011; JTvG))

Tassy, P., P. Anupandhanant, L. Ginsburg, P. Mein, B. Ratanasthien & V. Suteethorn (1992)- A new *Stegolophodon* (Proboscidea, Mammalia) from the Early Miocene of northern Thailand. Geobios 25, 4, p. 511-523.
(First complete molars of Stegolophodon from Miocene basins of N Thailand: complete M3 from Mae Mob (M Miocene) and partial mandible with M2-M3 from Na Sm (E Miocene))

Tattersall, I. & J.H. Schwartz (2009)- Evolution of the genus *Homo*. Annual Review Earth Planetary Sciences 37, p. 67-92.
(General review of hominid evolution in last ~2 Myrs. Heterogeneity among 'early African Homo erectus' and no clear link to Asian Homo erectus group. Pithecanthropus (now Homo) erectus now reckoned to be ~0.7- 1.5

Myr old. First truly cosmopolitan Homo is H. heidelbergensis, known from Africa, Europe and China 600 kyr ago. Homo sapiens originated in Africa)

Taufik, L., J.C. Teixeira, B. Llamas, H. Sudoyo, R. Tobler & G.A. Purnomo (2022)- Human genetic research in Wallacea and Sahul: Recent findings and future prospects. Genes (MDPI) 13, 12, 2373, p. 1-11.
(online at https://www.mdpi.com/2073-4425/13/12/2373)
(Review of genetic sequence data from human populations in E Indonesia and New Guinea- Australia. All present-day non-African human populations carry at least 2% Neanderthal genetic ancestry, with Denisovan DNA occurring at trace levels in S and E Asians, but considerably more (~2-6%) in populations living today E of Wallace Line)

Teixeira, J.C., G.S. Jacobs, C. Stringer, J. Tuke, G. Hudjashov, G.A. Purnomo, H. Sudoyo, M.P. Cox et al. (2021)- Widespread Denisovan ancestry in Island Southeast Asia but no evidence of substantial super-archaic hominin admixture. Nature Ecology & Evolution 5, p. 616-624
(online at: https://www.researchgate.net/publication/350299475_Widespread_Denisovan_ancestry_in_Island_Southeast_Asia_but_no_evidence_of_substantial_super-archaic_hominin_admixture)
(Hominin fossil record of Island SE Asia (ISEA) shows at least two endemic 'super-archaic' species (Homo luzonensis in Philippines and H. floresiensis on Flores) were present when anatomically modern humans arrived in region >50,000 years ago. Modern humans across SE Asia (and especially in Australo-Papuan populations) carry genomic traces of ancient interbreeding with Denisovans- a hominin lineage without any fossil record in ISEA. New genetic data corroborate widespread Denisovan ancestry in present-day SE Asia human populations, but no super-archaic admixture signals)

Ter Haar, C. (1934)- *Homo-soloensis*. De Ingenieur in Nederlandsch-Indie (IV) 1, 4, p. IV.51- IV.57.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1934-04.pdf)
(Summary of lecture. Discussion of geological setting of Homo soloensis discovery in 1932 in Solo River terrace deposits at Ngandong, Kendeng Hills, C Java. Associated with rich (Late?) Pleistocene mammal fauna, with Stegodon (primarily forest fauna). Ngandong yielded multiple human skulls, but, reamrably, only skull caps.)

Theunissen, B. (1989)- Eugene Dubois and the ape-man from Java; The history of the first "missing link" and its discoverer. Kluwer Academic Publishers, Dordrecht, p. 1-293.
(Study of the life and scientific contributions of Eugene Dubois, discoverer of 'Java Man' in 1891)

Theunissen, B., J. de Vos, P.Y. Sondaar & F. Aziz (1990)- The establishment of a chronological framework for the hominid-bearing deposits of Java: a historical survey. In: L.F. La Porte (ed.) Establishment of a geologic framework for paleoanthropology, Geological Society of America (GSA), Special Paper 242, p. 39-53.
(Brief historical survey of discovery of Pleistocene mammals and hominid-bearing deposits in Java since mid-1850's, and of attempts to establish chronological framework for Javanese hominids)

Thorne, A. & M.H. Wolpoff (1981)- Regional continuity in Australasian Pleistocene hominid evolution. American J. Physical Anthropology 55, 3, p. 337-349.
(Study of Sangiran 17 Homo erectus skull, interpreted to show similarities with Late Pleistocene Australian hominid)

Tiauzon, A. (2011)- Lithic technology in Song Terus during the late Middle Pleistocene and the early Upper Pleistocene. M.Sc. Thesis, Museum National Histoire Naturelle, Paris, p. 1-96.
(online at: http://hopsea.mnhn.fr/pc/thesis/M2%20Archie_TIAUZON.pdf)

Tjia, H.D. (2006)- Geological evidence for Quaternary land bridges in insular Southeast Asia. In: T. Simanjuntak et al. (eds.) Archaeology: Indonesian perspective, R.P. Soejono's Festschrift, LIPI Press, Jakarta, p. 71-78.
(During most of Quaternary (<1.8 Ma) four major land bridges provided access from SE Asia into Wallacea (= mobile region between Sundaland and Sahul-land): (1) Sabah/ Borneo via Palawan-Mindoro (Philippines), (2)

Sabah/ Borneo via Sulu Archipelago to Mindanao (Philippines), (3) Kangean Gp via Paternoster platform to S Sulawesi, (4) Java/S Kalimantan via Banda Arc to Alor- Timor or via Sumba- Savu-Rote to Timor. Today these land bridges contain passages >200m deep, but probably result of recent tectonic subsidence, not older than 100 ka when modern humans began to populate region)

Tobias, P. (1966)- A re-examination of the Kedung Brubus mandible. Zoologische Mededelingen, Leiden, 41, p. 307-320.
(online at: http://dare.uva.nl/cgi/arno/show.cgi?fid=150000)
(Description of fragment of Pithecanthropus erectus jawbone collected by Dubois at Kedung Brubus in 1890 and described in 1924. Probably from juvenile individual)

Tobias, P.V. (1971)- The brain in hominid evolution. Columbia University Press, New York, p. 1-170.
(online at: https://ia800208.us.archive.org/21/items/braininhominidev38tobi/braininhominidev38tobi.pdf)
(With chapter 5 on Java Homo erectus. Brain capacity of Sangiran and Trinil Homo erectus average 860 cc))

Tobias, P.V. & G.H.R. von Koenigswald (1964)- A comparison between the Olduvai hominines and those of Java and some implications for hominid phylogeny. Nature 204, 4958, p. 515-518.

Tocheri, M.W., C. Orr, S. Larson, T. Sutikna, Jatmiko, E.W. Saptomo, R.A. Due, T Djubiantono, M.J. Morwood & W.L. Jungers (2007)- The primitive wrist of *Homo floresiensis* and its implications for hominin evolution. Science 317, 5845, p. 1743-1745.
(online at: www.researchgate.net/publication/5958398_The_Primitive_Wrist_of_Homo_floresiensis_and_Its_Implications_for_Hominin_Evolution)
(Wrist bones of holotype of Homo floresiensis (LB1) retains morphology primitive for African ape-human clade, and different from Neandertals and modern humans, indicating LB1 is species descended from hominin ancestor that branched off before origin of the clade that includes modern humans, Neandertals, etc.)

Tocheri, M.W., E.G. Veatch, Jatmiko, E.W. Saptomo & T. Sutikna (2022)- *Homo floresiensis*. In: C.F.W. Higham & N.C. Kim (eds.) The Oxford handbook of Early Southeast Asia, Chapter 2, p. 38-69.

Tokunaga, S., H. Oshima, A.A. Polhaupessy & Y. Ito (1985)- A palynological study of the Pucangan and Kabuh Formations in the Sangiran area. In: N. Watanabe & D. Kadar (eds.) Quaternary geology of the hominid fossil bearing formations in Java. Geological Research Development Centre (GRDC), Bandung, Special Publ. 4, p. 199-217.
(Preliminary palynological study of Pleistocene of Sangiran, C Java. Rel. common grass pollen from M-U Pucangan and Lower Kabuh Fms)

Turvey, S.T., J.J. Crees, J. Hansford, T.E. Jeffree, N. Crumpton, I. Kurniawan, E. Setiyabudi et al. (2017)- Quaternary vertebrate faunas from Sumba, Indonesia: implications for Wallacean biogeography and evolution. Proc. Royal Society (London), B, 284, 20171278, p. 1-10.
(online at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5577490/pdf/rspb20171278.pdf)
(New vertebrate fossil deposits on Sumba. Pleistocene deposit at Lewapaku in interior highlands may be close to 1 Ma old, with small Stegodon sumbaensis, tooth of Varanus komodoensis and fragments of giant murids. Holocene cave deposits at Mahaniwa (~2000-3500 BP) with large rats and extinct frugivorous Varanus hooijeri. Sumba Quaternary vertebrates typical island fauna, comparable to fauna of Flores)

Tyler, D.E. (1991)- A taxonomy of Javan hominid mandibles. J. Human Evolution 6, p. 401-420.
(presumably same as Tyler (1992), below). Seven human mandibular remains from Java (Kedung Brubus, Sangiran 1, 5, 6, 8, 9, and 22) are all from single species, Homo erectus)

Tyler, D.E. (1992)- A taxonomy of Javan hominid mandibles. Acta Anthropologica Sinica 11, 4, p. 292-299 *(in Chinese)*.
(online at: http://www.anthropol.ac.cn/EN/Y1992/V11/I04/285)

Tyler, D.E. (1995)- The current picture of hominid evolution in Java. Acta Anthropologica Sinica 14, 4, p. 313-323.
(online at: http://www.anthropol.ac.cn/EN/abstract/abstract806.shtml#1)
(Brief review of Home erectus and Meganthropus from Java. Present-day 'negritos' hunters-gatherers living in remote rainforests of SE Asia may have descended from Ngandong hominids (Homo soloensis), then became largely displaced by wave of farmers (modern Indonesians) from mainland Asia since ~5000 years ago)

Tyler, D.E. (1997)- New and significant fossil finds from Sangiran, Central Java. In: N. Jablonski (ed.) Changing face of East Asia during the Tertiary and Quaternary, Proc. Fourth Conference Evolution of the East Asian Environment, University of Hong Kong Press, p. 498-515.

Tyler, D.E. (2001)- "*Meganthropus*" cranial fossils from Java. J. Human Evolution 16, 2, p. 81-101.
(3 of 12 Homo erectus skulls from E-M Pleistocene of Sangiran may represent 'Meganthropus'. Meganthropus I, II, and III more massive than any known H. erectus specimens, also higher vaulted, smaller brained, and have thick lower occipital planes. May represent species that separated from H. erectus upon its arrival to Java)

Tyler, D.E. (2001)- Three new *Homo erectus* mandibles from Java. J. Human Evolution 16, 2, p. 103-115.
(online at: https://www.academia.edu/27285526/Three_newHomo_erectus_mandibles_from_Java)
(Eleven known Homo erectus manidublar pieces from E-M Pleistocene of Java, all from Sangiran, one from Kedung Brubus. Sangiran 21 (E), Sangiran 22 (F), and Sangiran 37 (G) first described here. Sangiran 21, 22, and 27 from U Pucangan Fm and dated as ~1.2 Ma)

Tyler, D.E. (2001)- Two new "*Meganthropus*" mandibles from Java. J. Human. Evolution 16, p. 151-158.
(Two new mandibles from U Pucangan Fm of Sangiran, C Java. Dated as ~1.2- 1.4 Ma. Morphologically compatible with other"Meganthropus" mandibles described from Java)

Tyler, D.E. (2003)- Sangiran 5 ("*Pithecanthropus dubius*"), *Homo erectus*, "*Meganthropus*" or *Pongo*? J. Human Evolution 18, 3-4, p. 229-241.
(Eleven E-M Pleistocene jaw fragments now known from Java, all but one from Sangiran. Morphologically, they are a mixture of undoubted H. erectus, 'H. meganthropus' and possibly a pongid. 'Pithecanthropus dubius' (Sangiran 5) may not be hominid, but if it is, must be placed with 'H. meganthropus', not H. erectus)

Tyler, D.E. (2004)- An examination of the taxonomic status of the fragmentary mandible Sangiran 5, *(Pithecanthropus dubius), Homo erectus, "Meganthropus"*, or *Pongo*? Quaternary International 117, p. 125-130.
(Morphology of Sangiran 5 mandible fossil, initially named Sangiran 1939 and recovered by Von Koenigswald, with characteristics beyond known range of any H. erectus and must be pongid ape)

Tyler, D.E., N. Jablonski & S. Sartono (1995)- Earliest known monkey fossil from the Indonesian Archipelago: an announcement. In: J.R.F. Bower & S. Sartono (eds.) Palaeo-anthropology: evolution and ecology of *Homo erectus*, Pithecanthropus Centennial Foundation, Leiden University, p. 213-216.
(Monkey fossil from Sangiran. See also Jablonski & Tyler1999 and Larick et al. 2000: not as old as assumed?)

Tyler, D.E., G.S. Krantz & S. Sartono (1995)- The taxonomic status of the '*Meganthropus*' cranial (Sangiran 31) and the '*Meganthropus*' occipital fragment III. In: J.R.F. Bower & S. Sartono (eds.) Palaeo-anthropology: evolution and ecology of *Homo erectus*, Pithecanthropus Centennial Foundation, Leiden, 1, p. 189-202.

Tyler, D.E. & S. Sartono (2001)- A new *Homo erectus* cranium from Sangiran, Java. Human Evolution 16, 1, p. 13-25.
(New H. erectus cranium found by local farmer in 1993 at Sangiran M Pucangan Fm, ~1.6-1.8 Ma in age. Braincase and most of face. Skull longer and narrower than Trinil, possibly female counterpart to Sangiran 17)

Urciuoli, A., J. Kubat, L. Schisanowski, F. Schrenk, B. Zipfel, M. Tawane, L. Bam et al. (2021)- Cochlear morphology of Indonesian *Homo erectus* from Sangiran. J. Human Evolution 165, 103163, p. 1-16.

(Cochlear morphology (bony labyrinth of inner ear) of two Indonesian H. erectus (Sangiran 2 and 4). Most characters found in later humans already present in Indonesian H. erectus, with exception of Neanderthals, which display an autapomorphic condition relative to other Homo species)

Van den Bergh, G.D. (1988)- *Duboisia santeng*, the endemic Pleistocene bovid from Java (Indonesia). Masters Dissertation, University of Utrecht, p. 1-72.
(Duboisia santeng (Dubois' antelope) extinct antelope-like bovid that was endemic to Indonesia in Pleistocene (since then also found in Thailand; JTvG))

Van den Bergh, G.D. (1990)- Sulawesi (Celebes): faunistisch en geologisch ontmoetingspunt der aardschollen. Cranium 7, 2, p. 68-76.
(online at: http://natuurtijdschriften.nl/download?type=document;docid=523240)
('Sulawesi: faunistic and geologic junction of earth plates'. Discussion of present-day endemic fauna of Sulawesi and low diversity Pleistocene 'island fauna' in SW Sulawesi with endemic pig (Celebochoerus heekereni) and dwarfed proboscideans (Elephas celebensis, Stegodon sompoensis, S. cf. trigonocephalus))

Van den Bergh, G.D. (1999)- The Late Neogene elephantoid-bearing faunas of Indonesia and their paleozoogeographic implications. A study of the terrestrial faunal succession of Sulawesi, Flores and Java, including evidence for early hominid dispersal east of Wallace's Line. Scripta Geologica, Leiden, 117, p. 1-419.
(online at: www.repository.naturalis.nl/document/45937)

Van den Bergh, G.D., B.V. Alloway, M. Storey, R. Setiawan, D. Yurnaldi, I. Kurniawan, M.W. Moore, Jatmiko et al. (2022)- An integrative geochronological framework for the Pleistocene So'a Basin (Flores, Indonesia), and its implications for faunal turnover and hominin arrival. Quaternary Science Reviews 294, 107721, p. 1-41.
(online at: https://www.sciencedirect.com/science/article/pii/S0277379122003523)
(Major review of Pleistocene stratigraphy, fossils and stone artefacts of Soa basin, Flores. Age of Homo floresiensis at Liang Bua 100-50 ka. Giant tortoise and dwarf Stegodon sondaari last occurred at Tangi Talo at ~1.3 Ma, not 0.9 Ma as previously thought, disappearing before, or coincident with, earliest hominin arrival (first records of lithic artefacts directly below 1 Ma Wolo Sege Tephra). Artefacts common in younger layers, associated with mid-sized Stegodon florensis and giant rat Hooijeromys nusatenggara. Paleoenvironment of So'a Basin became drier around time of observed faunal transition and arrival of hominins, possibly related to climate change at onset of M-Pleistocene Transition (~1.25 Ma), with increased aridity and monsoonal intensity. Mata Menge site with H. floresiensis-like fossils dated to~0.7 Ma)

Van den Bergh, G.D. & F. Aziz (1991)- Fossil vertebrates from the Walanae Formation and younger Pleistocene deposits, South Sulawesi. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 1, 3, p. 8-11.
(>1500 Late Pleistocene vertebrate fossils recovered from Walanae Fm grey clay layer, deposited on floodplain of N-flowing Walanae River, S Sulawesi. Most abundant is large pig Celebochoerus heekereni, also dwarf Stegodon, giant tortoise Geochelone atlas and large crocodile. Endemic species show S Sulawesi was isolated from SE Asian mainland)

Van den Bergh, G.D., F. Aziz, P.Y. Sondaar & S.T. Hussain (1992)- Taxonomy, stratigraphy, and paleozoogeography of Plio-Pleistocene Proboscideans from the Indonesian islands. Bull. Geological Research Development Centre (GRDC), Bandung, Seri Paleontologi 7, p. 28-58.
(Taxonomic revision of Plio-Pleistocene proboscideans (elephants) from Indonesian region (Java, Sulawesi, Flores, Timor, Sumba, Sangihe, Sumatra))

Van den Bergh, G.D., F. Aziz, P.Y. Sondaar & J. de Vos (1994)- The first *Stegodon* fossils from Central Sulawesi and a new advanced *Elephas* species from South Sulawesi. Bull. Geological Research Development Centre (GRDC) 17, p. 22-39.
(First Stegodon from M-L Pleistocene Napu Fm near Betue, Poso District, C Sulawesi Very similar to S. trigonocephalus from Java. Also new Elephas species from M-L Pleistocene Tanrung Fm of S Sulawesi)

Van den Bergh, G.D., J. de Vos, F. Aziz & M.J. Morwood (2001)- Elephantoidea in the Indonesian region: new *Stegodon* findings from Flores. In: G. Cavaretta et al. (eds.) The World of Elephants, Proc. 1 Int. Congress, Rome, CNRS p. 623-627.
(online at: http://www.sovraintendenzaroma.it/sites/default/files/storage/original/application/7a278b6cbdcf3e33490dcd342e5f6676.pdf)
(Recent discoveries in 1996-1999 of fossil Stegodon on Flores confirm earlier discoveries: (1) Early Pleistocene island assemblage, dated at 0.9 Ma with dwarfed Stegodon sondaari with Varanus komodoensis and giant tortoise remains; (2) M Pleistocene assemblage dated at 0.85- 0.7 Ma, with intermediate to large-Stegodon florensis, with giant rat Hooijeromys nusatenggara and V. komodoensis. This fauna is associated with early human stone tools at various localities. S. florensis closely related to S. trigonocephalus group.)

Van den Bergh, G.D., J. de Vos & P.Y. Sondaar (2001)- The Late Quaternary palaeogeography of mammal evolution in the Indonesian Archipelago. Palaeogeogr. Palaeoclim. Palaeoecology 171, p. 385-408.
(online at: http://users.clas.ufl.edu/krigbaum/6930/vandenbergh_etal_p3_2001.pdf)
(Recent sub-faunas from Sundaland preserved broad connections with SE Asia mainland during Last Glacial: all balanced and mainly mainland elements. Wallacea (Sulawesi, Lesser Sunda islands) was always geographically isolated from mainland, with unbalanced faunas with endemic elements. Significant faunal turnover throughout region around 0.8 Ma (also in Japan, Taiwan). On Java Late Pliocene- E Pleistocene Satir Fauna indicates island conditions)

Van den Bergh, G.D., J. de Vos, P.Y. Sondaar & F. Aziz (1995)- *Homo erectus* in S.E. Asia: time space and migration routes, I. The Flores case. In: J. Gibert et al. (eds.) Proc. Int. Conf. The hominids and their environment during the Lower and Middle Pleistocene of Eurasia, Orce 1995, p. 353-362.

Van den Bergh, G.D., J. de Vos, P.Y. Sondaar & F. Aziz (1996)- Pleistocene zoogeographic evolution of Java (Indonesia) and glacio-eustatic sea-level fluctuations: a background for the presence of *Homo*. Bull. Indo-Pacific Prehistory Association (IPPA) 14 (Chiang Mai Papers), 1, p. 7-21.
(online at: http://ejournal.anu.edu.au/index.php/bippa/article/view/425/414)
(Relation of Java mammal faunal changes in last 2.5 My to global sea level changes not clear. No proof for presence of mammals on Java during first marked glacio-eustatic sea level lowering at 2.4 Ma. Oldest recognizable fauna is Satir fauna, age between 2- 1.5 Ma and indicates island conditions. Isolated conditions continue until ~0.8 Ma, suggested by unbalanced Ci Saat (1.2 Ma) and Trinil faunas (0.9 Ma), when Homo erectus arrived. Major faunal immigration at 0.8 Ma with Kedung Brubus fauna, corresponding with marked lowering of glacio-eustatic sea level)

Van Den Bergh, G.D., R. Due Awe, M.J. Morwood, T. Sutikna, Jatmiko & E.W.Saptomo (2008)- The youngest *Stegodon* remains in Southeast Asia from the Late Pleistocene archaeological site Liang Bua, Flores, Indonesia. Quaternary International 182, p. 16-48.
(online at: www.academia.edu/18237457/The_youngest_stegodon_remains_in_Southeast_Asia_from Etc.)
(Stegodon remains from Late Pleistocene of Liang Bua cave, NW Flores, described as new endemic dwarf subspecies: Stegodon florensis insularis. Hominin activities likely played role in Stegodon bone accumulation at cave. Disappearance of Stegodon from Liang Bua sequence coincides with tuff layer in cave around 11 ka)

Van Den Bergh, G.D., Y. Kaifu, I. Kurniawan, R.T. Kono, A. Brumm, E. Setiyabudi, F. Aziz & M.J. Morwood (2016)- *Homo floresiensis*-like fossils from the early Middle Pleistocene of Flores. Nature 534, 7606, p. 245-248.
(online at: http://udel.edu/~mcdonald/vandenbergh2016.pdf)
(Hominin fossils excavated in 2014 from early M Pleistocene Mata Menge site, Soa Basin, C Flores, include mandible fragment and isolated teeth, similar to Late Pleistocene H. floresiensis of Liang Bua cave. Dated as ~0.7Ma, oldest known hominin remains from Flores. Mata Menge fossils tend to support view that H. floresiensis is dwarfed descendent of early Asian Homo erectus)

Van Den Bergh, G.D., I. Kurniawan, M.J. Morwood, C.J. Lentfer, Suyono, R. Setiawan & F. Aziz (2009)- Environmental reconstruction of the Middle Pleistocene archaeological/ palaeontological site Mata Menge,

Central Flores, Indonesia. In: F. Aziz et al. (eds.) Geology, palaeontology and archaeology of the Pleistocene Soa Basin, Central Flores, Indonesia, Chapter 4, Pusat Survei Geologi, Bandung, Special Publ. 36, p. 59-94.
(M Pleistocene paleontological- archeological Mata Menge site in Soa basin represents lake shore deposits, aged ~0.80- 0.88 Ma. Common Stegodon florensis bones from aged animals that probably died natural death. Despite abundance of stone artefacts in same layers no evidence for butchering. FT age of overlying white tuff 0.75 ± 0.07 Ma (Morwood et al. 1998). Fossils all in 0.5- 1.3m thick 'Unit B'; overlying Unit C tuffaceous siltstone and sand unfossiliferous)

Van den Bergh, G.D., Bo Li, A. Brumm, R. Grun, D. Yurnaldi, M.W. Moore, I. Kurniawan, R. Setiawan, F. Aziz, R.G. Roberts, Suyono, M. Storey, E. Setiabudi & M.J. Morwood (2016)- Earliest hominin occupation of Sulawesi, Indonesia. Nature 529, 7585, p. 208-211.
(online at: https://www.academia.edu/26879805/Earliest_hominin_occupation_of_Sulawesi_Indonesia)
(New excavations at Talepu in Walanae Basin, NE of Maros, with hominid stone artefacts and fossil megafauna (Bubalus, Stegodon, Celebochoerus) from sediments deposited from before 200 ka until ~100ka)

Van den Bergh, G.D., H.J.M. Meijer, R.A. Due, M.J. Morwood, K. Szabo, L.W. van den Hoek Ostende, T. Sutikna, E.W. Saptomo, P.J. Piper & K.M. Dobney (2009)- The Liang Bua faunal remains: a 95 k.yr. sequence from Flores, East Indonesia. J. Human Evolution 57, 5, p. 527-537.
(online at: www.academia.edu/18237450/The_Liang_Bua_faunal_remains_a_95k_yr_sequence_from_F Etc.)
(Excavations at Liang Bua limestone cave on Flores faunal sequence spanning the last 95 ky. Major climatic fluctuations, and two human species: (1) H. floresiensis from 95- 17 ka, and (2) modern humans from 11 ka-Present. Faunal assemblage comprises island gigantism in small mammals and dwarfing of large taxa. Confirms long-term isolation, impoverishment, and phylogenetic continuity of Flores faunal community)

Van den Bergh, G.D., B. Mubroto, F. Aziz, P.Y. Sondaar & J. de Vos (1996)- Did *Homo erectus* reach the island of Flores? In: P. Belwood (ed.) Bull. Indo-Pacific Prehistory Association (IPPA) 14 (Chiang Mai Papers-1), p. 27-34.
(online at: https://journals.lib.washington.edu/index.php/BIPPA/article/view/11585)
(Primitive E- M Pleistocene stone artifacts from Ola Bula Fm near Mata Menge, Central Flores (first discovered by T. Verhoeven in 1956), too old for Homo sapiens, probably made by Homo erectus)

Van den Bergh, G.D., P.Y. Sondaar, J. de Vos & F. Aziz (1996)- The Proboscideans of the South-East Asian islands. In: J. Shoshani & P. Tassy (eds.) The Proboscidea; evolution and palaeoecology of elephants and their relatives, Oxford University Press, chapter 23, p. 240-248.
(Present-day elephants are excellent swimmers. Presence of Pleistocene fossil proboscidean remains on several islands surrounded by deep seas suggests they crossed sea barriers. Proboscideans rel. common in fossil impoverished endemic island faunas, sometimes as the only large mammal. In the Lesser Sunda Arc, only Stegodontidae reported as large mammals on islands Flores, Sumba, and Timor)

Van den Bergh, G.D., U. P. Wibowo, E. Setiyabudi, I. Kurniawan, M. Storey, I.Y.P. Suharyogo & M. Puspaningrum (2019)- The Early Pleistocene terrestrial vertebrate faunal sequence of Java, Indonesia. 79th Annual Meeting Society of Vertebrate Paleontology (SVP), Brisbane 2019, 1p. *(Poster)*
(online at: www.researchgate.net/publication/340984466_The_Early_Pleistocene_terrestrial_vertebrate_faunal_sequence_of_Java_Indonesia)
(Summary of work on oldest mammal-bearing beds in Java, the 575m thick Kali Glagah Fm near hamlets Satir and Cisaat, Bumiayu, largely corroborating 1925-1930 and later work. KGF overlies shallow marine Kali Biuk Fm and is overlain by volcanoclastic Mengger Fm. 39Ar/40Ar dating on hornblendes from pumice clast inMengger Fm 1.53 ± 0.08 Ma (providing minimum age for KGF mammal assemblages). Max. age of basal KGF 2.14 Myr, based on published paleomagnetic work. Oldest mammal fauna on Java is Satir Fauna, from basal 50m of KGF, with age estimated at 2 Ma. Satir Fauna succeeded by Ci Saat composite "fauna", based on scattered finds from middle and upper fluvial KGF)

Van den Brink, L.M. (1982)- On the mammal fauna of the Wajak Cave, Java (Indonesia). Modern Quaternary Research in Southeast Asia 7, Balkema, Rotterdam, p. 177-193.
(On Late Pleistocene mammal and hominid fossils from Wajak cave, Lawah Hill, S of Kediri, Java. Rel. modern fauna with Rusa timorensis, Muntiacus muntjac, Sus scrofa, Tapirus indicus, Rhinoceros sondaicus, Hystrix javanica, Panthera tigris, Homo sapiens and Presbytis cristatus)

Van den Hoek Ostende, L., G.D. van den Bergh & R. Awe Due (2006)- First fossil insectivores from Flores. Hellenic J. Geosciences 41, p. 67-72.
(online at: https://www.repository.naturalis.nl/document/160872)
(Hominid bearing strata from Liang Bua cave on Flores yielded common microvertebrate remains, incl. mandibles of shrews)

Van der Geer, A., G. Lyras, J. de Vos & M. Dermitzakis (2010)- Evolution of island mammals: adaptation and extinction of placental mammals on islands. Wiley and Sons, p. 1-461.
(With reviews of Pleistocene mammal localities and biozones of Java (Ch. 12, p. 172-189), Flores (Ch. 13, p. 190-205), Sulawesi (Ch. 14, p. 206-215), The Philippines (Ch. 15, p. 216-227))

Van der Geer, A., G.A. Lyras & R. Volmer (2018)- Insular dwarfism in canids on Java (Indonesia) and its implication for the environment of *Homo erectus* during the Early and earliest Middle Pleistocene. Palaeogeogr. Palaeoclim., Palaeoecology 507, p. 168-179.
(On 'Trinil Dog' from earliest M Pleistocene of Java. Large Xenocyon (previously Megacyon) present in E Pleistocene and replaced by small Xenocyon (previously Mececyon) during earliest M Pleistocene. Change in body size likely effect of increased competition among carnivores on island. Endemism of terrestrial mammals of earliest M Pleistocene horizon of site Trinil indicates Java marginally isolated)

Van der Geer, A.A.E., G.D. van den Bergh, G.A. Lyras, U.W. Prasetyo, R. Awe Due, E. Setiyabudi & H. Drinia (2016)- The effect of area and isolation on insular dwarf proboscideans. J. of Biogeography 43, 8, p. 1656-1666.
(online at: www.academia.edu/27262337/The_effect_of_area_and_isolation_on_insular_dwarf_proboscideans)
(Study of body size evolution of 22 palaeo-insular fossil elephantoid species across 17 islands support ' island rule', as all but one of the elephants experienced dwarfism on islands. Smallest islands generally harbour smallest elephants. Average body size decrease higher when competitors are present)

Van der Hoop, A.N.J. Thomassen a Thuessink (1932)- Megalithic remains in South-Sumatra. Thesis, W.J. Thieme, Zutphen, p. 1-191.
(Originally published as 'Megalithische oudheden in Zuid-Sumatra', Ph.D. thesis, University of Utrecht. Classic treatise of bronze-age? megalithic statues on the Pasemah Plateau in Barisan Range of S Sumatra. (full name of Dr. van der Hoop is rathr long: Abraham Nicolaas Jan Thomassen a Thuessink van der Hoop)

Van der Hoop, A.N.J. Thomassen a Thuessink (1938)- De Praehistorie. In: F.W. Stapel (ed.) Geschiedenis van Nederlandsch Indie, vol. 1, Uitgeversmaatschappij Joost van den Vondel, Amsterdam, p. 9-111.
(online at: https://www.dbnl.org/tekst/thom052gesc01_01/thom052gesc01_01_0001.php)
(Review of knowledge of prehistory of Netherlands Indies)

Van der Kaars, W.A. & M.A.C. Dam (1995)- A 135,000-year record of vegetational and climatic change from the Bandung area, West-Java, Indonesia. Palaeogeogr. Palaeoclim., Palaeoecology 117, p. 55-72.
(online at: www.researchgate.net/profile/Sander-Van-Der-Kaars/publication/248289473_A_135000-year_record_of_vegetational_and_climatic_change_from_the_Bandung_area_West-Java_Indonesia Etc.)
(Sediment cores from Bandung intramontane basin provide paleoclimatic record for Java for last 135,000 years. Anomalously dry conditions in penultimate glacial period, around 135 ka, and very warm and humid interglacial conditions from 126- 81 ka. Reduction in Asplenium ferns from 81-74 ka suggests drier conditions, while increased numbers indicate slightly wetter climate from 74- 47 ka. Distinctly cooler and possibly drier climate from 47- 20 ka. For Last Glacial Maximum 4-7 °C lower temperatures recorded)

Van der Maarel, F.H. (1931)- Mammalia. In: B.G. Escher et al. (eds.) De palaeontologie en stratigraphie van Nederlandsch Oost-Indie ('Feestbundel K. Martin'), Leidsche Geologische Mededelingen 5, p. 471-484.
(online at: www.repository.naturalis.nl/document/549246)
(Listings of Pleistocene mammal species as known from Indonesia in 1931, with bibliography)

Van der Maarel, F.H. (1932)- Contribution to the knowledge of the fossil mammalian fauna of Java. Wetenschappelijke Mededeelingen Dienst Mijnbouw Nederlandsch-Indie 15, p. 1-208.
(online at: https://resolver.kb.nl/resolve?urn=MMKB18:002206000:pdf)
(Also as Doct. Thesis University of Utrecht, under Prof. L. Rutten. Descriptions of vertebrate fossils collected by the Dienst van den Mijnbouw (Geological Survey) in 1925/26 in residencies of Pekalongan, Bojonegoro, Rembang, Madiun and Solo. First description of E Pleistocene fauna from Bumiayu with Tetralophodon, deer and giant tortoise Geochelone. Unlike Trinil and younger fauna, the Bumiayu (Kali Glagah) faunas do not contain any recent species and are in folded beds (now commonly viewed as 'island fauna'; JTvG))

Van der Meulen, A.J. & G.G. Musser (1999)- New paleontological data from the continental Plio-Pleistocene of Java. In: J.W.F. Reumer & J. de Vos (eds.) Elephants have a snorkel! Papers in honour of Paul Y. Sondaar. Deinsea, 7, p. 361-368. *(online at: www.hetnatuurhistorisch.nl/fileadmin/user_upload/documents-nmr/Publicaties/Deinsea/Deinsea_07/Deinsea_7_028_van_der_Meulen.pdf)*
(Preliminary study of Murinae (rats and mice) from continental Plio-Pleistocene of C and E Java. Correlation of Grenzbank in Sangiran Dome with Trinil H.K. layer confirmed. High numbers of Rattus spp. indicate murid fauna unlike present Sundaic murid fauna and has more Indochinese character. Murids from Satir fauna differ from Trinil H.K. fauna, possibly due to age difference, as indicated by Sondaar et al. biostrat scheme)

Van der Plas, M. (2007)- A new model for the evolution of *Homo sapiens* from the Wallacean islands. PalArch's J. Vertebrate Paleontology 1, 1, p. 1-121.

Van der Vlerk, I.M. (1923)- Een nieuwe vindplaats van fossiele werveldieren op Java. De Mijningenieur 4, p. 67-68.
('A new locality of vertebrate fossils on Java')

Van Es, L.J.C. (1927)- Geologische waarnemingen op Java. II. Trinil. De Mijningenieur. No. 2, 23-24.
('Geological observations on Java, II. Trinil')

Van Es, L.J.C. (1929)- Trinil. Fourth Pacific Science Congress, Batavia/Bandung 1929, Excursion E5, p. 1-14.
(Field guide to Trinil hominid site, C Java)

Van Es, L.J.C. (1929)- The Prehistoric remains in Sampoeng cave, Residency of Ponorogo, Java. Fourth Pacific Science Congress, Batavia/Bandung 1929, 3, Biological Papers, p. 329-340.
(First description of prehistoric arrow heads, bone spatulas, etc. from Goewo Lowo, a rock shelter W of Sampoeng, E of Ponorogo, E Java)

Van Es, L.J.C. (1931)- The age of *Pithecanthropus*. Ph.D. Thesis, Technische Hogeschool Delft, Martinus Nijhoff, The Hague, p. 1-142.
(online at: https://www.delpher.nl/nl/boeken/view?identifier=MMKB18:001677000:00159)
(Thorough review of Pleistocene age of Trinil Beds of C Java, distribution of fossil vertebrates in Java, geology of Kendeng zone, etc.)

Van Es, L.J.C. (1936)- De fossiele menschen van Java. Handelingen 7e Nederlandsch-Indisch Natuurwetenschappelijk Congres, Batavia 1935, p. 60-69.
('The fossil hominids of Java')

Van Es, L.J.C. (1936)- L'homme fossile aux Indes neerlandaises. In: International Geological Congress, Report of XVI Session, Washington, 2, p. 1231-1237.
('Fossil Man in the Netherlands Indies; In French')

Van Essen, H., G. van den Bergh & J. de Vos (2006)- The final solution of the *Cryptomastodon* problem. Morphological correlations between supernumerary teeth in stegodonts and elephants (Proboscidea, Mammalia). Courier Forschungsinstitut Senckenberg, Frankfurt, 256, p. 29-41.
(Response to Hooijer 1984. Specimens of Cryptomastodon martini Von Koenigswald 1933 from Sangiran and Mt. Pati Ayam here viewed as supernumerary tooth of Stegodon trigonocephalus)

Van Heekeren, H.R. (1949)- Voorlopige mededeling over Palaeolitische vondsten in Zuid-Celebes. Oudheidkundig Verslag 1941-1947, Oudheidkundige Dienst in Indonesia, Jakarta, p. 109-110.
(online at: https://www.delpher.nl/nl/tijdschriften/view?coll=dts&identifier=MMKITLV3:002517002:00126)
('Preliminary communication about Paleolithic finds in South Sulawesi'.Important late 1947-early 1948 discovery of Pleistocene bones, molars and artefacts by personnel of the 'Oudheidkundige Dienst' (Archeological Survey) along Walanae River, E of Tjabenge village, NE of Makassar, S Sulawesi. Rel. simple stone tools (scrapers) reminescent of Sangiran artefacts, as described by Von Koenigswald (see also Hooijer 1948-1949 papers on associated mammals))

Van Heekeren, H.R. (1955)- Prehistoric life in Indonesia- A new short guide to the prehistoric collection of the Lembaga Kebudajaan Indonesia "Bataviaasch Genootschap van Kunsten en Wetenschappen", Djakarta. Penerbit Soeroengan, Jakarta, p. 1-59.
(Brief guide to the Prehistoric Indonesia and prehistoric collections in what is now Museum Nasional Indonesia, Jakarta: (1) Paleolithic (Sangiran, etc. with Pithecanthropuso erectus, Homo soloensis, Homo wadjakensis, etc.); (2) Mesolithic (Hoabinhian shell mounds of E Sumatra, etc., with early Homo sapiens; Toalean cave dwellers in Sulawesi, Sampung in E Java, etc.) and (3) Neolithic (arrival of proto-Malayan people around 1500-1000 B.C.; across Java, etc.) and (4) Early Metal Age/ Bronze Age (with kettle drums, megalithic tradition, etc.))

Van Heekeren, H.R. (1957)- The stone age of Indonesia. Verhandelingen Koninklijk Instituut Taal Land en Volkenkunde (KITLV) 21, p. 1-141. *(also 1972 second edition, published as Verhandelingen 61, 247p.)*
(online at: www.oxis.org/books/verhandelingen/heekeren-1957.pdf)
(Review of Pleistocene faunas, hominids and Paleolithic, Mesolithic and Neolithic stone tool industries in Indonesia)

Van Heekeren, H.R. (1958)- The bronze-iron age of Indonesia. Verhandelingen Koninklijk Instituut Taal Land en Volkenkunde (KITLV) 22, p. 1-158.
(online at: www.oxis.org/books/verhandelingen/heekeren-1958.pdf)
(Review of numerous archeological objects from 'Bronze-Iron Age' period, which follows Neolithic in Indonesia)

Van Heekeren, H.R. (1958)- The Tjabenge flake industry from South Celebes. Asian Perspectives 2, p. 77-81.

Van Heekeren, H.R. (1972)- The stone age of Indonesia, 2nd Ed. Verhandelingen Koninklijk Instituut Taal Land en Volkenkunde (KITLV) 61, M. Nijhoff, The Hague, p. 1-247.
(online at: http://booksandjournals.brillonline.com/content/books/9789004286917
(Classic review of archeological discoveries in Indonesia prior to 1972)

Van Heekeren, H.R. (1975)- Chronology of the Indonesian prehistory. In: G.J. Bartstra & W. Arnold (eds.) Modern Quaternary Research in Southeast Asia, Balkema, Rotterdam, p. 47-53.
(Two tables with ages of Pleistocene and Holocene prehistoric sites in Indonesia)

Van Heteren, A.H. (2008)- *Homo floresiensis* as an island form. PalArch's J. Vertebrate Palaeontology 5, 2, p. 1-12.
(online at: http://www.leatherandshoes.nl/wp-content/uploads/2020/12/heteren_2008_5_2.pdf)
(Small-bodied Homo floresiensis displays several island adaptations, also observed among other typical island faunas, indicating that Homo floresiensis might very well have been an endemic island form)

Van Heteren, A.H. & J. de Vos (2012)- Stone implements from Java and Flores: a history of the discoveries. Comptes Rendus Palevol 11, 2-3, p. 181-189.
(online at: https://www.sciencedirect.com/journal/comptes-rendus-palevol/vol/11/issue/2)
(*Stone implements found on Flores similar to those found in Sangiran, C Java, by Von Koenisgswald and Ghosh (1973), confirming crossing to Flores by Homo erectus. Primitive stone artefacts at Mata Menge and Boa Leza associated with Stegodon florensis)*

Van Stein Callenfels, P.V. (1925)- Het eerste Paleolithische werktuig in den Archipel. Oudheidkundig Verslag 1924, Oudheidkundige Dienst in Nederlands Indie, Batavia, p. 127-133.
('The first Paleolithic tool in the Archipel'. Hoabinhian biface axe from NE Sumatra, excavated by missionary Neumann near Batu Kemang in the Deli district of NE Sumatra. Now viewed as Late Paleolithic or Mesolithic)

Van Stein Callenfels, P.V. (1925)- Shell middens in North Sumatra. Illustrated London News, p. 127-133.
(One of the first archeological excavations of 'Hoabinhian'; shell middens of NE Sumatra lowlands, at Sukajadi, with common unifacial and chopper-type pebbles (described earlier by Witkamp (1920) and later dated to ~7500 B.P. (McKinnon, 1991))

Van Stein Callenfels, P.V. (1926)- Bijdrage tot de chronologie van het Neolithicum in Zuid-Oost Azie. Oudheidkundig Verslag 1926, Oudheidkundige Dienst in Nederlands Indie, Batavia, p. 174-180.
('Contribution to the chronology of the Neolithic in SE Asia')

Van Stein Callenfels, P.V. (1929)- A trip to the pre-historic remains in Poenoeng, Sampoeng and Dander. Fourth Pacific Science Congress, Batavia/Bandung 1929, Excursion Guide E5, p. 1-7.
(Brief report on trip to Neolithic stone implement workshop at Punung and cave with Neolithic debris at Sampung (first discovered by Van Es in 1926) in Southern Mountains of Central-East Java)

Van Stein Callenfels, P.V. (1932)- Note preliminaire sur les fouilles dans l'abri-sous-roche du Guwa Lawa a Sampung. In: Hommage du Service Archeologique des Indes Neerlandaises au Premier Congres des Prehistoriens d'Extreme-Orient a Hanoi, 1932, p. 16-32.
('Preliminary note on the excavations in the rock shelter of Guwa Lawa in Sampung, contributed by the Netherlands Indies delegation to the first congress of prehistorians in the Far East, in Hanoi 1932'. Classic text on first systematic archeological excavation in C Java, in remnants of Mesolithic-Neolithic cave dwelling. The 'Sampoeng' cave deposits yielded remnants of Papuan/ Melanesoid humans with large teeth, arrow-heads made of deer antlers, remnants of fireplaces, mollusc shells (mixture of marine and non-marine species, numerous mammal fossils, etc.)

Van Stein Callenfels, P.V. (1934)- Implements used by Solo Man at least 40,000 years ago?: tools associated with the mid-Pleistocene Javanese. Illustrated London News 185, 4974, p. 254-255.

Van Stein Callenfels, P.V. (1935)- An advance in Far Eastern prehistory. Prehistoric kitchen-middens in the Straits Settlements. Illustrated London News, 5 January 1935, p. 13-16.
(Mid-Holocene? prehistoric shell-mounds at Guak Kepah, with relics of an AustraloMelanesoid culture in coastal zone of the 'Wellesley Province' in NW Malay Peninsula. Similar tokitchen middens in NE Sumatra, formed by human 'waste disposal' activities on former beach ridges in the coastal plain)

Van Stein Callenfels, P.V. (1936)- L'industrie osseuse de Ngandong. L'Anthropologie 46, p. 359-362.
('The bone tools industry of Ngandong', C Java)

Van Stein Callenfels, P.V. (1936)- The Melanesoid civilization of Eastern Asia. Bulletin Raffles Museum, Singapore, Series B, 1, p. 41-51.

Van Stein Callenfels, P.V. (1938)- Archeologisch onderzoek in Celebes. Tijdschrift Koninklijk Nederlands Aardrijkskundig Genootschap, Ser. 2, 55, p. 138-142.

('Archeological investigations in Celebes (Sulawesi)')

Van Weers, D.J. (1985)- *Hystrix gigantea*, a new fossil porcupine species from Java (Rodentia: Hystricidae). Senckenbergiana Lethaea 66, p. 111-119.
(Fossil porcupine molars from Sangiran (C Java), collected by Von Koenigswald. Three assigned to new species (Hystrix gigantea n. sp.), one may belong to Hystrix brachyura Linnaeus)

Van Weers, D.J. (1992)- *Hystrix vanbreei* n. sp., a new fossil porcupine from the Pleistocene of Java, with notes on the extant species of the Indonesian Archipelago. Senckenbergiana Lethaea 72, p. 189-197.
(Isolated porcupine teeth and two mandible fragments collected by Van Koenigswald from M Pleistocene of Sangiran, C Java, represent new species Hystrix vanbreei. Also present at Trinil)

Van Weers, D.J. (2003)- The porcupine *Hystrix (Acanthion) brachyura punungensis* subsp. nov. from Late Pleistocene fissure deposits near Punung, Java. Scripta Geologica, Leiden, 126, p. 217-225.
(online at: http://repository.naturalis.nl/document/46244)
(Cheek teeth and mandibular fragment of porcupines from Late Pleistocene fissure deposits near Punung considered new subspecies, Hystrix brachyura punungensis subsp. nov. M Pleistocene specimens allocated to Hystrix brachyura subsp.; those from the Holocene to Hystrix javanica)

Veach, E.G., M.W. Tocheri, T. Sutikna, K. McGrath, E.W. Saptomo, Jatmiko & K.M. Helgen (2019)- Temporal shifts in the distribution of murine rodent body size classes at Liang Bua (Flores, Indonesia) reveal new insights into the paleoecology of *Homo floresiensis* and associated fauna. J. Human Evolution 130, p. 45-60.
(Liang Bua cave stratigraphic sequence spans past ~190 kyr. Numerous vertebrate elements, ~80% rats of six genera: Papagomys, Spelaeomys, Hooijeromys, Komodomys, Paulamys, and Rattus. Changes in size distribution due to shifts in abundances of open habitat-adapted medium-sized rats versus closed habitat smaller ones. Modest increase in open habitats at ~3 ka, likely result of farming by modern human populations. At ~60 ka shift from more open habitats to more closed conditions, along with disappearance of H. floresiensis, Stegodon florensis insularis, Varanus komodoensis, Leptoptilos robustus (marabou stork), and Trigonoceps sp.)

Verhoeven, T. (1952)- Stenen werktuigen uit Flores. Anthropos 47, 1-2, p. 95-98.
(First report on discovery of human stone stools in caves and on hilltops in Manggarai, W Flores, in 1950)

Verhoeven, T. (1953)- Eine Mikrolithenkultur in Mittel- und West-Flores. Anthropos 48, 3-4, p. 597-612.
(Pleistocene hominid stone tool artifacts from 'abris' and caves in limestone areas of W and C Flores, discovered during 1950-1952 surveys (see also Van Heekeren (1950s) and Bednarik, 2000)

Verhoeven, T. (1958)- Proto-Negrito in den Grotten auf Flores (Indonesie). Anthropos 53, 1-2, p. 229-232.
('Proto-Negrito in the caves of Flores'. Discussion of multiple Late Pleistocene or E Holocene hominid skeletal remains of relatively small humans (1.46m), found in caves in Flores in 1952-1955. Interpreted as likely 'negrito' or 'proto-negrito' remnants by Huizinga and Von Koenigswald (humans are associated with common 'flake-culture' tools; and their main diet was probably giant rats and freshwater molluscs (see also Hooijer, 1950s)

Verhoeven, T. (1958)- Pleistozane Funde in Flores. Anthropos 53, 1-2, p. 264-265.
(First report of large fossilized E Pleistocene mammal bones Stegodon molars in Ola Bula area, northern C Flores, associated with E Paleolithic hominid stone tools. Includes molars, identified (from photos) by Von Koenigswald and Hooijer as Pleistocene Stegodon trigonocephalus, one of the first reports of crossings of Asian fauna across the Wallace Line. Also report finding Pleistocene tektite near Boawae)

Verhoeven, T. (1959)- Die Klingenkultur der Insel Timor. Anthropos 54, 5-6, p. 970-972.
(online at: https://ur.booksc.me/book/47106085/042bb6)
('The stone tool culture of Timor Island'. Stone tools first found on Timor by F. Sarasin and A. Buhler in caves of Niki-Niki in C Timor. In 1954 Timor expedition additional tools identified in 15 additional caves)

Verhoeven, T. (1964)- *Stegodon*-fossilien auf der Insel Timor. Anthropos 59, 3-4, p. 634-636.
('Stegodon fossils on the island of Timor'.Discovery of molars, tusk and bones of Pleistocene Stegodon elephantoi, E of Atambua in 1964. Associated with Early Paleolithic stone tool, and overlying clay rich in freshwater molluscs (see also Maringer & Verhoeven, 1970))

Verhoeven, T. (1968)- Vorgeschichtliche Forschungen auf Flores, Timor und Sumba. In: Gedenkschrift P.W. Schmidt, Anthropica, Studia Instituti Anthropos 21, 3/4, p. 393-403.
('Prehistoric investigations on Flores, Timor and Sumba')

Veth, P., M. Spriggs, A. Jatmiko & S. O'Connor (1998)- Bridging Sunda and Sahul: The archaeological significance of the Aru Islands, southern Moluccas In: G.J. Bartstra (ed.) Bird's Head approaches; Irian Jaya studies; a programme for interdisciplinary research, Modern Quaternary Research in Southeast Asia 15, Balkema, Rotterdam, p. 157-177.
(online at: https://www.papuaerfgoed.org/sites/default/files/collectie/files/2006-05/Bartstra_1998_Birds.pdf)
(Review of archeological research in Aru Islands and surrounding Maluku islands)

Volmer, R., C. Hertler & A. van der Geer (2015)- Niche overlap and competition potential among tigers (*Panthera tigris*), sabertoothed cats (*Homotherium ultimum, Hemimachairodus zwierzyckii*) and Merriam's Dog (*Megacyon merriami*) in the Pleistocene of Java. Palaeogeogr. Palaeoclim. Palaeoecology 441, 4, p. 901-911.
(online at: www.academia.edu/34299156/Niche_overlap_and_competition_potential_among_tigers Etc.)
(M Pleistocene site of Sangiran where tigers co-occurred with machairodonts (Hemimachairodus zwierzyckii and Homotherium ultimum) and large Merriam's Dog (Megacyon merriami). Tigers did not increase body mass before Ngandong faunal level)

Volmer, R., A.A.E. van der Geer, P.A. Cabrera, U.P. Wibowo & I. Kurniawan (2019)- When did Cuon reach Java? Reinvestigation of canid fossils from *Homo erectus* faunas. Geobios 55, p. 89-102.
(online at: https://www.academia.edu/44261113/When_did_Cuon_reach_Java_Reinvestigation_of_canid_fossils_from_Homo_erectus_faunas)
(Revision of canid (dog) fossi lmaterial from Sangiran Dome. Previously described species Cuon sangiranensis, C. crassidens, C. alpinus, C. javanicus, and C. trinilensis. Cuon already present in Java during E-M Pleistocene and probably arrived as part of immigration events during Kedung Brubus faunal stage)

Volz, W. (1907)- Das geologische Alter der *Pithecanthropus*-Schichten bei Trinil, Ost-Java. Neues Jahrbuch Mineralogie Geologie Palaontologie, 100-year Festband 1907, p. 256-271.
(online at: https://www.biodiversitylibrary.org/item/155807#page/290/mode/1up)
('The geological age of the Pithecanthropus Beds near Trinil. East Java'. Age of Pithecanthropus beds probably Middle Diluvium (=M Pleistocene). Interprets rel. massive, volcanics-rich Pleistocene hominid-bearing Trinil sands as lahar-style mudflow deposits from Lawu volcano. With 3 profile sections (from E. Dubois?))

Von Heine-Geldern, R. (1945)- Prehistoric research in the Netherlands Indies. In: P. Honig & F. Verdoorn (eds.) Science and scientists in the Netherlands Indies, Board for Netherlands Indies, Surinam and Curacao, New York, p. 129-167.
(online at: https://archive.org/details/in.ernet.dli.2015.183759)
(Thorough review of prehistoric research in Indonesian Archipelago until 1940, by Austrian anthropologist-archaeologist Prof. Robert Freiherr von Heine-Geldern, originally from Vienna. "In memory of P.V. van Stein Callenfels")

Von Koenigswald, G.H.R. (1933)- Beitrag zur Kenntnis der fossilen Wirbeltiere Javas I. Wetenschappelijke Mededeelingen Dienst Mijnbouw Nederlandsch-Indie 23, p. 1-127.
('Contribution to the knowledge of the fossil vertebrate fauns of Java, part 1'. Studies of Pleistocene vertebrate faunas of Java, mainly collected during geological survey Java mapping)

Von Koenigswald, G.H.R. (1933)- Ein neuer Urmensch aus dem Diluvium Javas. Centralblatt Mineralogie Geologie Palaontologie 1933, B, 1, p. 29-42.
(On new Pleistocene hominid Homo (Javanthropus) soloensis, discovered recently by Oppenoorth)

Von Koenigswald, G.H.R. (1934)- Zur Stratigraphie des javanischen Pleistocan. De Ingenieur in Nederlandsch-Indie (IV) 1, 11, p. IV.185- IV.201.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1934-11.pdf)
('On the stratigraphy of the Pleistocene of Java'. On characteristics and stratigraphic position of Pleistocene Djetis, Trinil, Ngandong mammal faunas. With map of mammal localities in C and NE Java, etc.)

Von Koenigswald, G.H.R. (1934)- Die Spezialisation des lncisivengebisses bei den javanischen Hippopotamidae. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 37, 9, p. 653-659.
(online at: https://dwc.knaw.nl/DL/publications/PU00016621.pdf)
('The specialization of the incisors of Javanese Hippopotamidae'. Several species of Pleistoceen hippopotamids with reduced incisor teeth)

Von Koenigswald, G.H.R. (1935)- Die fossilen Saugetierfaunen Javas. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 38, 2, p. 188-198.
(online at: https://dwc.knaw.nl/DL/publications/PU00016677.pdf)
('The fossil mammal faunas of Java'. Pliocene-Pleistocene mammals grouped into 7 faunas, from young to old: 1. Sampoeng, 2. Ngandong, 3. Trinil, 4. Djetis/ Jetis, 5. Kali Glagah (Late Pliocene), 6. Tji Djoelang (S of Cirebon; M Pliocene; with anthracotheriid Merycopotamus) and 7. Tji Sande (3 km NW of Loerahgoeng; S of Cirebon; with Aceratherium boschi rhinoceros molar= latest Miocene- earliest Pliocene and oldest mammal fossil from Java). No figures)

Von Koenigswald, G.H.R. (1935)- Bemerkungen zur fossilen Saugetierfauna Javas, I. Uber das Vorkommen von *Nestotherium* und *Hyaena* in der Djetis-Fauna Mittel-Javas. De Ingenieur in Nederlandsch- Indie (IV) 2, 7, p. IV.67- IV.70.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1935-07.pdf)
('Remarks on the fossil mammal faunas of Java, I' On the occurrence of Nestoritherium and Hyena in the Jetis fauna of Central Java'. Perissodactyl fossils from Sangiran, Hyeana from Trinil)

Von Koenigswald, G.H.R. (1935)- Bemerkungen zur fossilen Saugetierfauna Javas, II. Eine Djetis-Fauna aus den Tambakan-Schichten der Gegend von Soebang, West-Java. De Ingenieur in Nederlandsch-Indie (IV) 2, 10, p. IV.85- IV.88.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1935-10.pdf)
('Remarks on the fossil mammal faunas of Java, II. On a Pleistocene Djetis fauna from the Tambakan beds SE of Subang, W Java'. Collected by C. Harloff. With Stegodon, Rhinoceros, Hippopotamus, Bubalus, etc.))

Von Koenigswald, G.H.R. (1935)- Over enkele fossiele zoogdieren van Java. Tijdschrift Koninklijk Nederlands Aardrijkskundig Genootschap, 2, 52, 4, p. 539-543.
(online at: http://resolver.kb.nl/resolve?urn=MMUBA13:001683001:pdf)
('On some fossil mammals of Java'. Popular review of skeletons of mammals from Java in various museums, incuding Late Pliocene hippo Hippopotamus (Hexaprotodon) simplex from Kali Glagah/ Bumiayu), M Pleistocene deer Cervus lydekkeri from Trinil, C Java and Late Pleistocene extinct elephantoid Stegodon trigonocephalusand giant water buffalo Bos bubalis palaeokerabau from Ngandong near Ngawi, C Java)

Von Koenigswald, G.H.R. (1935)- Das Neolithicum der Umgebung von Bandoeng. Tijdschrift Indische Taal Land en Volkenkunde, Batavia, 75, 3, p. 394-419.
(online at: https://ia802905.us.archive.org/12/items/in.ernet.dli.2015.283914/2015.283914.Tijdschrift-Voor.pdf)
('The Neolithic in the surroundings of Bandung')

Von Koenigswald, G.H.R. (1935)- Rapport over de vindplaatsen van fossiele gewervelde dieren uit de Solo-vallei. Dienst van den Mijnbouw, Bandung. *(Unpublished report)*
('Report on the discovery localities of fossil vertebrate animals from the Solo valley')

Von Koenigswald, G.H.R. (1936)- Ein fossiler Hominide aus dem Altpleistocan Ostjavas. De Ingenieur in Nederlandsch-Indie (IV) 3, 8, p. IV.149- IV.157.
(online at: https://www.stichtingblauwelijn.nl/assets/files/1936-08.pdf)
('A fossil hominid from the Early Pleistocene of East Java'. Discovery of a small hominid skull from Upper Pucangan beds near Mojokerto area in February 1936 in area being mapped by J. Duyfjes. Probably skull of a child. Von K here proposes the new name Homo modjokertensis)

Von Koenigswald, G.H.R. (1935)- Eine fossile Saugetierfauna mit *Simia* aus Sudchina. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 38, 8, p. 872-879.
(online at: http://https://dwc.knaw.nl/DL/publications/PU00016780.pdf)
('A fossil mammal fauna with Simia from South China'. On Pleistocene fossils from Chinese pharmacies in Hong Kong and SE Asia. With teeth of various ape species, including giant ape Gigantopithecus blacki n.gen., n.sp. All fossils believed to originate from sites in southern China)

Von Koenigswald, G.H.R. (1936)- Erste Mitteilung uber einen fossilen Hominiden aus dem Altpleistocan Ostjavas. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 39, 8, p. 1000-1009.
(online at: https://dwc.knaw.nl/DL/publications/PU00016947.pdf)
('First communication on a fossil hominid from the Early Pleistocene of East Java'. Same paper as above on discovery of fossil child skull in U Pucangan beds at N flank Kedung-Waru anticline near Mojokerto, named Homo modjokertensis)

Von Koenigswald, G.H.R. (1936)- Uber Altpalaeolithisch Artefakte von Java. Tijdschrift Koninklijk Nederlands Aardrijkskundig Genootschap (2) 53, p. 41-44.
(online at: https://resolver.kb.nl/resolve?urn=MMUBA13:001680001:pdf)
('About Early Paleolithic artifacts from Java'. First reported stone tools from Solo River terrace near Ngandong and Kuwung (von Koenigswald also found stone tools in Trinil Horizon near Solo in 1934, but not published yet))

Von Koenigswald, G.H.R. (1936)- Early Palaeolithic stone implements from Java. Bull. Raffles Museum, Singapore, B, 1, p. 52-60.
(Discoveries of small stone flake tools in fluvial M Pleistocene Kabuh Fm deposits of Ngebung Hills, NW Sangiran Dome, presumably made by Pithecanthropus erectus, mainly from chalcedony. Also Paleolithic stone implements from Solo River High Terrace gravels at Ngandong, C Java. 'Early Acheulian' lithic tools from Baksoka River Valley near Pacitan (for occurrence and position of Sangiran artifacts see also Bartstra, 1974))

Von Koenigswald, G.H.R. (1937)- Ein Unterkieferfragment des *Pithecanthropus* aus den Trinilschichten Mitteljavas. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, 40, 10, p. 883-893.
(online at: https://dwc.knaw.nl/DL/publications/PU00017126.pdf)
(On new lower jaw fragment of Pithecanthropus from Trinil Beds at Sangiran Dome, C Java)

Von Koenigswald, G.H.R. (1938)- Ein neuer *Pithecanthropus* Schadel. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam 41, 2, p. 185-192.
(online at: https://dwc.knaw.nl/DL/publications/PU00017154.pdf)
('A new Pithecanthropus skull'. New hominid skull from basal Trinil Beds at Sangiran Dome, C. Java)

Von Koenigswald, G.H.R. (1938)- Nieuwe *Pithecanthropus*-vondsten uit Midden-Java. Natuurkundig Tijdschrift voor Nederlandsch-Indie 98, p. 195-207.
('New Pithecanthropus finds from Central Java'. Review of known hominid fossils from Java)

Von Koenigswald, G.H.R. (1938)- De nieuwe *Pithecanthropus*-vondsten en hun betekenis voor de afstamming van den mensch. Handelingen 8e Nederlandsch-Indisch Natuurwetenschappelijk Congres, Soerabaia, July 1938, p. 486-488.
(online at: https://resolver.kb.nl/resolve?urn=MMKB31:042287000:pdf)
('The new Pithecanthropus discoveries and their significance for the origin of humans'. Very brief review of discoveries at Mojokerto, Trinil, etc.))

Von Koenigswald, G.H.R. (1939)- Das Pleistocan Javas. In: R. Grahmann & L.F. Zotz (eds.) Quartar- Jahrbuch Erforschung des Eiszeitalters und seiner Kulturen, De Gruyter, Berlin, 2, p. 28-53.
(online at: https://journals.ub.uni-heidelberg.de/index.php/qu/article/download/82655/76869)
('The Pleistocene of Java'. Overview of Java Pleistocene stratigraphy and vertebrate/ hominid finds, including 'Punung Fauna' not described elsewhere)

Von Koenigswald, G.H.R. (1940)- Neue *Pithecanthropus*-Funde 1936-1938, ein Beitrag zur Kenntnis der Praehominiden. Wetenschappelijke Mededeelingen Dienst Mijnbouw Nederlandsch-Indie 28, p. 1-233.
('New Pithecanthropus finds 1936-1938; a contribution to the knowledge of the pre-hominids')

Von Koenigswald, G.H.R. (1948)- Aantekeningen bij het *Pithecanthropus*-vraagstuk. Oratie Buitengewoon Hoogleraar in Stratigrafie en Paleontologie, Universiteit van Utrecht (Inaugural Lecture), 29.11.1948, p. 1-19.
('Notes about the Pithecanthropus question'. Inaugural lecture at appointment as Extrordinary Professor in Stratigraphy and Paleontology at the University of Utrecht (followed by promotion to Full Professor in 1951, until retirement in 1968))

Von Koenigswald, G.H.R. (1949)- The fossil hominids of Java. In: R.W. van Bemmelen, The geology of Indonesia and adjacent archipelagos, Nijhoff, The Hague, 1, p. 106-107.

Von Koenigswald, G.H.R. (1949)- Vertebrate stratigraphy of the Neogene and Quaternary. In: R.W. van Bemmelen, The geology of Indonesia and adjacent archipelagos, Nijhoff, The Hague, 1, p. 91-94.

Von Koenigswald, G.H.R. (1950)- Ein Elephant der *planifrons* Gruppe aus dem Pliocaen West-Javas. Eclogae Geologicae Helvetiae 43, 2, p. 268-274.
(online at: https://www.e-periodica.ch/digbib/view?pid=egh-001:1950:43#322)
('An elephant of the planifrons group from the Pliocene of W Java'. Late Pliocene elephant teeth from Kali Glagah Fm beds in 'Tji Djulang' (Tji Djoelang, Cijulang) river, collected in 1935, in Bumiayu area, S of Tegal. Rel. small and more primitive representative of 'A. planifrons' group from Siwaliks, India, described as Archidiskodon praeplanifrons n. sp. Associated with Pliocene 'Cheribonian' mollusc assemblage)

Von Koenigswald, G.H.R. (1950)- Fossil hominids from the Lower Pleistocene of Java. Reports 18th International Geological Congress, Great Britain 1948, 9, p. 59-61.

Von Koenigswald, G.H.R. (1951)- History of the discovery and geology and paleontology of Solo man. Introduction to: F. Weidenreich, Morphology of Solo man, Anthropological Papers American Museum Natural History 43, 3, p. 211-221.
(online at: https://digitallibrary.amnh.org/handle/2246/297)
(Review of discovery of skull fragments of Solo Man and associated mammal faunas at Ngandong, along Solo River in Kendeng zone, by Geological Survey geologist C. ter Haar in 1931 (the first Elephas molar from the 14m terrace deposits had already been reported by Elbert of the Selenka Expedition). Subsequent skull material excavated by Oppenoorth and Von Koenigswald in 1931-1933. Suggestions of cannibalism by Solo Man)

Von Koenigswald, G.H.R. (1952)- Fossil Sirenians from Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen B 55, 5, p. 610-613.
(On fossil sea cows ('dugong') from Java, incl. molar of new species Indosiren javanense, from U Miocene at Tji Paringan, Nyalindung, W Java. Also Eocene rib bone from U Eocene of Nanggulan?)

Von Koenigswald, G.H.R. (1952)- Evidence of a prehistoric Australo-melanesoid population in Malaya and Indonesia. Southwestern J. of Anthropology 8, 1, p. 92-96.
(Comment on Hooijer (1950) paper of same title, who suggested presence of big-teeth people resembling Australian aboriginals and Melanesians, who once lived in Malaya-Indonesian region, were result of local evolution. Von K disputes this, and argues that a mainly macrodont human population of Australo-Melanesian affinities was well established across SE Asia, then became extinct or was pushed to the E by the first Indonesians from E Asia in late Neolithic time (most authors believe this happened at ~2000 BC (this two-wave human migration into Malaysia-Indonesia was called the 'Two Layer' model by Prof. Teuku Jacob in 1967)

Von Koenigswald, G.H.R. (1952)- *Gigantopithecus blacki* von Koenigswald, a giant fossil hominoid from the Pleistocene of southern China. Anthropological Papers American Museum Natural History, New York, 43, 4, p. 291-325.
(online at: http://digitallibrary.amnh.org/handle/2246/298)
(Gigantopithecus blacki von Koenigswald known from four molars bought in Chinese drugstores in Hong Kong and Canton. Molars are largest known of any higher primate. Cannot be regarded as ancestral to man. Part of the (early) M Pleistocene Ailuropoda-orang fauna of S China)

Von Koenigswald, G.H.R. (1952)- Grepen uit de praehistorie van Indonesie. Sticusa Jaarboek 1952, p. 23-36.
('Topics from the prehistory of Indonesia'. Hard to find publication)

Von Koenigswald, G.H.R. (1953)- Extreme- Orient. In: H.V. Vallois & H. Movius, Catalogue des hommes fossiles, Comptes Rendus 19th Session International Geological Congress, Algeria 1952, 5, p. 329-341.
(Far East chapter in Catalogue of fossil humans)

Von Koenigswald, G.H.R. (1954)- Fossil hominids from Java. Indonesian J. Natural Science (Majalah Ilmu Alam untuk Indonesia) 110, p. 194-199.
(Listing of known hominid discoveries on Java, reprinted from Catalog of hominid fossils published at 19th International Geological Congress, Alger 1952. Localities include Kedung Brubus, Mojokerto, Ngandong, Sonde, Trinil and Wajak)

Von Koenigswald, G.H.R. (1955)- The Pleistocene of Java and the Plio-Pleistocene boundary. Actes IV Congres Association Internationale pour l' l'Etude du Quaternaire (INQUA), Rome 1953, p. 1-11.

Von Koenigswald, G.H.R. (1956)- Remarks on the correlation of mammalian faunas of Java and India and the Plio-Pleistocene boundary. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 59, p. 204-210.

Von Koenigswald, G.H.R. (1956)- The geological age of Wadjak man from Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 59, p. 455-457.

Von Koenigswald, G.H.R. (1956)- Der Solo-Mensch von Java: ein tropischer Neanderthaler. In G.H.R. Von Koenigswald (ed.) Hundert Jahre Neanderthaler, Bohlau, Koln, p. 21-26.
('Solo Man from Java: a tropical Neanderthaler')

Von Koenigswald, G.H.R. (1956)- Meeting prehistoric man. Thames and Hudson, London, p. 1-208.
(Popular book on discovery of early hominids, particularly on Java)

Von Koenigswald, G.H.R. (1956)- Fossil mammals from the Philippines. Proc. 4th Far-Eastern Prehistoric Congress, National Research Council of the Philippines, Quezon City, 1, Paper 22, p. 339-369.
(online at: www.rhinoresourcecenter.com/pdf_files/124/1248937364.pdf)
(Descriptions of Pleistocene mammal fossils collected by O. Beyer. With first descriptions of Rhinoceros philippinensis, Stegodon luzonensis, Elephas beyeri. Also present Stegodon cf. trigonocephalus, S. mindanensis, etc. Incl. probable M Pleistocene association of Stegodon, pre-Neolithic stone implements and tektites on Luzon, off Manila- Ipo road)

Von Koenigswald, G.H.R. (1958)- Preliminary report on a newly-discovered Stone Age culture from northern Luzon, Philippine Islands. Asian Perspectives 2, 2, p. 69-70.
(Common remains of Pleistocene mammals found during visit by Von K to island of Anda in late 1953, Pangasinan, incl. elephants, Stegodon, Bos, Sus and Cervus, in layers that also contained "astonishing assemblage of tektites" ('Anda-type' of Prof. O. Beyer). This is second occurrence of tektites in association with fossil mammals (first in upper Trinil layer at Sangiran, C Java). Specialization of Elephas and Stegodon on Anda suggests both sites are contemporaneous. In late 1957 von K also collected 'Cabalwanian' stone tools in little known site in Cagayan valley, N Luzon, which he believed were made by Homo erectus)

Von Koenigswald, G.H.R. (1959)- A mastodon and other fossil mammals from Thailand. Royal Department of Mines, Bangkok, Report of Investigation 2, p. 25-34.

Von Koenigswald, G.H.R. (1962)- Das absolute Alter des *Pithecanthropus erectus* Dubois. In: G.Kurth (ed.) Evolution und hominisation (Festschrift G. Heberer), Fischer, Stuttgart, p. 112-119.
('The absolute age of Pithecanthropus erectus Dubois')

Von Koenigswald, G.H.R. (1963)- Fossil pygmy Suidae from Java and China. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 66, p. 192-197.

Von Koenigswald, G.H.R. (1964)- Potassium-Argon dates and early man: Trinil. Reports 6th. Int. Congress Quaternary, Warsaw 1961, 4, p. 325-327.

Von Koenigswald, G.H.R. (1968)- Observations upon two *Pithecanthropus* mandibles from Sangiran, Central Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 71, 2, p. 99-107.

Von Koenigswald, G.H.R. (1968)- The real date of Java Man. In: G. Kurth (ed.) Evolution und hominisation, 2nd ed., G. Fischer, Stuttgart, p. 117-125.
(Incl. mention of 710,000 yr radiometric age of tektite horizon in hominid-bearing Trinil Beds (Kabuh Fm) in Sangiran, C Java)

Von Koenigswald, G.H.R. (1973)- The oldest hominid fossils from Asia and their relation to human evolution. Proc. Symposium L'Origine dell'uomo, Rome 1971, Accademia Nazionale dei Lince 182, p. 97-118.

Von Koenigswald, G.H.R. (1974)- Fossil mammals of Java. VI, Machairodontinae from the Lower Pleistocene of Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 77, p. 267-273.
(Description of jaw and teeth of two genera of Pleistocene sabre tooth cat from Jetis Fauna, Sangiran, C. Java)

Von Koenigswald, G.H.R. (1975)- Early Man in Java: catalogue and problems. In: R.H. Tuttle (ed.) Paleoanthropology, morhology and paleoecology, Mouton Publishers, The Hague, p. 303-309.
(Ngandong fauna with Homo soloensis ('Solo Man') is Java representative of Neanderthal Man. Associated with bird fossils suggesting colder climate than today, and water buffaloes with horn spreads >2m, indicating more open country, not typical rainforest conditions. Advanced bone culture and completely rounded stone balls also point to younger hominid population than Sangiran/ Trinil Homo erectus. Etc.)

Von Koenigswald, G.H.R. (1976)- Climatic changes in Java and Sumatra during the Upper Pleistocene. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B79, 4, p. 232-234.
(Mammal fauna of U Pleistocene of Java (Ngandong) not typical of tropical forest, but more open country. Also crane birds on Java and mountain goat on Sumatra suggest 6-9° C temperature drop during last glaciation)

Von Koenigswald, G.H.R. (1976)- Evolution of man. University of Michigan Press, Ann Arbor, Revised edition, p. 1-158.

Von Koenigswald, G.H.R. (1976)- The importance of Java for the early history of man. J. Medical Sciences 8, 3, p. 87-90.
(online at: https://jurnal.ugm.ac.id/bik/article/view/4757/4009)
(Brief review of five different types of early hominids known from Java. Summary of lecture given after receiving Dr. h.c. degree from Gadjah Mada University, Yogyakarta. Oldest mammalian fossils on Java in West Java (Cijulang, Kali Glagah). Two main episodes of mammalian migration: (1) Sivamalayan migration, from India region; (2) Sinomalayan migration, from China region via Philippines, Kalimantan)

Von Koenigswald, G.H.R. (1978)- Selachia (Pisces) from the black clay of Sangiran, Central Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 81, 3, p. 364-369.
(Lower Pleistocene 'black clay' of Sangiran deposited in fresh water lake according to mollusc fauna. Presence of fossil teeth of marine fish Pristiopsis cf P. microdon (sawfish) and Eulamia gangetica (river shark) that can enter and live in fresh water indicates likely outlet of lake into open sea)

Von Koenigswald, G.H.R. (1978)- Lithic industries of *Pithecanthropus erectus* of Java. In: F. Ikawa-Smith (ed.) Early Palaeolithic in South and East Asia, Mouton Publishers, The Hague, p. 23-27.
(Sangiran area of C Java yielded five skulls of Pithecanthropus erectus since 1937, with average K-Ar ages of 830,000 years. Stone implements known from M Pleistocene fluviatile part of Sangiran section (most not found in-situ). Flakes rarely larger than 5-6 cm)

Von Koenigswald, G.H.R. (1980)- Neue Einblicke in die Geschichte der Hominiden. Annalen Naturhistorischen Museums Wien, 83, p. 181-195.
(online at: https://www.zobodat.at/pdf/ANNA_83_0181-0195.pdf)
('New insights in the history of the hominids'. Review of recent developments in hominid studies in Africa, China and SE Asia. One of the more surprising developments is rel. high K/Ar radiometric age (1.9 Ma) of tuffs associated with 'Mojokerto child' Homo erectus by Curtis & Jacob. Tektites in upper tuffs of Sangiran, 20m above Pithecanthropus II horizon, dated at ~0.72 Ma)

Von Koenigswald, G.H.R. & A.K. Ghosh (1973)- Stone implements from the Trinil Beds of Sangiran, Central Java. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B76, 1, p. 1-34.
(Crude stone tools from Trinil Beds at Sangiran believed to be of human origin (Pithecanthropus erectus; first discovered by vK in 1936, but these finds were initiallly dismissed until similar tools of similar age were found in Flores in 1970 and 1990s; JTvG))

Von Koenigswald, G.H.R. & F. Weidenreich (1938)- Discovery of an additional *Pithecanthropus* skull. Nature 142 (3598), p. 715.
(Discovery of almost complete brain case of Pithecanthropus in 1937, in Trinil Fm of Sangiran, C Java)

Von Koenigswald, G.H.R. & F. Weidenreich (1939)- The relationship between *Pithecanthropus* and *Sinanthropus*. Nature 144 (3657), p. 926-929.
(online at: https://www.nature.com/articles/144926a0.pdf)
(Great similarities between first skull of Sinanthropus (S. pekinensis) found in China and Pithecanthropus materials from Central Java (P. erectus, from Trinil, Sangiran). Possibly same genus/ species)

Walters, I. (2002)- Early hominids in SE Asia: older, younger, smarter and more. In: P. Kershaw et al. (eds.) Bridging Wallace's Line: the environmental and cultural history and dynamics of the SE Asian- Australian region, Advances in Geology 34, Catena Verlag, p. 255-263

Wastl, J. (1939)- Praehistorische Menschenreste aus dem Muschelhugel von Bindjai-Tamiang in Nord-Sumatra. In: Kultur und Rasse, Otto Reche zum 60. Geburtstag gewidmet von Schulern und Freunden, Munchen, p. 237-243.
('Prehistoric remains from the shell midden of Binjai-Tamiang in North Sumatra'. According to Dr. Josef Wastl from Vienna, the human remains at Binjai Tamiang shell mound in N Sumatra are part of a Papuan/ Melanesoid population (Widianto & Noerwidi (2023) concur with australo-melanesoid interpretation)

Watanabe, N. & D. Kadar (eds.) (1985)- Quaternary geology of the hominid fossil bearing formations in Java. Geological Research Development Centre (GRDC), Bandung, Special Publ. 4, p. 1-378.
(Extensive report of multi-year Indonesian- Japanese Research Project 1976-1979, mainly on Pleistocene of Sangiran area, C Java. Project did not find new hominid fossils, but good documentation of Sangiran stratigraphy, faunas, radiometric and paleomagnetic studies, etc.)

Weesie, P.D.M. (1982)- The fossil bird remains in the Dubois collection. In: G. Batstra & W. Casparie (eds.) Modern Quaternary Research in Southeast Asia 7, Balkema, Rotterdam, p. 87-90.
(Pleistocene bird fossils from Trinil, C Java, incl. Branta, Ephippiorhynchus, Leptoptilos, Pavo, Tadorna. Assemblage include species now restricted to more northern regions, suggests cooler climate than present-day during deposition of Trinil beds)

Weidenreich, F. (1942)- Early man in Indonesia. The Far Eastern Quarterly 2, 1, p. 58-65.
(Early review of Pleistocene hominids in Indonesia)

Weidenreich, F. (1945)- The puzzle of *Pithecanthropus*. In: P. Honig & F. Verdoorn (eds.) Science and scientists in the Netherlands Indies, Board for the Netherlands Indies, Surinam and Curacao, New York, p. 380-390.
(online at: https://archive.org/details/in.ernet.dli.2015.183759)
(Review of research works on 'Java Man' by Dubois, Von Koenigswald, etc.)

Weidenreich, F. (1945)- Giant early man from Java and South China. American Museum Natural History Anthropological Papers 40, 1, p. 5-134.
(online at: digitallibrary.amnh.org/handle/2246/263)
(Discussion of unusually large hominids (Pithecanthropus robustus Weidenreich from China, Meganthropus palaeojavanicus from Sangiran, Java). Partly based on unpublished material from Von Koenigswald)

Weidenreich, F. (1951)- Morphology of Solo man. Anthropological Papers American Museum Natural History 43, 3, p. 203-288.
(online at: http://digitallibrary.amnh.org/handle/2246/297)
(Description of 11 skulls of Homo soloensis, collected between 1931 and 1941 from Late Pleistocene Solo river 20m terrace at Ngandong, left bank of Solo River, 6 miles N of Ngawi, C Java. Associated with rich mammalian fauna (Sus terhaari, Cervus javanicus) and possible bone tools and stone balls. Hominins believed to be younger than those from other parts of Java (Sangiran, Mojokerto, Trinil). Remarkable absence of other hominin body parts. With introduction by Von Koenigswald on geology and associated fauna (NB: unfinished study due to Weidenreich's death in 1948; see also Santa Luca 1980))

Weinand, D.C. (2005)- A reeevaluation of the paleoenvironmental reconstructions associated with *Homo erectus* from Java, Indonesia, based on the functional morphology of fossil bovid Astragali. Ph.D. Thesis University of Tennessee, p. 1-173.
(online at: https://trace.tennessee.edu/utk_graddiss/2315)
(Study of bovid astragali ('cow knuckle bones') from hominid-bearing sites in Java indicate paleoenvironment at Trinil site (~1 Ma) dominated by vegetated river valleys and upland forests, broken by grasslands. Grasslands probably expanded at time of Kedung Brubus locality (~0.8 Ma). Environmental change and immigration of new species important to evolutionary success of Homo erectus in M Pleistocene)

Westaway, K.E. (2006)- Reconstructing the Quatermary landscape evolution of Western Flores: an environmental and chronological context for an archaeological site. Ph.D. Thesis University of Wollongong, p. 1-411.
(online at: http://ro.uow.edu.au/theses/562/)
(Age range for occupation of Liang Bua cave by Homo floresiensis 95-11 ka, most intensive phases of occupation 74-61 and 17-11 ka, depositional age of holotype skeleton 36-14 ka, and age of oldest human skeletal remains found on Flores 95-74 ka)

Westaway, K.E. & C.P. Groves (2009)- The mark of ancient Java is on none of them. Archaeology in Oceania 44, 2, p. 84-95.
(Suggested links between Javanese E Pleistocene Homo erectus and Australian Late Pleistocene Homo sapiens crania (Thorne, etc.) questionable. Hybridization of two species unlikely: no chronological overlap and phylogenetic analysis indicate no close genetic relationship between Ngandong-like population from Java and late Pleistocene Australian fossils from Willandra Lakes)

Westaway, K.E., J. Louys, R. Due Awe, M.J. Morwood, G J. Price, J.X. Zhao, M. Aubert, R. Joannes-Boyau, T.M. Smith et al. (2017)- An early modern human presence in Sumatra 73,000-63,000 years ago. Nature 548, 7667, p. 322-325. *(online at: www.academia.edu/39125826/An_early_modern_human_presence_in_Sumatra_73_000_63_000_years_ago)*
(Reinvestgation of Lida Ajer cave in Padang Highlands, W Sumatra, which yielded Late Pleistocene human teeth (Dubois 1890), associated with rich rainforest fauna. Enamel-dentine junction morphology, enamel thickness and comparative morphology show teeth belong to anatomically modern humans (Homo sapiens). Dating of bone-bearing sediments and U-series/ electron spin resonance dating of Pongo mammalian teeth place modern humans in Sumatra between 73-63 ka. Evidence of rainforest occupation by H. sapiens at ~70 ka is ~20 ka earlier than assumed timing of dispersal of modern humans across SE Asia)

Westaway, K.E., J.M. Morwood, R.G. Roberts, A.D. Rokus, J.X. Zhao, P. Storm, F. Aziz, G. van den Bergh, P. Hadi, Jatmiko & J. de Vos (2007)- Age and biostratigraphic significance of the Punung rainforest fauna, East Java, Indonesia, and implications for *Pongo* and *Homo*. J. Human Evolution 53, p. 709-717.
(Punung Fauna of Java represents faunal turnover when Stegodon and other archaic mammal species were replaced by modern fauna, including rainforest-dependent species such as Pongo pygmaeus (orangutan). Dated as early Last Interglacial age (between 128±15 and 118±3 ka))

Westaway, K.E., M.J. Morwood, R.G. Roberts, J.X. Zhao, T. Sutikna et al. (2007)- Establishing the time of initial human occupation of Liang Bua, western Flores, Indonesia. In: R. Grun & R.G. Roberts (eds.) 11th Int. Conf. Luminescence and electron spin resonance dating (LED 2005), Quaternary Geochronology 2, p. 337-343.
(Liang Bua limestone cave in W Flores existed as subterranean chamber since 400 ka and has archeological-faunal sequence known to span last 95 ka (type locality for small-bodied hominid Homo floresiensis))

Westaway, K.E., M.J. Morwood, T. Sutikna, M.W. Moore, A.D. Rokus, G.D. van den Bergh, R.G. Roberts & E.W. Saptomo (2009)- *Homo floresiensis* and the late Pleistocene environments of eastern Indonesia: defining the nature of the relationship. Quaternary Science Reviews 28, p. 2897-2912.
(online at: www.academia.edu/466213/Homo_floresiensis_and_the_late_Pleistocene_environments_of Etc.)
(Occupation deposits in Liang Bua limestone cave on Flores span ~95 kyrs and contain abundant stone artefacts, well preserved faunal remains and evidence for endemic hominin Homo floresiensis. H. floresiensis endured rapidly fluctuating environmental conditions over last ~100 ka. Peaks in occupation at 100-95, 74-61 and 18-17 ka correlate with episodes of channel formation and erosion in cave, which in turn correspond with high rainfall, thick soils and high bio-productivity outside)

Westaway, K.E., R.G. Roberts, T. Sutikna, M.J. Morwood, R. Drysdale, J.X. Zhao & A.R. Chivas (2009)- The evolving landscape and climate of western Flores: an environmental context for the archaeological site of Liang Bua. J. Human Evolution 57, 5, p. 450-464.
(Reconstruction of landscape history of Liang Bua cave and Wae Racang valley, W Flores, using speleothems, geological structure, tectonic uplift, karst, cave, and terrace development)

Westaway, K.E., J.X. Zhao, R.G. Roberts, A.R. Chivas, M.J. Morwood & T. Sutikna (2007)- Initial speleothem results from western Flores and eastern Java, Indonesia: were climate changes from 47 to 5 ka responsible for the extinction of *Homo floresiensis*? J. Quaternary Science 22, 5, p. 429-438.
(O and C isotopic shifts in stalagmites from W Flores and E Java suggest rapid increase in rainfall around 13 ka or 17-16.5 ka, and may be related to abrupt disappearance of Stegodon and Homo floresiensis in W Flores)

Westaway, M.C. (2002)- Preliminary observations on the taphonomic processes at Ngandong and some implications for a late *Homo erectus* survivor model. Tempus 7, p. 189-193.
(At Late Pleistocene Ngandong site (Solo River, Java) different taphonomic alterations between human and non-human skeletal elements. Homo erectus remains may be older than many non-hominin fossils)

Westaway, M.C., A.C. Durband, C.P. Groves & M. Collard (2015)- Mandibular evidence supports *Homo floresiensis* as a distinct species. Proc. National Academy of Sciences USA (PNAS) 112, 7, p. E604-E605.
(online at: www.pnas.org/content/112/7/E604.full.pdf)
(Mandibular characteristics of Homo floresiensis from Liang Bua, Flores, close to early hominins, and not pathological H. sapiens as suggested in Henneberg et al. and Eckhardt et al. 2014 papers)

Westaway, M.C., A. Durband & D. Lambert (2015)- Human evolution in Sunda and Sahul and the continuing contributions of Professor Colin Groves. In: A.M. Berle & M.F. Oxenham (eds.) Taxonomic tapestries: the threads of evolutionary, behavioural and conservation research, Australian National University (ANU) Press, p. 249-276.
(online at: www.jstor.org/stable/pdf/j.ctt169wd9c.16.pdf)
('Late' Homo erectus from sites like Ngandong/ Sambungmacan/ Ngawi, Java, often viewed as advanced H. erectus or sometimes as 'archaic' H. sapiens, but may be separate species Homo soloensis. Etc.)

Westaway, M., T. Jacob, F. Aziz, H. Otsuka & H. Baba (2003)- Faunal taphonomy and biostratigraphy at Ngandong, Java, Indonesia and its implications for the late survival of *Homo erectus*. American J. Physical Anthropology 120, Abstract, p. 222–223.

Weston, E.M. & A.M. Lister (2009)- Insular dwarfism in hippos and a model for brain size reduction in *Homo floresiensis*. Nature 459, 7243, p. 85-88.

Wetmore, A. (1940)- Avian remains from the Pleistocene of Central Java. Journal of Paleontology 14, p. 447-450.
(Pleistocene bird remains from Ngandong Terrace of Solo River at Watoealang, 5km W of Ngawi, incl. Leptoptilos titan (marabou stork) and humerus of crane (Grus grus), indicating rel. cool temperatures in U Pleistocene (cranes today not found S of China))

Wibowo, U.P. (2016)- Walking with Indonesian elephants: attribution of isolated proboscidean femurs and tibias to genus based on morphological differences. Masters Thesis, School of Earth and Environmental Sciences, University of Wollongong, p. 1-185.
(online at: https://ro.uow.edu.au/theses/4803/)
(Pleistocene- Recent deposits of Java, Sulawesi, Flores, Timor and Borneo contain elephantoid fossils, mainly of genera Sinomastodon, Stegodon and Elephas. In Late Pleistocene- Recent only Elephas maximus (Sumatra))

Wibowo, U.P., A. Ferdianto, N. Laili, D. Yurnaldi & R. Setiawan (2019)- Vestige of the hominid at the Pleistocene ancient estuarine fossils bearing site of Cisaar Valley, Sumedang, West Java. Purbawidya, J. Penelitian Pengembangan Arkeologi 8, 2, p. 65-78.
(online at: https://purbawidya.kemdikbud.go.id/index.php/jurnal/article/view/P8%282%292019-1)
(Cisaar Valley outcrops of Plio-Pleistocene sediments of Kaliwangu and Citalang Fms. Paleoenvironments change upward from shallow marine, estuarine to fluvial. Two horizons with vertebrate faunas:(1) Cisaat fauna (>1 Ma old) and (2) Trinil fauna, associated with stone artifacts (~1-0.9 Ma))

Wibowo, U.P., B. Hakim & A.M. Saiful (2019)- New find of *Stegodon sompoensis* maxilla from Ciangkange Soppeng, South Sulawesi. Walennae (J. Arkeologi Sulawesi Selatan dan Tenggara) 17, 2, p. 143-154.
(online at: https://walennae.kemdikbud.go.id/index.php/walennae/article/view/366/342)
(New maxilla fragment with molar root teeth M1 of archaic elephantoid Stegodon sompoensis from Walanae Fm conglomeratic sandstone at Cangkange, 4 km E of Cabenge Archeological site of S Sulawesi. Likely age ~2.5 Ma (Late Pliocene to E Pleistocene), based on vertebrate fauna biostratigraphy of S Sulawesi)

Wibowo, U.P., E. Setiyabudi & I. Kurniawan (2018)- A *Stegodon* mandible from Cipanaruban, Subang, West Java; description and its position in the Java vertebrate biostratigraphy. Jurnal Geologi dan Sumberdaya Mineral (JGSM) 19, 1, p. 9-14.
(online at: http://jgsm.geologi.esdm.go.id/index.php/JGSM/article/view/126/333)
(Mandible of Stegodon trigonocephalus at Cipanaruban River near Pasir Cabe paleontological site (Von Koenigswald 1935), ~6 km E of Subang, W Java. Presumably part of Cisaat Fauna, E Pleistocene)

Wibowo, U.P., E. Setiyabudi, I. Kurniawan & H. Insani (2015)- Indonesian Archipelago paleogeography as the natural laboratory of the Proboscidean migration and adaptation pattern. Proc. Joint Convention HAGI-IAGI-IAFMI-IATMI, Balikpapan, JCB2015-355, p. 1-4.
(online at: www.researchgate.net/publication/285482789_Indonesian_Archipelago_Paleogeography_as_the_Natural_Laboratory_of_the_Proboscidean_Migration_and_Adaptation_Pattern)
(History of migration elephantoids (Sinomastodon, Stegoloxodon, Stegodon, Elephas) in Pleistocene- Holocene from mainland Asia to Indonesian Archipelago (mainly Java, also Sulawesi, Flores, Timor, etc.). Various degrees of dwarfing, depending on geological conditions, i.e. in island faunas)

Wibowo, U.P., I.Y.P. Suharyogi & E. Setiyabudi (2017)- The enigma of the existence of vertebrate fossils in the Flores Island. Proc. Joint Convention HAGI-IAGI-IAFMI-IATMI (JCM 2017), Malang, p. 1-3. *(Extended Abstract)*
(online at: https://www.iagi.or.id/web/digital/5/2017_IAGI_Malang_The-Enigma-of-the-Existence.pdf)
(Fossil analyses indicate Pleistocene vertebrate faunas of Flores most likely originated from Java, not Sulawesi (characteristics of Stegodon, giant tortoise and hominids that look like descendants of Homo erectus))

Widianto, H. (1986)- Posisi stratigrafi dan teknologi alat serpih Sangiran. Berkala Arkeologi 7, 1, p. 1-14.
(online at: https://berkalaarkeologi.kemdikbud.go.id/index.php/berkalaarkeologi/article/view/447)
('The stratigraphic position and technology of the Sangiran stone tools')

Widianto, H. (1993)- Unite et diversite des hominides fossils de Java: presentation de restes humains fossils inedits. Doct. Thesis, Museum National Histoire Naturelle, Institut Paleontologie Humaine, Paris, p. 1-277. *(Unpublished)*
('Unity and diversity of Java fossil hominids: presentation of unpublished human fossil remains'. Cranial remains show three Homo erectus groups: robust group, Trinil-Sangiran and Ngandong. Mandibular remains show only one genus, Homo, and one species, erectus, with two subspecies defined here: Homo erectus paleojavanicus and Homo erectus javanicus)

Widianto H. (2001)- Searching for *Homo erectus* artifacts. In: E. Indriati (ed.) A scientific life: papers in honor of Prof. Dr. T. Jacob, Bigraf Publishing, Yogyakarta, p. 75-89.

Widianto, H. (2001)- The perspective on the evolution of Javanese *Homo erectus* based on morphological and stratigraphic characteristics. In: T. Simanjuntak et al. (eds.) Sangiran: man, culture, and environment in Pleistocene times, Proc. International Colloquium on Sangiran, Solo- Indonesia, Solo 1998, Jakarta, National Research Centre of Archaeology, Yayasan Obor Indonesia, Jakarta, p. 24-45.

Widianto, H. (2011)- Human path after Sangiran era. Conservation office Sangiran Early Man site, p. 1-148.

Widianto, H. (2012)- Sangiran. The breadth of hominid sites. Directorate General of History and Archaeology, Conservation Office of Sangiran Early Man Site, Java, p. 1-141.

Widianto, H. & D. Grimaud-Herve (2000)- Un nouveau crane humain fossile dans le dome de Sangiran (Java, Indonesie). Comptes Rendus Academie Sciences, Paris, Ser. IIA, 330, 12, p. 883-888.
('A recently discovered fossil human skull near Grogol Wetan village in Kabuh Fm of in Sangiran dome, Java'. New human remains from Kabuh Fm at Grogol Wetan, Sangiran dome, with morphological characters very similar to other hominids of same horizon. Part of homogeneous population of asiatic Homo erectus)

Widianto, H. & D. Grimaud-Herve (2014)- The latest discoveries of *Homo erectus* in Java. In: N. Amano et al. (eds.) Southeast Asia: human evolution, dispersals and adaptations, 17th Congress International Union Prehistoric and Protohistoric Sciences (UISPP), Burgos 2014, p. 102-106.

Widianto, H., D. Grimaud & S. Sartono (2001)- The evolutionary position of the Ngawi calvaria. Bull. Indo-Pacific Prehistory Association (IPPA) 21, p. 162-169.
(online at: https://journals.lib.washington.edu/index.php/BIPPA/article/view/11778/10407)
(Ngawi 1 hominid skull originally described by Sartono 1991 from left bank Solo River near Selopuro, possibly derived from fluvial Pitu terraces, 5 km W of Ngawi. Probably member of Ngandong and Sambungmacan group of M-U Pleistocene Homo erectus)

Widianto, H.& S. Noerwidi (2020)- 'It is time to look at the West: a new interpretation on *Homo erectus* findings distribution on Java. Berkala Arkeologi 40, 2, p. 153-178.
(online at: https://berkalaarkeologi.kemdikbud.go.id/index.php/berkalaarkeologi/article/view/598/544)
(Occupation of Java by 'Asian' mammals started at Plio-Pleistocene boundary, around 2.4 Ma. Oldest Homo erectus found in Sangiran, around 1.5 Ma. Recent Pleistocene discoveries from W Java (Rancah, Semedo, Bumiayu) shows oldest Homo erectus in W Java between 1.8-1.7 Ma) (Indonesian version "Saatnya menengok ke barat: sebuah interpretasi barutentang distribusi temuan Homo erectus di Jawa" online at: https://berkalaarkeologi.kemdikbud.go.id/index.php/berkalaarkeologi/article/view/598/545))

Widianto, H.& S. Noerwidi (2023)- Long journey of Indonesian *Homo erectus*: Arrival and dispersal in Java Island. L'Anthropologie 127, 3 (Special Issue Prehistoire de l'Indonesie), 103167, p.
(Review paper. Three evolutive stages among Javanese Homo erectus during Pleistocene: archaic Homo erectus, typical Homo erectus, and progressive Homo erectus. Earlier taxonomic names, such as Pithecanthropus (erectus, robustus, dubius, modjokertensis, soloensis) and Meganthropus paleojavanicus do not indicate genus or species differences; there is only one genus (Homo) and one species (erectus) on Java. No evolutive link between last Homo erectus and first H. sapiens in Indonesia can be seen)

Widianto, H.& S. Noerwidi (2023)- The emergence and distribution of early modern human in Indonesia. L'Anthropologie 127, 3 (Special Issue Prehistoire de l'Indonesie), 103161, p. 1-25.
(Discussion of 'Multiregional Evolution" versus "Out-of-Africa" models of Homo sapiens evolution and migration. Early anatomically modern human remains in Indonesia found since second half of U Pleistocene, including Wajak man (age uncertain), Lida Ajer (W Sumatra) and Punung (Java) remains, and also dwarf Homo floresiensis from Liang Bua, Flores)

Widianto, H.& S. Noerwidi (2023)- Australo-melanesian: Human population in Indonesian Archipelago during the Pleisto-Holocene transition. L'Anthropologie 127, 3 (Special Issue Prehistoire de l'Indonesie), 103157, p. 1-19.
(Since the end of the Ice Age (beginning of Holocene, 11,000 years ago), the Indonesian Archipelago has shown significant waves of Early Modern Human migration. Indonesia's current population characterized by Mongoloid group in W and Australo-Melanesian race in E, with common intermixing. Early Australo-Melanesian remains widespread between ~10,000-4,000 years ago, from N Sumatra (Bukit Kerang shell midden), Java (Gunung Sewu), Sulawesi (Toalean culture), Lesser Sunda islands (Liang Toge), etc. Many were cave dwellers. After ~4000 years ago, in W Indonesia, Australo-Melanesia poplation replacedby Mongoloids)

Widianto, H.& S. Noerwidi (2023)- Austronesian diaspora in Indonesian archipelago arrived at the last migration wave. L'Anthropologie 127, 3 (Special Issue Prehistoire de l'Indonesie), 103162, p. 1-15.
(Australo-Melanesian humans dominated in Indonesian Archipelago in first half of Holocene, as revealed by discoveries in Sumatra, Java, and S Kalimantan. Provisional dating data indicates this population has lived in region since around 13,000 (Simanjuntak, 2002) to be slowly replaced by new population of Austronesians around 4000 years ago)

Widianto, H.& S. Noerwidi (2023)- Long journey of Indonesian *Homo erectus*: Arrival and dispersal in Java Island. L'Anthropologie 127, 3 (Special Issue Prehistoire de l'Indonesie), 103167, p. 1-25.

(Review of Early to Late Pleistocene Homo erectus on Java from ~1.5 Ma to 100,000-300,000 years ago. Pithecanthropus (erectus, robustus, dubius, modjokertensis, soloensis) and Meganthropus paleojavanicus all viewed here as one species Homo erectus)

Widianto, H., S. Noerwidi & A.T. Hascarya (2023)- New hominin calvaria discovery from Grenzbank layer of Sangiran Dome (Java, Indonesia): The last archaic *Homo erectus* lived in Java. L'Anthropologie 127, 3 (Special Issue Prehistoire de l'Indonesie), 103165, p. 1-20.
(New find in 2016 by locals of incomplete skullcap in Bokong, Sangiran, believed to be from Grenzbank conglomerate horizon. Bjg 1602 similar morphology and size as robust Sangiran 4 skullcap (first known as Pithecanthropus robustus), which is one of the robust Homo erectus members from E Pleistocene between 1.6-1.2 Ma. Bjg 1602 may represent last survivor of robust Homo erectus in Java in E-M Pleistocene (~0.9 Ma)).

Widianto, H. & T. Simanjuntak (2010)- Sangiran answering the world. Conservation office Sangiran Early Man site, p. 1-146.

Widianto, H., B. Toha & T. Simanjuntak (2001)- The discovery of stone implements in the Grenzbank: new insights into the chronology of the Sangiran flake industry. Bull. Indo-Pacific Prehistory Association 21, p. 157-169.
(online at: http://ejournal.anu.edu.au/index.php/bippa/article/view/273/263)
(Sangiran flake industry stone tools made from chalcedony and silicified tuff found in situ in 'Grenzbank' layer between Kabuh and Pucangan Fms. Age at least 800,000 years ago)

Widianto, H., B. Toha, T. Simanjuntak & M. Hidayat (1997)- Laporan penelitian Situs Sangiran: Proses sedimentasi, posisi stratigrafi dan kronologi artefak pada endapan purba Seri Kabuh dan Seri Notopuro. Berita Penelitian Arkeologi 1, p. 1-136.
('Sangiran site research report: Sedimentation process, stratigraphic position and chronology of artifacts in the ancient deposits of the Kabuh and Notopuro series')

Widianto, H. & V. Zeitoun (2003)- Morphological description, biometry and phylogenetic position of the skull of Ngawi 1 (East Java, Indonesia). Int. J. Osteoarchaeology 13, 6, p. 339-351.
(Rel. complete and well-preserved human skull of Ngawi 1 (Solo River near Selopuro village; Sartono 1991) morphologically closer to Ngandong-Sambungmacan (40,000 yrs) than to Trinil-Sangiran series. Question is whether skull belongs to subspecies of H. sapiens, or to H. soloensis. After local volcano-tectonic events at 71 ka and catastrophic events at 780 ka, first inhabitants of Java may have disappeared and Ngawi 1 may be new invader from Asia. Indonesian human group may have evolved at same time as Neandertals in Europe)

Widiasmoro (1998)- Late Tertiary- Early Quaternary magmatic arc and its relationship to the sedimentation processes in Sangiran, Central Java. In: H.T. Simanjuntak (ed.) Proc. International Colloquium on Sangiran: man, culture and environment in Pleistocene times, Solo 1998, Japan Foundation and National Research Center for Archaeology, Jakarta, p. 45-46.

Widiasmoro & S. Boedhisampurno (2001)- Relationship between the late Tertiary- early Quaternary magmatic activity and the paleoecology of the *Pithecanthropus* in Central-East Java. In: E. Indriati (ed.) A scientific life, Papers in Honor of Prof. Dr. T. Jacob, Proc. Conference on Man: Past, Present and Future, BIGRAF Publishing, Yogyakarta, p. 119-136.

Willemsen, G.F. (1986)- *Lutrogale palaeoleptonyx* (Dubois, 1908), a fossil otter from Java in the Dubois collection. Proc. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, B 89, 2, p. 195-200.
(M Pleistocene otter fossil from fluvial deposits Kedung Brubus, C Java)

Williams, E. (1957)- *Hardella isoclina* Dubois redescribed. Zoologische Mededelingen (Leiden), 35, 17, p. 235-240.
(online at: https://repository.naturalis.nl/pub/319310)

(Re-description and first illustration of well-preserved Pleistocene emydine freshwater turtle shell from Trinil Beds at Kedoeng Panas, Java (first briefly described by E. Dubois, 1908). Paper questions relation of H. isoclina with Hardella, and prefers assignment to genus Clemmys? isoclina (Dubois) (see also Das (1997))

Wiradnyana, K., T. Setiawan & H. Forestier (2023)- Le Hoabinhien et la reconstruction de la paleogeographie du Nord de Sumatra. L'Anthropologie 127, 3 (Special Issue Prehistoire de l'Indonesie), 103149, p. 1-12.
('The Hoabinhian and the reconstruction of the paleogeography of North Sumatra'. Hoabinhian Late Pleistocene-Holocene stone tool technocomplex in N Sumatra and mainland SE Asia characterized by presence of unifacial pebbles (sumatraliths). In N Sumatra Hoabinhian sites present as shell-middens in coastal areas and in caves in mountainous areas. Recent 14C dating of several Hoabihian sites in N Sumatra show occupation between 12,000- 3,000 years BP)

Witkamp, H. (1920)- Kjokkenmoddinger ter Oostkust van Sumatra. Tijdschrift Koninklijk Nederlandsch Aardrijkskundig Genootschap 37, p. 572–574.
('Shell middens at the East coast of Sumatra'. First report of presence of several large mounds of sea shells, typically 20m wide and 3-5m high, NW of Bindjai in NE Sumatra. These represent 'kitchen waste' of Mesolithic hunter-gatherer peoples. Many of these prehistoric shell mounds have since been destroyed for use as road cover and in local lime kilns. (later archeological excavation work by P. van Stein Callenfels in 1925 was never properly documented; see also Schurmann, 1931; JTvG)

Wolpoff, M.H. (1984)- Evolution in *Homo erectus*: the question of stasis. Paleobiology 10, 4, p. 389-406.
(Analyses of Homo erectus fossils, incl. Indonesian material. Shows evolutionary changes in increased cranial capacity and mandibular and dental features. Late end of H. erectus range difficult to define, as evidenced by difficulty in agreeing on whether Ngandong (C Java) samples are H. erectus or H. soloensis or H. sapiens)

Wolpoff, M.H., A.G. Thome, J. Jelinek & Y. Zhang (1994)- The case for sinking *Homo erectus*. 100 years of *Pithecanthropus* is enough! In: J.L. Franzen (ed.) 4th International Conference 100 years of *Pithecanthropus*. The *Homo erectus* problem, Frankfurt 1991, p. 341-361.
(online at: https://www.researchgate.net/publication/285136980_The_case_for_sinking_Homo_erectus_100_years_of_Pithecanthropus_is_enough)
(Provocative and probably oversimplified hypothesis that Homo erectus and Homo sapiens are parts of single evolving Homo species in past two million years)

Wurster, C.M. & M.I. Bird (2015)- Barriers and bridges: early human dispersals in equatorial SE Asia. In: J. Harff et al. (eds.) Geology and archaeology: submerged landscapes of the continental shelf, Geological Society, London, Special Publ. 411, p. 235-250.
(Review of paleogeography of W Indonesian region during Last Glacial Period. Hominin fossil sites generally associated with areas of open vegetation. N-S savannah corridor probably existed on Sunda Shelf, facilitating rapid dispersal of early humans in SE Asia)

Yabuki, H. & M. Shima (1981)- Fission track age and chemical composition of tektite from the remain of *Pithecanthropus erectus*. Scientific Papers Institute of Physical and Chemical Research (Tokyo) 75, 2, p. 102-104.
(M Pleistocene age determination of tektite from Sangiran, C. Java)

Yokoyama, Y., C. Falgueres, F. Semah, T. Jacob & R. Grun (2008)- Gamma-ray spectrometric dating of late *Homo erectus* skulls from Ngandong and Sambungmacan, Central Java, Indonesia. J. Human Evolution 55, 2, p. 274-277.
(Hominid fossils from Ngandong and Sambungmacan, C Java, Indonesia considered youngest representatives of Homo erectus (but much younger than, and different from, typical Sangiran H. erectus; JTvG). Dating of three skulls established minimum age of ~40 ka, with upper age limit ~60-70 ka. Ages suggest Homo erectus of Java likely survived Toba eruption and was possibly contemporaneous with earliest Homo sapiens in SE Asia)

Yokoyama, T., S. Hadiwisastra, W. Hantoro, T. Matsuda & S. Nishimura (1980)- K-Ar age of the 'Lahar Tuff' lowest part of the Pucangan formation, Pleistocene of Sangiran, Central Java, Indonesia. J. Riset Geologi dan Pertambangan (LIPI) 3, 1, p. 1-7.
(K-Ar age of 2.06 ± 0.6 Ma from andesite from 'Lahar Tuff' of base of Pucangan Fm in W Sangiran Dome. Layer has normal magnetic polarity, maybe correlated with Olduvai Event (1.67-1.87 Ma))

Yokoyama, T. & I. Koizumi (1989)- Marine transgressions on the Pleistocene Pecangan Formation in the Sangiran area, central Java, Indonesia. Palaeogeogr. Palaeoclim. Palaeoecology 72, p. 177-193.
(Paleomagnetic survey, diatoms and electric conductivity suggest four marine transgressions in Pleistocene Pucangan Fm between 1.8- 0.73 Ma, reflecting glacial eustasy. First major transgression at ~1.5 Ma, after the Olduvai Event and next one just before the Jaramillo Event, presumably reflecting glacial eustasy)

Yudha, D.S. (2008)- Reevaluation du crane Ngawi 1 (*Homo erectus*, Java, Indonesie), apports de l'imagerie 3D et des analyses multivariees. Master Thesis Quaternaire et Prehistoire, Museum National Histoire Naturelle, Paris, p. 1-55.
(online at: http://hopsea.mnhn.fr/pc/thesis/M2_DONAN_S_Y.pdf)
('Reevaluation of the Ngawi 1 skull (Homo erectus, Java, Indonesia); 3D imaging and multivariate analyses'. M-L Pleistocene Ngawi 1 skull found in 1987 morphologically close to 'late Homo erectus' Ngandong and Sambungmacan hominids. Morphological characteristics of Ngawi skull not directly comparable to Chinese (Sinanthropus) and African Homo erectus (H. ergaster), but one African individual (Olduvai 9) fits well in Ngandong-Ngawi group)

Yurnaldi, D., R. Setiawan & E.Y. Patriani (2018)- The magnetostratigraphy and the age of So'a Basin fossil-bearing sequence, Flores, Indonesia. Indonesian J. on Geoscience (IJOG) 5, 3, p. 221-234.
(online at: https://ijog.geologi.esdm.go.id/index.php/IJOG/article/view/433/269)
(Three fossil-bearing intervals recognized in Pleistocene Soa Basin of Flores, with upper one holding important hominin fossils. Paleomagnetic sampling of four sections shows two reverse and two normal polarity zones. Reverse magnetozones part of Matuyama Chron, normal magnetozones are Jaramillo and Brunhes chrons. Tangi Talo fossil layer below base Jamarillo, older than previously thought (>1.07 Ma). Mata Menge fossil intervals above Matuyama-Brunhes boundary (<0.78 Ma))

Yuwono, J.S.E. (2009)- Late Pleistocene to Mid-Holocene coastal and inland interaction in the Gunung Sewu karst area, Yogyakarta. Bull. Indo-Pacific Prehistory Association (IPPA) 29 (Manila Congress), p. 33-44.
(online at: https://journals.lib.washington.edu/index.php/BIPPA/article/view/9475)
(On archeological rock shelters in Gunung Sewu (Southern Mountains of Java) karst and prehistoric settlement)

Zaim, Y. (1981)- Umur serta stratigrafi Formasi Pucangan daerah Perning, Mojokerto, Jawa Timur. Proc. 10th Annual Conv. Indonesian Association Geologists (IAGI), Bandung, p. 230-237.
(online at: https://www.iagi.or.id/web/digital/34/PIT-IAGI-1981-Vol-1-Paper-21.pdf
('Revision of age and stratigraphy of the Pucangan Fm in the Perning area, Mojokerto, E Java'. Top Pucangen Fm below mammal-bearing Kabuh Fm with Late Pliocene Globorotalia tosaensis (N21))

Zaim, Y. (1989)- Les formations 'volcano-sedimentaires' Quaternaires de la region de Patiayam (Central Java, Indonesie). Milieu de sedimentation et mineralogie. Doct. Thesis Museum National Histoire Naturelle, Universite de Paris, p. 1-264. *(Unpublished)*
(online at: https://horizon.documentation.ird.fr/exl-doc/pleins_textes/2022-07/010039879.pdf)
('The Quaternary 'volcano-sedimentary' formations of the Patiayam region (C Java, Indonesia); depositional environment and mineralogy'. Young volcanoclastics of Patiayam anticlinal dome in NE Java backarc area, ~20km SSE of Muria volcano. Shoshonitic volcanism ('lower laharic breccias') started at ~ 2 Ma, likely derived from Patiayam volcano and possibly also from Bawean/ Karimunjawa area to NE. Around 1.1- 0.9 Ma onset of Muria I or II-derived leucite-bearing potassic-ultrapotassic volcanoclastics. Underlain by Pliocene marine Jambe Fm mollusc-bearing claystones and associated with Pleistocene 'Trinil-type' mammal fossils (Van Es, 1931), with 1978 discovery of hominid molar identified as Homo erectus ngandongensis. Etc.)

Zaim, Y. (1996)- The age of Pitu Terrace of Watualang region, Ngawi (East Java), Indonesia. Buletin Geologi (Jurusan Geologi Teknik ITB) 26, 2/3, p. 31-36. *(online at: www.academia.edu/73141839/The_Age_of_Pitu_Terrace_of_Watualang_region_Ngawi_East_Java_Indonesia)*
(Six Solo River terraces in area of Watualang (Ngawi). New vertebrate fossils in Pitu Terrace include Buffelus bubalus var sondaicus fosilis (Asian water buffalo) in Terrace V, suggesting age not older than Late Pleistocene, equivalent of Ngandong fauna)

Zaim, Y. (2004)- A new discovery of *Stegodon* in Early Pleistocene sediments from the Sumedang area (West Jawa, Indonesia). 18th International Senckenberg Conference, Weimar 2004, 1p. *(Abstract only)*
(online at: www.senckenberg.de/fis/sngconf18/doc/abstracts/115_Zaim.pdf)
(Summary of Zaim 2002 paper. Dwarf Stegodon tooth from E Pleistocene of W Java, probaby of Satir or Ci Saat fauna indicates E Pleistocene island in this part of W Java)

Zaim, Y. (2010)- Geological evidence for the earliest appearance of hominins in Indonesia. In: J.G. Fleagle et al. (eds.) Out of Africa 1, The first hominin colonization of Eurasia, Vertebrate Paleobiology and Paleoanthropology 2, Springer Science, p. 97-110.
(online at: https://www.researchgate.net/publication/282854129_Geological_Evidence_for_the_Earliest_Appearance_of_Hominins_in_Indonesia_Yahdi_ZaiminOut_of_Africa_I_-_The_First_Hominin_Coloniz Etc.)
(Until end Tertiary most Indonesian regions still in marine environment. Tectonics and glacioeustatic changes during Pleistocene formed Indonesian Archipelago. Sunda Land acted as land bridge and migration route for Homo erectus and vertebrate faunas from Asia mainland to Java. First arrival of vertebrate faunas from Asia to Indonesia through Sundaland at end of Late Pliocene, followed by arrival of early hominin (Homo erectus paleojavanicus (= Meganthropus paleojavanicus)) to Java in Early Pleistocene (1.6-1.0 Ma))

Zaim, Y. & R. Ardan (1998)- A premolar of *Homo erectus* from Patiayam region, Central Java. Buletin Geologi (ITB) 28, p. 31-36.
(First lower permanent premolar of Homo erectus, in fluvial sandstones of E-M Pleistocene Slumprit Fm)

Zaim, Y., R.L. Ciochon, J. Polanski, F.E. Grine, E.A. Bettis, Y. Rizal, R. Larick, M. Heizler, Aswan, K.L. Eaves & H.E. Marsh (2011)- New 1.5 million-year-old *Homo erectus* maxilla from Sangiran (Central Java, Indonesia). J. Human Evolution 61, 4, p. 363-376.
(*online at: https://www.academia.edu/5900201/New_1_5_million_year_old_Homo_erectus_maxilla_from_Sangiran_Central_Java_Indonesia)*
(New H. erectus left maxilla fragment from cemented gravelly sands at base Grenzbank Zone at Bapang, Sangiran. Pumice hornblende 2m above locality with 40Ar/39Ar age of 1.51 Ma (N.B.: numerical ages proposed in this paper for Sangiran stratigraphy are significantly older than most other dating results and not in agreement with paleomagnetic interpretations, tektite horizon, etc.?; JTvG))

Zaim, Y. & M. Delaune (1990)- Nouvelles donnees sur la stratigraphie et le milieu de sedimentation des formations volcano-sedimentaires quaternaires de la region de Patiayam (Java- Indonesie). Geodynamique 5, 2, p. 135-150.
('New data on the stratigraphy and depositional environment of the volcano-sedimentary formations of the Patiayam region (Java, Indonesia)'. Shoshonitic volcanic activity of Patiayam Dome, S of Muria Volcano in N Java, initiated ~2 Ma ago. During Lower Pleistocene (0.9 Ma) still active and contemporaneous with beginning of activity at Muria 1)

Zaim, Y., J. de Vos, O.F. Huffman, F. Aziz, J. Kappelman & Y. Rizal (2003)- A new antler specimen from the 1936 Perning hominid site, East Jawa, Indonesia, attributable to *Axis lydekkeri* (Martin, 1886). Jurnal Teknologi Mineral, Bandung, 10, 2, p. 45-52.
(Nearly complete left antler, attributed to Axis lydekkeri, found in 2001 in excavation E of relocated site that produced Homo modjokertensis in 1936. Not reported previously from hominid-bearing bed)

Zaim, Y., A.T. Hascaryo, M.R. Puspaningrum, Aswan & Y. Rizal (2022)- Many ways for Quaternary hominin and fauna to reach Indonesian Archipelago. Proc. Workshop on Ecological drivers of Pleistocene hominin and faunal dispersal across Southeast Asia (EPHSEA 2022), Bangkok, p. 20. *(Abstract only)*
(online at: https://drive.google.com/drive/folders/1Xgtqkk-R6c: teD6tBnjFniH6QiERN7PcI)
(Two likely pathways for migration of Quaternary hominins and vertebrates from Asia mainland to SE Asia and Indonesian Archipelago:(1) Siva-Sino Malayan routes of Malay Peninsula- Sumatra- Bangka- Java and Kalimantan in W, and (2) track of Philippines- Sulawesi- Nusa Tenggara in E Indonesia, and vice versa)

Zaim, Y., R. Larick, R.L. Ciochon, Suminto, Y. Rizal & Sujatmiko (1999)- Karakteristik satuan Lahar Bawah dari formasi Pucangan di Sangiran, Jawa Tengah. Buletin Geologi. (ITB) 31, p. 67-84.
('Characteristics of the Lower Lahar unit in the Pucangan Fm in Sangiran'. See also Bettis et al. 2004)

Zaim, Y. & R. Marino (2002)- Pygmy *Stegodon* dari Desa Cariang, Kecamatan Tomo, Kab. Sumedang, Jawa Barat. Buletin Geologi (ITB) 34, 1, p. 45-52.
('Pygmy Stegodon from Cariang Village, Tomo District, Sumedang, West Java'. New species of small elephantoid Stegodon cariangensis in E Pleistocene lacustrine black clay in Majalengka, W Java (deposits unconformable above Late Pliocene marine clays of Kaliwangu Fm))

Zaim, Y., Y. Rizal & Aswan (2007)- The geological background of hominid colonization of Java. In: A.M. Semah & K. Setiagama (eds.) Proc. International Conference First islanders- Human origins patrimony in Southeast Asia, AsiaLink-HOPSea Programme, Paris, p. 92-98.
(online at: http://hopsea.mnhn.fr/pc/brochures/2007HOPseaFI.pdf)

Zaim, Y., Y. Rizal, Aswan & B.S. Fitriyana (2006)- S. Sartono: dari hominid ke delapsi dengan kontroversi. Penerbit Institut Teknologi Bandung, Bandung, p. 1-183.
('S. Sartono: from hominids to 'delapsi' with controversy'. Collection of papers)

Zanolli, C. (2013)- Additional evidence for morpho-dimensional tooth crown variation in a new Indonesian *H. erectus* sample from the Sangiran Dome (Central Java). PloS One 8, 7, e67233, p. 1-15.
(online at: http://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0067233&type=printable)
(Fifteen new Homo erectus fossil dental remains found in last two decades in Kabuh Fm of Sangiran Dome area, some from excavation of human occupation floors in basal Kabuh Fm)

Zanolli, C. (2015)- Molar crown inner structural organization in Javanese *Homo erectus*. American J. Physical Anthropology 156, 1, p. 148-157.
(Study of 7 Homo erectus permanent molar crowns from late E- early M Pleistocene Kabuh Fm of Sangiran, C Java. Features differ from penecontemporaneous African H. erectus/ergaster and H. heidelbergensis, as well as in Neanderthals, but occur in recent human populations)

Zanolli, C., L. Bondioli, L. Mancini, A. Mazurier, H. Widianto, R. Macchiarelli (2012)- Brief communication: Two human fossil deciduous molars from the Sangiran dome (Java, Indonesia): outer and inner morphology. American J. Physical Anthropology 147, 3, p. 472-481.
(Two deciduous crowns collected near Pucung, probably from E-M Pleistocene Kabuh Fm, or from E Pleistocene 'Grenzbank'. Probably from Homo erectus)

Zanolli, C., F.E. Grine, O, Kullmer, F, Schrenk & R, Macchiarelli (2015)- The early Pleistocene deciduous hominid molar FS-72 from the Sangiran dome of Java, Indonesia: A taxonomic reappraisal based on its comparative endostructural characterization. American J. Physical Anthropology 157, 4, p. 666-674.
(Among 10 fossil hominid deciduous teeth reported from Pleistocene of Sangiran Dome of Java are two isolated lower second molars: PCG.2 from Kabuh Fm and FS-72 from Pucangan Fm. FS-72 originally attributed to Meganthropus paleojavanicus because of large size, but internal morphology suggests more likely a dm2 of Pongo rather than Homo)

Zanolli, C., O. Kullmer, J. Kelley, A.M. Bacon, F. Demeter, J. Dumoncel, L. Fiorenza, F. Grine, J.J. Hublin et al. (2019)- Evidence for increased hominid diversity in the Early to Middle Pleistocene of Indonesia. Nature Ecology Evolution 3, 5, p. 755-764.
(Manuscript online at: https://hal.science/hal-02296699/file/Zanolli_%26_al_2019_NatureEcolEvol.pdf)
(Since first discovery of Pithecanthropus (Homo) erectus by E. Dubois at Trinil in 1891, over 200 hominid dentognathic remains collected from E-M Pleistocene deposits of Java. Dental structures confirm presence of Meganthropus as Pleistocene hominid distinct from Pongo, Gigantopithecus and Homo, and also reveal that Dubois's H. erectus paratype molars from 1891 are not hominin but Meganthropus)

Zeitoun, V. (2000)- Revision de l'espece *Homo erectus* (Dubois, 1893). Bulletins Memoires Societe Anthropologie de Paris, N.S., 12, 1-2, p. 1-200.
(online at: https://journals.openedition.org/bmsap/pdf/5963)
('Reappraisal of the species Homo erectus'. Mainly on details of skull morphology)

Zeitoun, V., V. Barriel & H. Widianto (2016)- Phylogenetic analysis of the calvaria of *Homo floresiensis*. Comptes Rendus Palevol 15, 5, p. 555-568.
(online at: www.sciencedirect.com/science/article/pii/S1631068316000130)
(Metrics of calvariae of human fossils from Liang Bua, Flores, indicate LB1 is part of Homo erectus clade, and less similar to Sambungmacan-Ngandong-Ngawi group. Homo floresiensis not pathological modern human)

Zeitoun, V., F. Detroit, D. Grimaud-Herve & H. Widianto (2010)- Solo man in question: convergent views to split Indonesian *Homo erectus* in two categories. Quaternary International 223-224, p. 281-292.
(online at: https://www.academia.edu/23025566/Solo_man_in_question_Convergent_views_to_split_Indonesian_Homo_erectus_in_two_categories)
(Homo (Javanthropus) soloensis Oppenoorth 1932 from Solo River bank terraces thought to belong to either archaic Homo sapiens, or (most paleoanthropologists:) an evolved Homo erectus. Chronological gap splits two categories, posing question of possibly two separate species. Catastrophic event around Brunhes/Matuyama geomagnetic reversal (Australasian tektite strewn field 770,000 or 803 ka) must have triggered shifting among Asian populations and environments. Toba eruption of 17 ka probably also had drastic global consequences on human ecology and evolution)

Zeitoun, V., H. Widianto & T. Djubiantono (2007)- The phylogeny of the Flores Man: the cladistic answer. In E. Indriati (ed.) Proc. International Seminar Southeast Asian paleoanthropology: Recent advances on Southeast Asian paleoanthropology and archaeology, Gadjah Mada University, Yogyakarta, p. 54-60.

Zeitoun, V., C. Winayalai, P. Auetrakulvit, R. Debruyne, J.B. Mallye & A. Lenoble (2023)- Transgressing the limits of palaeoenvironmental data for Southeast Asian Pleistocene faunal assemblages: A critical review to go further. Annales de Paleontologie 109, 4, 102657, p. 1-18.
(Critical review of Pleistocene 'savannah corridor' hypothesis for SE Asia mammal migrations)

Zhang, H., T. Pape & A.M. Lister (2028)- On the type material of *Elephas hysudrindicus* Dubois, 1908 (Mammalia, Proboscidea). J. Vertebrate Paleontology 38, 1, e1425211, p. 1-3.
(online at: https://www.researchgate.net/publication/323863868_On_the_type_material_of_Elephas_hysudrindicus_Dubois_1908_Mammalia_Proboscidea)
(In a study of a collection of elephantid molars, etc., from multiple location in Kendeng Hills of Java, Dubois (1908), saw E. hysudrindicus from Java as evolutionary intermediate between E Pleistocene E. hysudricus Falconer and Cautley1845 from Siwalik Hills of India-Pakistan and extant Asian elephant E. "indicus" (now E. maximus Linnaeus, 1758))

Zijlstra, J.S., L.W. van den Hoek Ostende & R. Awe Due (2008)- Verhoeven's giant rat of Flores (*Papagomys theodorverhoeveni*, Muridae) extinct after all? Contributions to Zoology, 77, 1, p. 25-31.
(online at: https://www.repository.naturalis.nl/document/98741)

(Giant rat species from Flores Papagomys theodorverhoeveni believed to be extinct, but reported by Suyanto and Watts (2002) as still extant, on basis of single museum specimen. Identification of that specimen disputed here)

Zwierzycki, J. (1926)- Een vondst uit de Palaeolithische cultuurperiode in een grot in Boven-Djambi. De Mijningenieur 7, 4, p. 64-67.
('A discovery from the Paleolithic culture period in a cave in Upper Jambi'. Tianko Panjang cave excavation in early 1920's yielded some animal bones (deer, pangolin) and obidian artifacts (see also Sarasin 1914, Bronson & Asmar 1975))

Zwierzycki, J. (1926)- De beteekenis van nieuwe fossiele werveldiervondsten bij Boemiajoe. De Mijningenieur 7, 12, p. 229-234.
('The significance of new fossil mammal discoveries near Bumiayu'. Localities in river valleys of Kali Glagah and Ci Saat, 'West Kendeng Zone', C Java. Ongoing excavations of mammal bones in several horizons in ~1200m thick series of coarse sandstones, conglomerates and clays with fresh-water molluscs. Fossil elephants (incl. Stegodon, Elephas), hippopotamus, etc., described by Stehlin (1925). Age believed to be Late Pliocene, possibly E Pleistocene like Trinil (lecture summary; no figures) (for more on Bumiayu locality info see Van der Maarel (1932) and Semah (1997))

Zwierzycki, J. (1948)- Przedhistoryczne typy ludzkie na Jawie. Wiadomosci Museum Ziemi 4, Warsaw, p. 137-172.
('Prehistoric types of Man on Java'. In Polish with English summary. Five vertebrate horizons in C and E Java: (1) Tjisande, (2) Tjidjoelang, (3) Lower Boemiajoe, (4) Upper Boemiajoe and (5) Trinil)



Bibliography of Indonesian Geology